

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 March 2006 (16.03.2006)

PCT

(10) International Publication Number
WO 2006/029196 A1

(51) International Patent Classification:
C12N 15/864 (2006.01)

(21) International Application Number:
PCT/US2005/031837

(22) International Filing Date:
8 September 2005 (08.09.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/607,854 8 September 2004 (08.09.2004) US

(71) Applicant (for all designated States except US): **THE GOVERNMENT OF THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES, NATIONAL INSTITUTES OF HEALTH** [US/US]; Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852-3804 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHIORINI, John, A.** [US/US]; 9611 Hillridge Drive, Kensington, MD 20895 (US). **PASQUALE, Giovanni, Di** [US/US]; 11701 Goodloe Road, Silver Spring, MD 20906 (US).

(74) Agents: **SPRATT, Gwendolyn, D. et al.**; Needle & Rosenberg, P.C., Suite 1000, 999 Peachtree Street, Atlanta, GA 30309-3915 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: **TRANSCYTOSIS OF ADENO-ASSOCIATED VIRUSES**



(57) Abstract: The present invention provides methods of transcytosing barrier epithelial cells using adeno-associated virus-4 (AAV4), adeno-associated virus-5 (AAV5), adeno-associated virus-7 (AAV7), bovine adeno-associated virus (BAAV), and vectors and particles derived therefrom. In addition, the present invention provides methods of delivering a nucleic acid across the barrier epithelia using the AAV4, AAV5, AAV7, and BAAV vectors and particles.

BEST AVAILABLE COPY

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 March 2006 (16.03.2006)

PCT

(10) International Publication Number
WO 2006/029196 A1

(51) International Patent Classification:
C12N 15/864 (2006.01)

(21) International Application Number:
PCT/US2005/031837

(22) International Filing Date:
8 September 2005 (08.09.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/607,854 8 September 2004 (08.09.2004) US

(71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES, NATIONAL INSTITUTES OF HEALTH [US/US]; Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852-3804 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CHIORINI, John, A. [US/US]; 9611 Hillridge Drive, Kensington, MD 20895 (US). PASQUALE, Giovanni, DI [US/US]; 11701 Goodloe Road, Silver Spring, MD 20906 (US).

(74) Agents: SPRATT, Gwendolyn, D. et al.; Needle & Rosenberg, P.C., Suite 1000, 999 Peachtree Street, Atlanta, GA 30309-3915 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: TRANSCYTOSIS OF ADENO-ASSOCIATED VIRUSES



(57) Abstract: The present invention provides methods of transcytosing barrier epithelial cells using adeno-associated virus-4 (AAV4), adeno-associated virus-5 (AAV5), adeno-associated virus-7 (AAV7), bovine adeno-associated virus (BAAV), and vectors and particles derived therefrom. In addition, the present invention provides methods of delivering a nucleic acid across the barrier epithelia using the AAV4, AAV5, AAV7, and BAAV vectors and particles.

WO 2006/029196 A1

WO 2006/029196 A1



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 March 2006 (16.03.2006)

PCT

(10) International Publication Number
WO 2006/029196 A1

(51) International Patent Classification:
C12N 15/864 (2006.01)

(21) International Application Number:
PCT/US2005/031837

(22) International Filing Date:
8 September 2005 (08.09.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/607,854 8 September 2004 (08.09.2004) US

(71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES, NATIONAL INSTITUTES OF HEALTH [US/US]; Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852-3804 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CHIORINI, John, A. [US/US]; 9611 Hillridge Drive, Kensington, MD 20895 (US). PASQUALE, Giovanni, Di [US/US]; 11701 Goodloe Road, Silver Spring, MD 20906 (US).

(74) Agents: SPRATT, Gwendolyn, D. et al.; Needle & Rosenberg, P.C., Suite 1000, 999 Peachtree Street, Atlanta, GA 30309-3915 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: TRANSCYTOSIS OF ADENO-ASSOCIATED VIRUSES



(57) Abstract: The present invention provides methods of transcytosing barrier epithelial cells using adeno-associated virus-4 (AAV4), adeno-associated virus-5 (AAV5), adeno-associated virus-7 (AAV7), bovine adeno-associated virus (BAAV), and vectors and particles derived therefrom. In addition, the present invention provides methods of delivering a nucleic acid across the barrier epithelia using the AAV4, AAV5, AAV7, and BAAV vectors and particles.

WO 2006/029196 A1



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5

TRANSCYTOSIS OF ADENO-ASSOCIATED VIRUSES**CROSS-REFERENCE TO RELATED APPLICATIONS**

This claims the benefit of U.S. Provisional Application No. 60/607,854, entitled
"Transcytosis of Adeno-Associated Viruses", filed September 8, 2004, by Chiorini *et al*,
10 which is herein incorporated by reference in its entirety.

BACKGROUND

The adeno-associated viruses (AAV) were originally classified according to size,
structure, and dependence upon a helper virus for replication. AAV is a member of the
Parvoviridae, a virus family characterized by a single stranded linear DNA genome and a
15 small icosahedral shaped capsid measuring about 20nm in diameter. AAV was first
described as a contaminant of tissue culture grown simian virus 15, a simian adeno virus
and was found dependent on adenovirus for measurable replication. This led to its name,
adeno-associated virus, and its classification in the genus Dependovirus. Because the
majority of AAV isolates were first identified as contaminants of laboratory stocks of
20 adenovirus, little is known about their natural tissue tropism. However *in vivo* experiments
suggest they are effective vectors for gene transfer applications. Currently eleven full-length
isolates have been cloned and their initial characterization indicates that each serotype has
unique binding/cell tropism characteristics.

Transcytosis is the transport of macromolecular cargo from one side of a cell to the
25 other within membrane-bounded carrier(s). It is a strategy used by multicellular organisms
to selectively move material between two different environments while maintaining the
distinct compositions of those environments. The ability of a pathogen to spread through a
tissue is a critical determinate of its virulence. The process of transcytosis has been reported
for a number of viruses. For example, HIV and poliovirus cross simple epithelial cells
30 without infection and are still infectious when they cross into the submucosa. Likewise, the
Epstein-Barr virus (EBV) forms a complex with mucosal immunoglobulins (IgA) that are
specific for gp350, a viral surface protein that is present in latently infected people. This
complex binds to the poly-immunoglobulin receptor at the basal surface of epithelial cells,
and is endocytosed and delivered apically without infection. To date, there is no report of
35 transcytosis by any AAV.

5

Provided herein are methods for transcytosis across barrier epithelial cells using AAV vectors. The ability of a non-pathogenic vector to transcytose barrier epithelial cells can be used to deliver genes to sub-epithelial targets. One important example includes the delivery of genes across the blood-brain-barrier without the need for direct injection into the brain. Furthermore, herein is described a method for re-directing virus that enters a cell by transcytosis to result in transduction of the cell by blocking exocytosis.

SUMMARY

In accordance with the purpose(s) of this invention, as embodied and broadly described herein, this invention, in one aspect, relates to a method of delivering a heterologous nucleic acid across an epithelial barrier comprising delivering to the epithelial barrier an AAV vector comprising the heterologous nucleic acid. The epithelial cells can be in the gut, lung, genitourinary tract, kidney, blood vessels or brain.

In another aspect, the invention relates to a method of transcytosing epithelial cells of a human subject comprising administering to the subject a viral vector comprising a heterologous nucleic acid, wherein the viral vector is selected from a group consisting of BAAV, AAV4, AAV5, or AAV7.

Additional advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the invention. The advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings, which are incorporated in and constitute a part of this specification, illustrate (one) several embodiment(s) of the invention and together with the description, serve to explain the principles of the invention.

Figure 1 shows that AAV4 transcytosed in CaCo-2, MDCKI, MDCKII, Human primary immortalized epithelial endometrial, Bovine brain primary endothelia cells (BBB). AAV5 transcytosed CaCo-2 cells, whereas BAAV transcytosed in MDCKs, Endometrial,

5 airways epithelia, and BBB. AAV6 did not transcytose in any of cell types tested. Hela cells do not form barrier epithelia and were used as a control.

Figure 2 shows that the treatment of the basal lateral surface of Human primary airways epithelial cell (HAE) with tannic acid blocked the transcytosis of BAAV vector containing a GFP expression cassette from the apical surface to the basal lateral.

10 Furthermore transduction dramatically increased when assayed at 24 hrs post inoculation. In contrast no change was observed in AAV2 transduction, which did not demonstrate any transcytosis activity and has limited binding activity on HAE.

Figure 3 shows AAV7 transcytosis assay on bovine brain endothelial cells. Virus DNA extracted from basal lateral medium after 3H incubation 2×10^9 DRP of AAV were
15 loaded on the apical side of the cell layer. AAV5 is used as a control.

DETAILED DESCRIPTION

The present invention may be understood more readily by reference to the following detailed description of the invention and the Examples included therein and to the Figures and their previous and following description.

20 Before the present compounds, compositions, articles, devices, and/or methods are disclosed and described, it is to be understood that this invention is not limited to specific synthetic methods, specific cell types, or to particular tissues, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

25 As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

Ranges may be expressed herein as from "about" one particular value, and/or to
30 "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint,
35 and independently of the other endpoint.

5 “Optional” or “optionally” as used herein means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances where it does not.

AAV Transcytosis

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier comprising delivering to the epithelial barrier an AAV vector comprising the heterologous nucleic acid. In one aspect of the method, the AAV is AAV4, AAV5, AAV7, or BAAV. The AAV capsid protein forming the viral particle is understood herein to confer upon the AAV particle the desired transcytosing ability. Thus, “AAV vector”, as used herein, refers to any virion, vector, or viral particle comprising or encoding at least one AAV capsid protein. As an example, an AAV4 vector can encode an AAV4 capsid protein and thus be encapsidated in said protein forming an AAV4 particle. Alternatively the AAV vector can comprise a nucleic acid encoding a modified AAV or a portion of an AAV capsid protein (a capsid protein fragment) that confers serotype-specific transcytotic activity. AAV capsids, capsid protein fragments and capsid modifications are disclosed, for example, in U.S. Patent Application No. 60/526786 (BAAV), U.S. Patent No. 6,468,524 (AAV4), U.S. Patent Application No. 09/717,789 (AAV5), U.S. Patent Application 2003/0228282 (AAV7), International Application No. PCT/US04/15534, filed May 19, 2004 (AAAV), and U.S. Patent Application No. 60/676604, filed April 29, 2005 (AAV-X1, AAV-X1b, AAV-X5, AAV-X19, AAV-X21, AAV-X22, AAV-X23, AAV-X24, AAV-X25, AAV-X26).

25 In another aspect of the method, the epithelial cells are in the gut, lung, genitourinary tract, kidney, blood vessels or brain. In another aspect of the method, the epithelial cells can be selected from a group consisting of bronchial, alveolar, tracheal or upper airway epithelial cells; absorptive enterocytes or M cells; endometrial or urinary epithelial cells; renal collecting duct or proximal tubule epithelial cells; cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells.

Further disclosed is a method of transcytosing epithelial cells of a human subject comprising administering to the subject an AAV vector comprising a heterologous nucleic acid. In one aspect of the method, the vector is AAV4, AAV5, AAV7, or BAAV. In another aspect of the method, the epithelial cells are selected from a group consisting of bronchial, alveolar, tracheal or upper airway epithelial cells; absorptive enterocytes or M cells;

35

5 endometrial or urinary epithelial cells; renal collecting duct or proximal tubule epithelial cells; cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells.

Further contemplated are methods for the delivery of molecules across epithelial cell barriers comprising coupling the molecules to non-recombinant (wild-type) AAV capsids or particles. In one aspect, the molecules are radioligands or enzymes.

10 The term "adeno-associated virus (AAV)" is used herein to refer to a genus of viruses in the family Parvoviridae which are all defective viruses (unable to replicate by themselves) and depend on the co-infection of their host cell by other, nondefective viruses to help them replicate.

Transcytosis refers to the transport of macromolecular cargo from one side of a cell
15 to the other, generally within a membrane-bounded carrier(s). Tuma and Hubbard provided a review of transcytosis (Tuma PL and Hubbard AL. 2003. *Physiol Rev.* 83:871-932), herein incorporated by reference for its teaching regarding the nature and uses for transcytosis. Transcytosis is a strategy used by multicellular organisms to selectively move material between two different environments while maintaining the distinct compositions of those
20 environments. N. Simionescu was the first to coin the term transcytosis to describe the vectorial transfer of macromolecular cargo within the plasmalemmal vesicles from the circulation across capillary endothelial cells to the interstitium of tissues. During this same period, another type of transcytosis was being discovered. Immunologists comparing the different types of immunoglobulins found in various secretions (e.g., serum, milk, saliva,
25 and the intestinal lumen) speculated that the form of IgA found in external secretions (called secretory IgA, due to the presence of an additional protein component) was selectively transported across the epithelial cell barrier. More is known about transcytosis as it is expressed in epithelial tissues, which form cellular barriers between two environments. In this polarized cell type, net movement of material can be in either direction, apical to
30 basolateral or the reverse, depending on the cargo and particular cellular context of the process. However, transcytosis is not restricted to only epithelial cells.

Since the 19th century dye experiments of Ehrlich, the brain has been known as a "privileged" organ where access is tightly regulated so that the environment remains chemically stable. The two principal gatekeepers of the brain are the cerebral capillary
35 endothelium and the cuboidal epithelial cells of the choroid plexus. These cellular barriers are specialized for the passage of different nutrients from the blood. The capillaries move

5 nutrients that are required rapidly and in large quantities, such as glucose and amino acids. These small molecules are transported by membrane carriers using facilitated diffusion. The choroid plexus supplies nutrients that are required less acutely and in lower quantities. These are folate and other vitamins, ascorbate, and deoxyribonucleotides.

There are two epithelial cells that participate in transcytosis in the intestine, M cells
10 and enterocytes (adsorptive columnar cells). These cells are very different from one another and the capillary endothelial cell. Depending on the species, M cells comprise a variable but small percentage of the epithelia overlying organized mucosal-associated lymphoid tissue, making them a very minor cell population in the gastrointestinal tract. The transcytotic route across M cells is thought to be part of the mechanism by which antigens are routinely
15 sampled along the entire mucosal surface. Not surprisingly, numerous pathogens have evolved mechanisms to exploit the transcytotic process as a means to invade and disseminate before a strong enough immune response can be mounted.

Absorptive enterocytes are simple columnar cells with several apical features in addition to their brush borders. Clathrin-coated pits are present at the base of microvilli, and
20 a thick glycocalyx composed of integral membrane proteins with glycosaminoglycan side chains emanates from the microvillar membrane. This latter structural feature as well as the rigidity of the microvilli are thought to prohibit microorganisms from attaching and invading enterocytes. The intracellular organization of these columnar epithelial cells is also polarized, with basally located nuclei, supranuclear Golgi, and an abundance of
25 pleiomorphic membrane compartments underlying the terminal web of the brush border. The basolateral-to-apical length of this cell is ~20 versus 0.2 μm for a capillary endothelial cell, making the transcytotic route across enterocytes potentially much longer. Furthermore, microtubules are an important structural element of the transcytotic pathway in enterocytes, but not in M or endothelial cells.

30 Transcytosis also occurs in the upper regions of the respiratory tract and has been demonstrated with two vector systems, pIgA-R and FcRn, but others could exist. Secretory IgA is a known constituent of the lung's immune defense system, with bronchial epithelial cells carrying out basolateral-to-apical transport of dIgA, which is secreted by local plasma cells in underlying lymphoid tissue. Albumin, which is found in lung fluid, is endocytosed
35 specifically at the apical surface of airway epithelia but is then subsequently degraded. At the alveolar level, the question of whether albumin is transcytosed intact is uncertain.

5 The methods and compositions described herein can be used to deliver heterologous nucleic acids to certain tissues. As used herein, the term "nucleic acid" refers to single-or multiple stranded molecules which may be DNA or RNA, or any combination thereof, including modifications to those nucleic acids. The nucleic acid may represent a coding strand or its complement, or any combination thereof. Nucleic acids may be identical in
10 sequence to the sequences which are naturally occurring for any of the novel genes discussed herein or may include alternative codons which encode the same amino acid as those provided herein, including that which is found in the naturally occurring sequence. These nucleic acids can also be modified from their typical structure. Such modifications include, but are not limited to, methylated nucleic acids, the substitution of a non-bridging
15 oxygen on the phosphate residue with either a sulfur (yielding phosphorothioate deoxynucleotides), selenium (yielding phosphorselenoate deoxynucleotides), or methyl groups (yielding methylphosphonate deoxynucleotides).

 As used herein, the term "isolated" refers to a nucleic acid separated or significantly free from at least some of the other components of the naturally occurring organism, for
20 example, the cell structural components or viral components commonly found associated with nucleic acids in the environment of the virus and/or other nucleic acids. The isolation of the native nucleic acids can be accomplished, for example, by techniques such as cell lysis followed by phenol plus chloroform extraction, followed by ethanol precipitation of the nucleic acids. The nucleic acids of this invention can be isolated from cells according to any
25 of many methods well known in the art.

 The AAV vectors disclose herein can comprise a heterologous nucleic acid functionally linked to the promoter. The term "heterologous" is used herein to refer to a nucleic acid which is derived from a different cell, tissue or organism. The nucleic acid can encode a polypeptide or protein or an antisense RNA, for example. By "functionally linked"
30 is meant such that the promoter can promote expression of the heterologous nucleic acid, as is known in the art, such as appropriate orientation of the promoter relative to the heterologous nucleic acid. Furthermore, the heterologous nucleic acid preferably has all appropriate sequences for expression of the nucleic acid, as known in the art, to functionally encode, *i.e.*, allow the nucleic acid to be expressed. The nucleic acid can include, for
35 example, expression control sequences, such as an enhancer, and necessary information processing sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites,

5 and transcriptional terminator sequences.

The heterologous nucleic acid can encode beneficial proteins that replace missing or defective proteins required by the subject into which the vector is transferred or can encode a cytotoxic polypeptide that can be directed, *e.g.*, to cancer cells or other cells whose death would be beneficial to the subject. The heterologous nucleic acid can also encode antisense
10 RNAs that can bind to, and thereby inactivate, mRNAs made by the subject that encode harmful proteins. In one embodiment, antisense polynucleotides can be produced from a heterologous expression cassette in an AAV4 viral construct where the expression cassette contains a sequence that promotes cell-type specific expression (Wirak *et al.*, 1991. *EMBO* 10:289). For general methods relating to antisense polynucleotides, see *Antisense RNA and*
15 *DNA*, D. A. Melton, Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1988).

Examples of heterologous nucleic acids which can be administered to a cell or subject as part of the present AAV4 vector can include, but are not limited to the following: nucleic acids encoding therapeutic agents, such as tumor necrosis factors (TNF), such as TNF- α ; interferons, such as interferon- α , interferon- β , and interferon- γ ; interleukins, such as
20 IL-1, IL-1 β , and ILs -2 through -14; GM-CSF; adenosine deaminase; cellular growth factors, such as lymphokines; soluble CD4; Factor VIII; Factor IX; T-cell receptors; LDL receptor; ApoE; ApoC; alpha-1 antitrypsin; ornithine transcarbamylase (OTC); cystic fibrosis transmembrane receptor (CFTR); insulin; Fc receptors for antigen binding domains of antibodies, such as immunoglobulins; and antisense sequences which inhibit viral
25 replication, such as antisense sequences which inhibit replication of hepatitis B or hepatitis non-A, non-B virus. The nucleic acid is chosen considering several factors, including the cell to be transfected. Where the target cell is a blood cell, for example, particularly useful nucleic acids to use are those which allow the blood cells to exert a therapeutic effect, such as a gene encoding a clotting factor for use in treatment of hemophilia. Furthermore, the
30 nucleic acid can encode more than one gene product, limited only, if the nucleic acid is to be packaged in a capsid, by the size of nucleic acid that can be packaged.

The term "polypeptide" as used herein refers to a polymer of amino acids and includes full-length proteins and fragments thereof. Thus, "protein," polypeptide," and "peptide" are often used interchangeably herein. Substitutions can be selected by known
35 parameters to be neutral (*see, e.g.*, Robinson WE Jr, and Mitchell WM., 1990. *AIDS* 4:S151-S162). As will be appreciated by those skilled in the art, the invention also includes

5 those polypeptides having slight variations in amino acid sequences or other properties. Such variations may arise naturally as allelic variations (*e.g.*, due to genetic polymorphism) or may be produced by human intervention (*e.g.*, by mutagenesis of cloned DNA sequences), such as induced point, deletion, insertion and substitution mutants. Minor changes in amino acid sequence are generally preferred, such as conservative amino acid
10 replacements, small internal deletions or insertions, and additions or deletions at the ends of the molecules. Substitutions may be designed based on, for example, the model of Dayhoff, *et al.* (in *Atlas of Protein Sequence and Structure 1978*, Nat'l Biomed. Res. Found., Washington, D.C.). These modifications can result in changes in the amino acid sequence, provide silent mutations, modify a restriction site, or provide other specific mutations.

15 The term "epithelia" is used herein to refer to cells which are linked tightly together by intercellular junctions to form a planar sheet. These sheets of cells form a barrier between two compartments. Epithelia therefore line all surfaces and cavities (including skin, peritoneum, linings of the intestine, airways, genitourinary tracts, glands, and blood vessels.

An epithelium has a free or apical surface facing the environment, or lumen of a
20 cavity, and a basal surface facing the underlying connective tissue. The boundary between the basal surface of an epithelium and the underlying connective tissue is usually very sharp, and is the site where the basal lamina (BL) is present. Most BL are too thin to be seen with the light microscope. However, the BL, together with a thin layer of connective tissue, is often times seen at the epithelial/connective tissue interface. This composite layer, visible
25 with the light microscope, was initially called the Basement Membrane. Application of the electron microscope revealed that, in most cases, this Basement Membrane actually consisted of the true basal lamina (lamina lucida plus lamina densa), along with a layer of adherent connective tissue.

For convenience of description, epithelia are classified into different types based on
30 the number of cell layers and the cell shape.

Epithelia which are 1 cell layer thick are called "simple" epithelia. Thus, each cell rests on the basal lamina, but also has a surface facing the lumen/outside world. Epithelia which are 2 or more cell layers thick are called "stratified" epithelia. In stratified epithelia, the basal layer of cells rests on the basal lamina, but subsequent layers do not, and are
35 simply stacked on top of the basal layer. The cells of the most superficial layer have a free surface. "squamous" cells are very flat, like a fried egg, where the yolk is the nucleus. The

5 nucleus is distinctly flattened, the cell is often so thin that this flattened nucleus bulges the cell surface outward. "cuboidal" cells range from true cuboidal where the cell is about as high as it is wide, to a flattened cuboidal where the cell is wider than high. In cuboidal cells the nucleus is usually round, and not flattened as in squamous. "columnar" cells are 2 or more times as high as wide. Nucleus is usually elongated in the long axis of the cell.

10 Squamous cells form the lining of cavities such as the mouth, blood vessels, heart and lungs and make up the outer layers of the skin. Cuboidal epithelium is found in glands and in the lining of the kidney tubules as well as in the ducts of the glands. They also constitute the germinal epithelium which produces the egg cells in the female ovary and the sperm cells in the male testes. Columnar epithelium forms the lining of the stomach and
15 intestines. Some columnar cells are specialized for sensory reception such as in the nose, ears and the taste buds of the tongue.

Ciliated columnar epithelial cells possess fine hair-like outgrowths, cilia on their free surfaces. These cilia are capable of rapid, rhythmic, wavelike beatings in a certain direction. Ciliated epithelium is usually found in the air passages like the nose. It is also found in the
20 uterus and Fallopian tubes of females.

Columnar epithelium with goblet cells is called glandular epithelium. Some parts of the glandular epithelium consist of such a large number of goblet cells that there are only a few normal epithelial cells left. Columnar and cuboidal epithelial cells often become specialized as gland cells which are capable of synthesizing and secreting certain substances
25 such as enzymes, hormones, milk, mucus, sweat, wax and saliva. Unicellular glands consist of single, isolated glandular cells such as the goblet cells. Sometimes a portion of the epithelial tissue becomes invaginated and a multicellular gland is formed. Multicellular glands are composed of clusters of cells. Most glands are multicellular including the salivary glands.

30 Where body linings have to withstand wear and tear, the epithelia are composed of several layers of cells and are then called compound or stratified epithelium. The top cells are flat and scaly and it may or may not be keratinized (i.e. containing a tough, resistant protein called keratin). The mammalian skin is an example of dry, keratinized, stratified epithelium. The lining of the mouth cavity is an example of an unkeratinized, stratified
35 epithelium.

5 *In vitro* Cell Models of Transcytosis

The use of *in vitro* cell models to study transcytosis has many advantages over *in vivo* systems. First, variation among animals is eliminated, as is the confounding issue of cargo possibly being modified or endocytosed by cell types other than the one under study. Moreover, *in vitro* systems can be manipulated in ways not possible *in vivo*, allowing
10 investigators to measure the effects of different variables (e.g., temperatures, pharmacological agents, etc.) with greater precision and to explore the molecular mechanisms of transcytosis.

The integrity of the monolayer is obviously vital to every study of transcytosis, and there are different methods for assessing it. Transepithelial electrical resistance (TER)
15 measurements are commonly used as an indication of tight junction integrity in a monolayer, and commercial instruments are available for these measurements.

Caco-2 cells, human primary colon carcinoma cells, are a well studied model of intestinal absorptive enterocytes. They are the most commonly used intestinal cell line because they differentiate furthest along the cryptto-villus axis and are the easiest to
20 transfect. Caco-2 cells have been especially used to model transcytosis of bacteria, which can cross barrier epithelia in the gut and brain (Zhang JR, et al., 2000. Cell 102(6):827-37), incorporated herein by reference.

There is little evidence for *in vivo* transcytosis of macromolecular cargo in kidney. Nonetheless, MDCK cells, which are derived from dog kidney, are the most-studied
25 epithelial cell model and have been used extensively to study transcytosis. These cells were originally developed by nephrologists for permeability and electrical studies. Their subsequent use by cell biologists for studies of the formation of tight junctions, establishment of polarity, and vesicle traffic have popularized MDCK cells. An advantage is that MDCK cells are easily cultured, easily transfected, and become polarized 3–5 days after
30 seeding. They were used in the now classical studies showing that enveloped viruses bud in a polarized fashion and that the newly synthesized viral membrane glycoproteins are targeted directly from the TGN to the appropriate PM domain. Furthermore, much of the current understanding of the IgA transcytotic pathway and the sorting signals in the pIgA-R comes from the elegant studies performed in MDCK cells. Two MDCK strains with very
35 different features were identified some time ago. The MDCK I cell has a high TER and characteristics reminiscent of the renal collecting duct, whereas the more commonly used

5 MDCK II strain, whose TER is one order of magnitude lower than that of MDCK I cells, has phenotypic features closer to those of the renal proximal tubule.

Both primary cells and cell lines, alone and in coculture with endothelial cells, are being used to study transcytosis in the lung. Clonetics bronchial/tracheal epithelial cell systems contain normal human bronchial/tracheal epithelial cells. This cell system has been
10 used for experimental applications in cancer research, respiratory disease, cellular function and differentiation.

The Clonetics® bovine Brain Microvascular Endothelial Cell System (bMVEC-B) is a model of the "Blood Brain Barrier". The system is designed to significantly improve a researcher's ability to study active and passive transport of drugs across the blood brain
15 barrier, to study brain endothelial cell tight junctions, and to study the basic biology of brain microvascular endothelial cells (Schinket AH, 1999. Advanced Drug Delivery Reviews 36:179-194; Tsukita S. et al., 1998. Molecular dissection of tight junctions: occluding and ZO-1 in Introduction to the Blood - Brain Barrier. Edited by William M Partridge; Inglis et al., 2004. Brain Research 998: 218-229), each of which is incorporated by reference for its
20 teaching of *in vitro* endothelial cell modeling of the blood-brain barrier.

Endometrial cells form an important barrier layer in the genitourinary tract. The cells used to model this system were developed by Kyo et al. and are derived from primary cells immortalized by the addition of the papillomavirus E6/E7 genes and human telomerase reverse transcriptase. The isolated cells have a normal chromosomes and retain
25 their responsiveness to sex-steroid hormones, exhibit glandular structure on three dimensional culture, and lack a transformed phenotype (Kyo S, et al. Am J Pathol., 2003. 163(6):2259-69), incorporated herein by reference for its teaching of this endometrial model.

Methods of Use

30 The use of AAVs to deliver genes to the lung would be of benefit in genetic diseases like cystic fibrosis, pseudohypoaldosteronism, and immotile cilia syndrome. Furthermore, delivering genes to the lung would be of impact in several non-genetic diseases. For example, delivering genes that make antibiotic like peptides to the cells underlying the epithelia would be useful to prevent or treat bronchitis; delivering genes that make growth
35 factors would be of value in common diseases like chronic bronchitis. Also, AAVs could be used to deliver genes that may play a role in asthma, like IL-10, or antibodies to IgE and

5 interleukins. The use of an AAV vector to deliver genes through the alveolar epithelia would be of benefit in genetic diseases like alpha-1-antitrypsin deficiency. Furthermore, delivering genes through the alveolar epithelia would be of significance in several pulmonary non-genetic diseases. For example, delivering genes that make antibiotic like peptides would be useful to prevent or treat pneumonia (perhaps of antibiotic-resistant
10 organisms); delivering genes that make growth factors would be of value in emphysema; delivering genes that over-express the epithelial sodium channel or the Na-K ATPase could be used to treat cardiogenic and non-cardiogenic pulmonary edema; delivering genes that have an anti-fibrosis effect like interferon for pulmonary fibrosis would also be useful. Also, AAVs could be used to deliver genes that may have a systemic effect like anti-hypertension
15 drugs, insulin, coagulation factors, antibiotics, growth factors, hormones and others.

The use of AAVs to deliver genes to the central nervous system (CNS)/ brain would be of benefit in neurological diseases, including Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia,
obsessive compulsive disorder, panic disorder, learning disabilities, ALS, triplet expansions
20 diseases, psychoses, autism, lysosomal storage diseases, Gaucher's disease, Hurler's disease, Krabbe's disease, battens disease, and altered behaviors (e.g., disorders in feeding, sleep patterns, balance, and perception).

The use of AAVs to deliver genes to the gastrointestinal system/ gut would be of benefit in treatment of diseases and/or Gastrointestinal Disorders such as colon cancers,
25 inflammatory bowel disease, diabetes, or Crohn's disease.

The use of AAVs to deliver genes to the genitourinary system would be of benefit in treatment of diseases of the female reproductive tract, molecular defects in implantation disorders, and gynecological cancers. These methods would also have contraceptive applications.

30 The use of AAVs to deliver genes to the kidney would be of benefit in treatment of inherited renal disorders such as polycystic kidney disease, Alport's syndrome, hereditary nephritis, primary hyperoxaluria, and cystinuria.

The use of AAVs for wide-spread delivery of genes across blood vessels into the muscle would be of benefit in neuromuscular diseases like muscular dystrophy and
35 Cardiovascular Disorders such as heart disease, restenosis, atherosclerosis, myocarditis, stoke, angina, or thrombosis.

5 The use of AAVs for wide-spread delivery of genes across blood vessels into any/all tissues of a subject would be of benefit in the treatment of certain cancers (e.g., gastric, ovarian, lung, bladder, liver, and breast).

 The use of AAVs for wide-spread delivery of genes across blood vessels into any/all tissues of a subject would be of benefit in the treatment of certain inflammatory disorders,
10 including, but not limited to, adrenalitis, alveolitis, angiocholecystitis, appendicitis, balanitis, blepharitis, bronchitis, bursitis, carditis, cellulitis, cervicitis, cholecystitis, chorditis, cochitis, colitis, conjunctivitis, cystitis, dermatitis, diverticulitis, encephalitis, endocarditis, esophagitis, eustachitis, fibrositis, folliculitis, gastritis, gastroenteritis, gingivitis, glossitis, hepatosplenitis, keratitis, labyrinthitis, laryngitis, lymphangitis, mastitis,
15 media otitis, meningitis, metritis, mucitis, myocarditis, myositis, myringitis, nephritis, neuritis, orchitis, osteochondritis, otitis, pericarditis, peritendonitis, peritonitis, pharyngitis, phlebitis, poliomyelitis, prostatitis, pulpitis, retinitis, rhinitis, salpingitis, scleritis, sclerochoroiditis, scrotitis, sinusitis, spondylitis, steatitis, stomatitis, synovitis, syringitis, tendonitis, tonsillitis, urethritis, and vaginitis; and disorders that are characterized by
20 inflammation such as hepatitis, rheumatoid arthritis, gout, trauma, pancreatitis, sarcoidosis, dermatitis, renal ischemia-reperfusion injury, Grave's disease, systemic lupus erythematosus, diabetes mellitus, and allogenic transplant rejection.

 The use of AAVs for wide-spread delivery of genes across blood vessels into any/all tissues of a subject would be of benefit in the treatment of other diseases, syndromes and
25 conditions, such as adenosine deaminase deficiency, sickle cell deficiency, thalassemia, hemophilia, diabetes, phenylketonuria, growth disorders, and defects of the immune system.

BAAV

 Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier of the lung, comprising delivering to the lung a BAAV vector comprising the nucleic
30 acid. In one aspect of the method, the epithelial barrier comprises human bronchial, alveolar, tracheal or upper airway epithelial cells. Thus, disclosed is a method of delivering a heterologous nucleic acid across human airway epithelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.

 Disclosed is a method of delivering a heterologous nucleic acid across an epithelial
35 barrier in the brain, comprising delivering to the brain a BAAV vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human cerebral

5 microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier. Thus, disclosed is a method of delivering a heterologous nucleic acid across human cerebral microvascular endothelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.

Disclosed is a method of delivering a heterologous nucleic acid across the epithelial
10 barrier of blood vessels into the muscle, comprising delivering to the blood stream a BAAV vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human vascular endothelial cells.

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier in the genitourinary tract, comprising delivering to the genitourinary tract a BAAV
15 vector comprising the nucleic acid genitourinary tract. In one aspect of the method, the epithelial barrier comprises human endometrial or urinary epithelial cells. Thus, disclosed is a method of delivering a heterologous nucleic acid across human endometrial epithelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial
20 barrier in the kidney, comprising delivering to the genitourinary tract a BAAV vector comprising the nucleic acid genitourinary tract. In one aspect of the method, the epithelial barrier comprises human renal collecting ducts or proximal tubules. Thus, disclosed is a method of delivering a heterologous nucleic acid across human kidney epithelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.

25 Disclosed is a method of transcytosing lung epithelial cells of a subject comprising contacting the lung epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human bronchial, tracheal, or upper airway epithelial cells.

Disclosed is a method of transcytosing CNS epithelial cells of a subject comprising
30 contacting the CNS epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.

Disclosed is a method of transcytosing vascular epithelial cells of a subject
35 comprising contacting the vascular epithelial cells of the subject with a BAAV vector

5 comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human vascular endothelial cells of the blood brain barrier.

Disclosed is a method of transcytosing genitourinary tract epithelial cells of a subject comprising contacting the genitourinary tract epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial
10 cells are human endometrial or urinary tract epithelial cells.

Disclosed is a method of transcytosing genitourinary tract epithelial cells of a subject comprising contacting the kidney epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human renal collecting ducts or proximal tubules

15 AAV5

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier in the gut, comprising delivering to the gut an AAV5 vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human absorptive enterocytes or M cells. Thus, disclosed is a method of delivering a heterologous nucleic acid
20 across human gut epithelial cells enterocytes, comprising delivering to the cells an AAV5 vector comprising the nucleic acid.

Disclosed is a method of transcytosing gut epithelial cells of a subject comprising contacting the gut epithelial cells of the subject with an AAV5 vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human
25 absorptive enterocytes.

AAV4

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier in the gut, comprising delivering to the gut an AAV4 vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human absorptive
30 enterocytes or M cells. Thus, disclosed is a method of delivering a heterologous nucleic acid across human gut epithelial cells enterocytes, comprising delivering to the cells an AAV4 vector comprising the nucleic acid.

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier in the lung, comprising delivering to the lung an AAV4 vector comprising the
35 nucleic acid. In one aspect of the method, the epithelial barrier comprises human bronchial,

5 tracheal, or upper airway epithelial cells. Thus, disclosed is a method of delivering a heterologous nucleic acid across human airway epithelial cells, comprising delivering to the cells an AAV4 vector comprising the nucleic acid.

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier in the CNS, comprising delivering to the CNS an AAV4 vector comprising the
10 nucleic acid. In one aspect of the method, the epithelial barrier comprises human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier. Thus, disclosed is a method of delivering a heterologous nucleic acid across human cerebral microvascular endothelial cells, comprising delivering to the cells a AAV4 vector comprising the nucleic acid.

15 Disclosed is a method of delivering a heterologous nucleic acid across the epithelial barrier of blood vessels into the muscle, comprising delivering to the blood stream an AAV4 vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human vascular endothelial cells of the blood brain barrier.

Disclosed is a method of delivering a heterologous nucleic acid across an epithelial
20 barrier in the genitourinary tract, comprising delivering to the genitourinary tract an AAV4 vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human endometrial or urinary epithelial cells. Thus, disclosed is a method of delivering a heterologous nucleic acid across human endometrial epithelial cells, comprising delivering to the cells an AAV4 vector comprising the nucleic acid.

25 Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier in the kidneys, comprising delivering to the kidneys an AAV4 vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human renal collecting ducts or proximal tubules. Thus, disclosed is a method of delivering a heterologous nucleic acid across human kidney epithelial cells, comprising delivering to the
30 cells an AAV4 vector comprising the nucleic acid.

Disclosed is a method of transcytosing lung epithelial cells of a subject comprising contacting the lung epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human bronchial, tracheal, or upper airway epithelial cells.

35 Disclosed is a method of transcytosing CNS epithelial cells of a subject comprising contacting the CNS epithelial cells of the subject with an AAV4 vector comprising a

5 heterologous nucleic acid. In one aspect of the method, the epithelial cells are human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.

Disclosed is a method of transcytosing vascular epithelial cells of a subject comprising contacting the vascular epithelial cells of the subject with an AAV4 vector
10 comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are vascular endothelial cells of the blood brain barrier.

Disclosed is a method of transcytosing genitourinary tract epithelial cells of a subject comprising contacting the genitourinary epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are
15 human endometrial or urinary epithelial cells.

Disclosed is a method of transcytosing kidney epithelial cells of a subject comprising contacting the kidney epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human renal collecting ducts or proximal tubules

20 Disclosed is a method of transcytosing gut epithelial cells of a subject comprising contacting the gut epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human absorptive enterocytes.

AAV7

25 Disclosed is a method of delivering a heterologous nucleic acid across an epithelial barrier in the CNS, comprising delivering to the CNS an AAV7 vector comprising the nucleic acid. In one aspect of the method, the epithelial barrier comprises human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier. Thus, disclosed is a method of delivering a heterologous nucleic acid across human
30 cerebral microvascular endothelial cells, comprising delivering to the cells an AAV7 vector comprising the nucleic acid.

Disclosed is a method of transcytosing CNS epithelial cells of a subject comprising contacting the CNS epithelial cells of the subject with an AAV7 vector comprising a heterologous nucleic acid. In one aspect of the method, the epithelial cells are human
35 cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.

5 **Inhibition of Transcytosis to Increase Transduction**

Described herein is a method for re-directing virus that enters a cell by transcytosis to result in transduction of the cell by blocking exocytosis. Thus, provided is a method of improving the efficiency of nucleic acid delivery to epithelial cells, comprising delivering to the cells an inhibitor of exocytosis and an AAV vector containing the nucleic acid. Also
10 provided is a method for transducing cells that have transcytosis activity but are normally resistant to transduction comprising administering to the cells inhibitors of exocytosis.

In one aspect of the methods, the AAV vector is derived from AAV4, AAV5, or BAAV. In a further aspect of the methods, the epithelial cell barriers are located in the kidney, gut, lung or vascular endothelium

15 Thus, disclosed is a method of delivering a heterologous nucleic acid to human airway epithelial cells, comprising delivering to the cells and an inhibitor of exocytosis and an AAV4 vector comprising the nucleic acid.

Further disclosed is a method of delivering a heterologous nucleic acid to human kidney epithelial cells, comprising delivering to the cells and an inhibitor of exocytosis and
20 an AAV4 vector comprising the nucleic acid.

Further disclosed is a method of delivering a heterologous nucleic acid to human vascular endothelial cells, comprising delivering to the cells and an inhibitor of exocytosis and an AAV4 vector comprising the nucleic acid.

Further disclosed is a method of delivering a heterologous nucleic acid to human
25 airway epithelial cells, comprising delivering to the cells and an inhibitor of exocytosis and a BAAV vector comprising the nucleic acid.

Further disclosed is a method of delivering a heterologous nucleic acid to human kidney epithelial cells, comprising delivering to the cells and an inhibitor of exocytosis and a BAAV vector comprising the nucleic acid.

30 Further disclosed is a method of delivering a heterologous nucleic acid to human vascular endothelial cells, comprising delivering to the cells and an inhibitor of exocytosis and a BAAV vector comprising the nucleic acid.

Further disclosed is a method of delivering a heterologous nucleic acid to human gut epithelial cells, comprising delivering to the cells and an inhibitor of exocytosis and an
35 AAV5 vector comprising the nucleic acid.

5 In one aspect of the disclosed methods, the inhibitors of exocytosis are chemical modifiers. In a further aspect of the methods, the chemical modifier is tannic acid, wherein the tannic acid is delivered to the basal lateral surface of the epithelial cells.

Compositions and methods for making AAV4 vectors

 Compositions and methods for making and using AAV4 vectors have been
10 previously described in U.S. Patent No. 6,468,524, which is hereby incorporated herein by reference for this teaching.

 Provided is the nucleotide sequence of the adeno-associated virus 4 (AAV4) genome and vectors and particles derived therefrom. Specifically, provided is a nucleic acid vector comprising a pair of AAV4 inverted terminal repeats (ITRs) and a promoter between the
15 inverted terminal repeats. The AAV4 ITRs are exemplified by the nucleotide sequence set forth in SEQ ID NO:6 and SEQ ID NO:20; however, these sequences can have minor modifications and still be contemplated to constitute AAV4 ITRs. The nucleic acid listed in SEQ ID NO:6 depicts the ITR in the "flip" orientation of the ITR. The nucleic acid listed in SEQ ID NO:20 depicts the ITR in the "flop" orientation of the ITR. Minor modifications in
20 an ITR of either orientation are those that will not interfere with the hairpin structure formed by the AAV4 ITR as described herein and known in the art. Furthermore, to be considered within the term "AAV4 ITRs" the nucleotide sequence must retain the Rep binding site described herein and exemplified in SEQ ID NO:6 and SEQ ID NO:20, *i.e.*, it must retain one or both features described herein that distinguish the AAV4 ITR from the AAV2 ITR:
25 (1) four (rather than three as in AAV2) "GAGC" repeats and (2) in the AAV4 ITR Rep binding site the fourth nucleotide in the first two "GAGC" repeats is a T rather than a C.

 The promoter can be any desired promoter, selected by known considerations, such as the level of expression of a nucleic acid functionally linked to the promoter and the cell type in which the vector is to be used. Promoters can be an exogenous or an endogenous
30 promoter. Promoters can include, for example, known strong promoters such as SV40 or the inducible metallothionein promoter, or an AAV promoter, such as an AAV p5 promoter. Additional examples of promoters include promoters derived from actin genes, immunoglobulin genes, cytomegalovirus (CMV), adenovirus, bovine papilloma virus, adenoviral promoters, such as the adenoviral major late promoter, an inducible heat shock
35 promoter, respiratory syncytial virus, Rous sarcomas virus (RSV), etc. Specifically, the promoter can be AAV2 p5 promoter or AAV4 p5 promoter. More specifically, the AAV4

5 p5 promoter can be about nucleotides 130 to 291 of SEQ ID NO: 1. Additionally, the p5 promoter may be enhanced by nucleotides 1-130. Furthermore, smaller fragments of p5 promoter that retain promoter activity can readily be determined by standard procedures including, for example, constructing a series of deletions in the p5 promoter, linking the deletion to a reporter gene, and determining whether the reporter gene is expressed, *i.e.*,
10 transcribed and/or translated.

The present invention also contemplates any unique fragment of these AAV4 nucleic acids, including the AAV4 nucleic acids set forth in SEQ ID NOs: 1, 3, 5, 6, 7, 12-15, 17 and 19. Fragments can be, for example, at least about 30, 40, 50, 75, 100, 200 or 500 nucleotides in length. The nucleic acid can be single or double stranded, depending upon the
15 purpose for which it is intended.

The present invention further provides an AAV4 Capsid polypeptide or a unique fragment thereof. AAV4 capsid polypeptide is encoded by ORF 2 of AAV4. Specifically, provided is an AAV4 Capsid protein comprising the amino acid sequence encoded by nucleotides 2260-4464 of the nucleotide sequence set forth in SEQ ID NO:1, or a unique
20 fragment of such protein. The present invention also provides an AAV4 Capsid protein consisting essentially of the amino acid sequence encoded by nucleotides 2260-4464 of the nucleotide sequence set forth in SEQ ID NO:1, or a unique fragment of such protein. The present invention further provides the individual AAV4 coat proteins, VP1, VP2 and VP3. Thus, provided is an isolated polypeptide having the amino acid sequence set forth in SEQ
25 ID NO:4 (VP1). The present invention additionally provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:16 (VP2). The present invention also provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:18 (VP3). By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by any AAV4 capsid gene that is of sufficient length to be unique to the AAV4 Capsid protein.
30 Substitutions and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide. However, an AAV4 Capsid polypeptide including all three coat proteins will have at least about 63% overall homology to the polypeptide encoded by nucleotides 2260-4464 of the sequence set forth in SEQ ID NO: 1. The protein can have about 65%, about
35 70%, about 75%, about 80%, about 85%, about 90%, about 95% or even 100% homology to the amino acid sequence encoded by the nucleotides 4467 of the sequence set forth in SEQ

5 ID NO:1. An AAV4 VP2 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90% about 95% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:16. An AAV4 VP3 polypeptide can have at least about 60%, about 70%, about 80%, about 90% about 95% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:18.

10 The herein described AAV4 nucleic acid vector can be encapsidated in an AAV particle. In particular, it can be encapsidated in an AAV1 particle, an AAV2 particle, an AAV3 particle, an AAV4 particle, or an AAV5 particle by standard methods using the appropriate capsid proteins in the encapsidation process, as long as the nucleic acid vector fits within the size limitation of the particle utilized. The encapsidation process itself is
15 standard in the art.

An AAV4 particle is a viral particle comprising an AAV4 capsid protein. An AAV4 capsid polypeptide encoding the entire VP1, VP2, and VP3 polypeptide can overall have at least about 63% homology to the polypeptide having the amino acid sequence encoded by nucleotides 2260-4464 set forth in SEQ ID NO:1 (AAV4 capsid protein). The capsid protein
20 can have about 70% homology, about 75% homology, 80% homology, 85% homology, 90% homology, 95% homology, 98% homology, 99% homology, or even 100% homology to the protein having the amino acid sequence encoded by nucleotides 2260-4464 set forth in SEQ ID NO:1. The particle can be a particle comprising both AAV4 and AAV2 capsid protein, *i.e.*, a chimeric protein. Variations in the amino acid sequence of the AAV4 capsid protein
25 are contemplated herein, as long as the resulting viral particle comprising the AAV4 capsid remains antigenically or immunologically distinct from AAV2, as can be routinely determined by standard methods. Specifically, for example, ELISA and Western blots can be used to determine whether a viral particle is antigenically or immunologically distinct from AAV2. Furthermore, the AAV4 viral particle preferably retains tissue tropism
30 distinction from AAV2, such as that exemplified in the examples herein, though an AAV4 chimeric particle comprising at least one AAV4 coat protein may have a different tissue tropism from that of an AAV4 particle consisting only of AAV4 coat proteins.

An AAV4 particle is a viral particle comprising an AAV4 capsid protein. An AAV4 capsid polypeptide encoding the entire VP1, VP2, and VP3 polypeptide can overall have at
35 least about 63% homology to the polypeptide having the amino acid sequence encoded by nucleotides 2260-4467 set forth in SEQ ID NO:1 (AAV4 capsid protein). The capsid protein

5 can have about 70% homology, about 75% homology, 80% homology, 85% homology, 90% homology, 95% homology, 98% homology, 99% homology, or even 100% homology to the protein having the amino acid sequence encoded by nucleotides 2260-4467 set forth in SEQ ID NO:1. The particle can comprise only VP1 and VP3 and still stably transduce cells. The particle can be a particle comprising both AAV4 and AAV2 capsid protein, *i.e.*, a
10 chimeric protein. Variations in the amino acid sequence of the AAV4 capsid protein are contemplated herein, as long as the resulting viral particle comprising the AAV4 capsid remains antigenically or immunologically distinct from AAV2, as can be routinely determined by standard methods. Specifically, for example, ELISA and Western blots can be used to determine whether a viral particle is antigenically or immunologically distinct
15 from AAV2. Furthermore, the AAV4 viral particle preferably retains tissue tropism distinction from AAV2, such as that exemplified in the examples herein, though an AAV4 chimeric particle comprising at least one AAV4 coat protein may have a different tissue tropism from that of an AAV4 particle consisting only of AAV4 coat proteins.

The invention further provides an AAV4 particle containing, *i.e.*, encapsidating, a
20 vector comprising a pair of AAV2 inverted terminal repeats. The nucleotide sequence of AAV2 ITRs is known in the art. Furthermore, the particle can be a particle comprising both AAV4 and AAV2 capsid protein, *i.e.*, a chimeric protein. The vector encapsidated in the particle can further comprise an exogenous nucleic acid inserted between the inverted terminal repeats.

25 The present invention further provides an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:1 (AAV4 genome). This nucleic acid, or portions thereof, can be inserted into other vectors, such as plasmids, yeast artificial chromosomes, or other viral vectors, if desired, by standard cloning methods. The present invention also provides an isolated nucleic acid consisting essentially of the nucleotide
30 sequence set forth in SEQ ID NO:1. The nucleotides of SEQ ID NO:1 can have minor modifications and still be contemplated by the present invention. For example, modifications that do not alter the amino acid encoded by any given codon (such as by modification of the third, "wobble," position in a codon) can readily be made, and such alterations are known in the art. Furthermore, modifications that cause a resulting neutral
35 amino acid substitution of a similar amino acid can be made in a coding region of the genome. Additionally, modifications as described herein for the AAV4 components, such as

5 the ITRs, the p5 promoter, etc. are contemplated in this invention.

The present invention additionally provides an isolated nucleic acid that selectively hybridizes with an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:1 (AAV4 genome). The present invention further provides an isolated nucleic acid that selectively hybridizes with an isolated nucleic acid comprising the
10 nucleotide sequence set forth in SEQ ID NO:1 (AAV4 genome). By "selectively hybridizes" as used in the claims is meant a nucleic acid that specifically hybridizes to the particular target nucleic acid under sufficient stringency conditions to selectively hybridize to the target nucleic acid without significant background hybridization to a nucleic acid encoding an unrelated protein, and particularly, without detectably hybridizing to AAV2. Thus, a
15 nucleic acid that selectively hybridizes with a nucleic acid of the present invention will not selectively hybridize under stringent conditions with a nucleic acid encoding a different protein, and vice versa. Therefore, nucleic acids for use, for example, as primers and probes to detect or amplify the target nucleic acids are contemplated herein. Nucleic acid fragments that selectively hybridize to any given nucleic acid can be used, *e.g.*, as primers and or
20 probes for further hybridization or for amplification methods (*e.g.*, polymerase chain reaction (PCR), ligase chain reaction (LCR)). Additionally, for example, a primer or probe can be designed that selectively hybridizes with both AAV4 and a gene of interest carried within the AAV4 vector (*i.e.*, a chimeric nucleic acid).

The present invention further provides an isolated nucleic acid encoding an adeno-
25 associated virus 4 Rep protein. The AAV4 Rep proteins are encoded by open reading frame (ORF) 1 of the AAV4 genome. The AAV4 Rep genes are exemplified by the nucleic acid set forth in SEQ ID NO:3 (AAV4 ORF1), and include a nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:3 and a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:3. The present invention also includes a nucleic
30 acid encoding the amino acid sequence set forth in SEQ ID NO: 2 (polypeptide encoded by AAV4 ORF1). However, the present invention includes that the Rep genes nucleic acid can include any one, two, three, or four of the four Rep proteins, in any order, in such a nucleic acid. Furthermore, minor modifications are contemplated in the nucleic acid, such as silent mutations in the coding sequences, mutations that make neutral or conservative changes in
35 the encoded amino acid sequence, and mutations in regulatory regions that do not disrupt the expression of the gene. Examples of other minor modifications are known in the art.

5 Further modifications can be made in the nucleic acid, such as to disrupt or alter expression of one or more of the Rep proteins in order to, for example, determine the effect of such a disruption; such as to mutate one or more of the Rep proteins to determine the resulting effect, etc. However, in general, a modified nucleic acid encoding all four Rep proteins will have at least about 90%, about 93%, about 95%, about 98% or 100% homology to the
10 sequence set forth in SEQ ID NO:3, and the Rep polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence set forth in SEQ ID NO:2.

The present invention also provides an isolated nucleic acid that selectively hybridizes with a nucleic acid consisting essentially of the nucleotide sequence set forth in
15 SEQ ID NO:3 and an isolated nucleic acid that selectively hybridizes with a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:3. "Selectively hybridizing" is defined elsewhere herein.

The present invention also provides each individual AAV4 Rep protein and the nucleic acid encoding each. Thus provided is the nucleic acid encoding a Rep 40 protein,
20 and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:12, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:12, and a nucleic acid encoding the adeno-associated virus 4 Rep protein having the amino acid sequence set forth in SEQ ID NO:8. The present invention also provides the nucleic acid encoding a Rep 52 protein, and in particular an isolated
25 nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:13, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:13, and a nucleic acid encoding the adeno-associated virus 4 Rep protein having the amino acid sequence set forth in SEQ ID NO:9. The present invention further provides the nucleic acid encoding a Rep 68 protein, and in particular an isolated nucleic acid comprising the
30 nucleotide sequence set forth in SEQ ID NO:14, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:14, and a nucleic acid encoding the adeno-associated virus 4 Rep protein having the amino acid sequence set forth in SEQ ID NO:10. And, further, provided is the nucleic acid encoding a Rep 78 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID
35 NO:15, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:15, and a nucleic acid encoding the adeno-associated virus 4 Rep protein

5 having the amino acid sequence set forth in SEQ ID NO:11. As described elsewhere herein, these nucleic acids can have minor modifications, including silent nucleotide substitutions, mutations causing neutral amino acid substitutions in the encoded proteins, and mutations in control regions that do not or minimally affect the encoded amino acid sequence.

The present invention further provides a nucleic acid encoding the entire AAV4
10 Capsid polypeptide. Specifically, provided is a nucleic acid having the nucleotide sequence set for the nucleotides 2260-4467 of SEQ ID NO:1. Furthermore, provided is a nucleic acid encoding each of the three AAV4 coat proteins, VP1, VP2, and VP3. Thus, provided is a nucleic acid encoding AAV4 VP1, a nucleic acid encoding AAV4 VP2, and a nucleic acid encoding AAV4 VP3. Thus, provided is a nucleic acid encoding the amino acid sequence
15 set forth in SEQ ID NO:4 (VP1); a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:16 (VP2), and a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:18 (VP3). The present invention also specifically provides a nucleic acid comprising SEQ ID NO:5 (VP1 gene); a nucleic acid comprising SEQ ID NO:17 (VP2 gene); and a nucleic acid comprising SEQ ID NO:19 (VP3 gene). The present invention also
20 specifically provides a nucleic acid consisting essentially of SEQ ID NO:5 (VP1 gene), a nucleic acid consisting essentially of SEQ ID NO:17 (VP2 gene), and a nucleic acid consisting essentially of SEQ ID NO:19 (VP3 gene). Furthermore, a nucleic acid encoding an AAV4 capsid protein VP1 is set forth as nucleotides 2260-4467 of SEQ ID NO:1; a nucleic acid encoding an AAV4 capsid protein VP2 is set forth as nucleotides 2668-4467 of
25 SEQ ID NO:1; and a nucleic acid encoding an AAV4 capsid protein VP3 is set forth as nucleotides 2848-4467 of SEQ ID NO:1. Minor modifications in the nucleotide sequences encoding the capsid, or coat, proteins are contemplated, as described above for other AAV4 nucleic acids.

Provided is an isolated AAV4 Rep protein. AAV4 Rep polypeptide is encoded by
30 ORF1 of AAV4. Specifically, provided is an AAV4 Rep polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2, or a unique fragment thereof. The present invention also provides an AAV4 Rep polypeptide consisting essentially of the amino acid sequence set forth in SEQ ID NO:2, or a unique fragment thereof. Additionally, nucleotides 291-2306 of the AAV4 genome, which genome is set forth in SEQ ID NO:1, encode the
35 AAV4 Rep polypeptide. The present invention also provides each AAV4 Rep protein. Thus provided is AAV4 Rep 40, or a unique fragment thereof. The present invention particularly

5 provides Rep 40 having the amino acid sequence set forth in SEQ ID NO:8. Provided is AAV4 Rep 52, or a unique fragment thereof. The present invention particularly provides Rep 52 having the amino acid sequence set forth in SEQ ID NO:9. Provided is AAV4 Rep 68, or a unique fragment thereof. The present invention particularly provides Rep 68 having the amino acid sequence set forth in SEQ ID NO:10. Provided is AAV4 Rep 78, or a unique
10 fragment thereof. The present invention particularly provides Rep 78 having the amino acid sequence set forth in SEQ ID NO:11. By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by AAV rep gene that is of sufficient length to be unique to the Rep polypeptide. Substitutions and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as
15 glycosylation, to the polypeptide. However, a polypeptide including all four Rep proteins will encode a polypeptide having at least about 91% overall homology to the sequence set forth in SEQ ID NO:2, and it can have about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence set forth in SEQ ID NO:2.

The present invention further provides an AAV4 Capsid polypeptide or a unique
20 fragment thereof. AAV4 capsid polypeptide is encoded by ORF 2 of AAV4. Specifically, provided is an AAV4 Capsid protein comprising the amino acid sequence encoded by nucleotides 2260-4467 of the nucleotide sequence set forth in SEQ ID NO:1, or a unique fragment of such protein. The present invention also provides an AAV4 Capsid protein consisting essentially of the amino acid sequence encoded by nucleotides 2260-4467 of the
25 nucleotide sequence set forth in SEQ ID NO:1, or a unique fragment of such protein. The present invention further provides the individual AAV4 coat proteins, VP1, VP2 and VP3. Thus, provided is an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:4 (VP1). The present invention additionally provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:16 (VP2). The present invention also
30 provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:18 (VP3). By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by any AAV4 capsid gene that is of sufficient length to be unique to the AAV4 Capsid protein. Substitutions and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the
35 polypeptide. However, an AAV4 Capsid polypeptide including all three coat proteins will have at least about 63% overall homology to the polypeptide encoded by nucleotides 2260-

5 4467 of the sequence set forth in SEQ ID NO: 1. The protein can have about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% or even 100% homology to the amino acid sequence encoded by the nucleotides 2260-4467 of the sequence set forth in SEQ ID NO:4. An AAV4 VP2 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90% about 95% or about 100% homology to the amino acid
10 sequence set forth in SEQ ID NO:16. An AAV4 VP3 polypeptide can have at least about 60%, about 70%, about 80%, about 90% about 95% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:18.

The AAV inverted terminal repeats in the vector for the herein described delivery methods can be AAV4 inverted terminal repeats. Specifically, they can comprise the nucleic
15 acid whose nucleotide sequence is set forth in SEQ ID NO:6 or the nucleic acid whose nucleotide sequence is set forth in SEQ ID NO:20, or any fragment thereof demonstrated to have ITR functioning. The ITRs can also consist essentially of the nucleic acid whose nucleotide sequence is set forth in SEQ ID NO:6 or the nucleic acid whose nucleotide sequence is set forth in SEQ ID NO:20. Furthermore, the AAV inverted terminal repeats in
20 the vector for the herein described nucleic acid delivery methods can also comprise AAV2 inverted terminal repeats. Additionally, the AAV inverted terminal repeats in the vector for this delivery method can also consist essentially of AAV2 inverted terminal repeats.

Compositions and methods for making AAV5 vectors

Compositions and methods for making and using AAV5 vectors have been
25 previously described in U.S. Patent Application No. 09/717,789, filed November 21, 2000, and in U.S. Patent No. 6,855,314, which are hereby incorporated herein by reference for this teaching.

The present application provides a recombinant adeno-associated virus 5 (AAV5). This virus has one or more of the characteristics described below. The compositions of the
30 present invention do not include wild-type AAV5. The methods of the present invention can use either wild-type AAV5 or recombinant AAV5-based delivery.

Provided are novel AAV5 particles, recombinant AAV5 vectors, recombinant AAV5 virions and novel AAV5 nucleic acids and polypeptides. An AAV5 particle is a viral
particle comprising an AAV5 capsid protein. A recombinant AAV5 vector is a nucleic acid
35 construct that comprises at least one unique nucleic acid of AAV5. A recombinant AAV5 virion is a particle containing a recombinant AAV5 vector, wherein the particle can be either

5 an AAV5 particle as described herein or a non-AAV5 particle. Alternatively, the recombinant AAV5 virion is an AAV5 particle containing a recombinant vector, wherein the vector can be either an AAV5 vector as described herein or a non-AAV5 vector. These vectors, particles, virions, nucleic acids and polypeptides are described below.

Provided is the nucleotide sequence of the AAV5 genome and vectors and particles
10 derived therefrom. Specifically, provided is a nucleic acid vector comprising a pair of AAV5 inverted terminal repeats (ITRs) and a promoter between the inverted terminal repeats. While the rep proteins of AAV2 and AAV5 will bind to either a type 2 ITR or a type 5 ITR, efficient genome replication only occurs when type 2 Rep replicates a type 2 ITR and a type 5 Rep replicates a type 5 ITR. This specificity is the result of a difference in
15 DNA cleavage specificity of the two Reps which is necessary for replication. AAV5 Rep cleaves at CGGT[^]GTGA (SEQ ID NO: 43) and AAV2 Rep cleaves at CGGT[^]TGAG (SEQ ID NO: 44) (Chiorini et al., 1999. J. Virol. 73 (5) 4293-4298). Mapping of the AAV5 ITR terminal resolution site (TRS) identified this distinct cleavage site, CGGT[^]GTGA, which is absent from the ITRs of other AAV serotypes. Therefore, the minimum sequence necessary
20 to distinguish AAV5 from AAV2 is the TRS site where Rep cleaves in order to replicate the virus. Examples of the type 5 ITRs are shown in SEQ ID NO: 41 and SEQ ID NO: 42, AAV5 ITR "flip" and AAV5 "flop", respectively. Minor modifications in an ITR of either orientation are contemplated and are those that will not interfere with the hairpin structure formed by the AAV5 ITR as described herein. Furthermore, to be considered within the
25 term "AAV5 ITR" the nucleotide sequence must retain one or more features described herein that distinguish the AAV5 ITR from the ITRs of other serotypes, e.g. it must retain the Rep binding site described herein.

The D- region of the AAV5 ITR (SEQ ID NO: 45), a single stranded region of the ITR, inboard of the TRS site, has been shown to bind a factor which depending on its
30 phosphorylation state correlates with the conversion of the AAV from a single stranded genome to a transcriptionally active form that allows for expression of the viral DNA. This region is conserved between AAV2, 3, 4, and 6 but is divergent in AAV5. The D+ region is the reverse complement of the D- region.

The promoter can be any desired promoter, selected by known considerations, such
35 as the level of expression of a nucleic acid functionally linked to the promoter and the cell type in which the vector is to be used. That is, the promoter can be tissue/cell-specific.

5 Promoters can be prokaryotic, eukaryotic, fungal, nuclear, mitochondrial, viral or plant promoters. Promoters can be exogenous or endogenous to the cell type being transduced by the vector. Promoters can include, for example, bacterial promoters, known strong promoters such as SV40 or the inducible metallothionein promoter, or an AAV promoter, such as an AAV p5 promoter. Additionally, chimeric regulatory promoters for targeted gene
10 expression can be utilized. Examples of these regulatory systems, which are known in the art, include the tetracycline based regulatory system which utilizes the tet transactivator protein (tTA), a chimeric protein containing the VP16 activation domain fused to the tet repressor of Escherichia coli, the IPTG based regulatory system, the CID based regulatory system, and the Ecdysone based regulatory system. Other promoters include promoters
15 derived from actin genes, immunoglobulin genes, cytomegalovirus (CMV), adenovirus, bovine papilloma virus, adenoviral promoters, such as the adenoviral major late promoter, an inducible heat shock promoter, respiratory syncytial virus, Rous sarcomas virus (RSV), etc., specifically, the promoter can be AAV2 p5 promoter or AAV5 p5 promoter. More specifically, the AAV5 p5 promoter can be about same location in SEQ ID NO: 23 as the
20 AAV2 p5 promoter, in the corresponding AAV2 published sequence. An example of an AAV5 p5 promoter is nucleotides 220-338 of SEQ ID NO: 23. Additionally, the p5 promoter may be enhanced by nucleotides 1-130 of SEQ ID NO: 23. Furthermore, smaller fragments of p5 promoter that retain promoter activity can readily be determined by standard procedures including, for example, constructing a series of deletions in the p5 promoter,
25 linking the deletion to a reporter gene, and determining whether the reporter gene is expressed, i.e., transcribed and/or translated. The promoter can be the promoter of any of the AAV serotypes, and can be the p19 promoter (SEQ ID NO: 38) or the p40 promoter set forth in the sequence listing as SEQ ID NO: 39.

It should be recognized that any errors in any of the nucleotide sequences disclosed
30 herein can be corrected, for example, by using the hybridization procedure described below with various probes derived from the described sequences such that the coding sequence can be reisolated and resequenced. Rapid screening for point mutations can also be achieved with the use of polymerase chain reaction single strand conformation polymorphism (PCR SSCP). The corresponding amino acid sequence can then be corrected accordingly.

35 The AAV5-derived vector can include any normally occurring AAV5 sequences in addition to an ITR and promoter. Examples of vector constructs are provided below.

5 The present vector or AAV5 particle or recombinant AAV5 virion can utilize any unique fragment of the present AAV5 nucleic acids, including the AAV5 nucleic acids set forth in SEQ ID NOS: 23 and 29-33, 35, 37, 38, 39 and 40. To be unique, the fragment must be of sufficient size to distinguish it from other known sequences, most readily determined by comparing any nucleic acid fragment to the nucleotide sequences of nucleic acids in
10 computer databases, such as GenBank. Such comparative searches are standard in the art. Typically, a unique fragment useful as a primer or probe will be at least about 8 or 10, preferable at least 20 or 25 nucleotides in length, depending upon the specific nucleotide content of the sequence. Additionally, fragments can be, for example, at least about 30, 40, 50, 75, 100, 200 or 500 nucleotides in length and can encode polypeptides or be probes. The
15 nucleic acid can be single or double stranded, depending upon the purpose for which it is intended. Where desired, the nucleic acid can be RNA.

 The present invention further provides an isolated AAV5 capsid protein to contain the vector. In particular, provided is not only a polypeptide comprising all three AAV5 coat proteins, i.e., VP1, VP2 and VP3, but also a polypeptide comprising each AAV5 coat
20 protein individually, SEQ ID NOS: 26, 27, and 28, respectively. Thus an AAV5 particle comprising an AAV5 capsid protein comprises at least one AAV5 coat protein VP1, VP2 or VP3. An AAV5 particle comprising an AAV5 capsid protein can be utilized to deliver a nucleic acid vector to a cell, tissue or subject. For example, the herein described AAV5 vectors can be encapsidated in an AAV5 capsid-derived particle and utilized in a gene
25 delivery method. Furthermore, other viral nucleic acids can be encapsidated in the AAV5 particle and utilized in such delivery methods. For example, an AAV1, 2,3,4, or 6 vector (e.g. AAV1,2,3,4, or 6 ITR and nucleic acid of interest) can be encapsidated in an AAV5 particle and administered. Furthermore, an AAV5 chimeric capsid incorporating both AAV2 capsid and AAV5 capsid sequences can be generated, by standard cloning methods,
30 selecting regions from the known sequences of each protein as desired. For example, particularly antigenic regions of the AAV2 capsid protein can be replaced with the corresponding region of the AAV5 capsid protein. In addition to chimeric capsids incorporating AAV2 capsid sequences, chimeric capsids incorporating AAV1, 3, 4, or 6 and AAV5 capsid sequences can be generated, by standard cloning methods, selecting regions
35 from the known sequences of each protein as desired. The particle can also comprise only VP1 and VP3 capsid proteins.

5 The capsids can also be modified to alter their specific tropism by genetically altering the capsid to encode a specific ligand to a cell surface receptor. Alternatively, the capsid can be chemically modified by conjugating a ligand to a cell surface receptor. By genetically or chemically altering the capsids, the tropism can be modified to direct AAV5 to a particular cell or population of cells. The capsids can also be altered immunologically
10 by conjugating the capsid to an antibody that recognizes a specific protein on the target cell or population of cells.

 The capsids can also be assembled into empty particles by expression in mammalian, bacterial, fungal or insect cells. For example, AAV2 particles are known to be made from VP3 and VP2 capsid proteins in baculovirus. The same basic protocol can produce an empty
15 AAV5 particle comprising an AAV5 capsid protein.

 The herein described recombinant AAV5 nucleic acid derived vector can be encapsidated in an AAV particle. In particular, it can be encapsidated in an AAV1 particle, an AAV2 particle, an AAV3 particle, an AAV4 particle, an AAV5 particle or an AAV6 particle, a portion of any of these capsids, or a chimeric capsid particle as described above,
20 by standard methods using the appropriate capsid proteins in the encapsidation process, as long as the nucleic acid vector fits within the size limitation of the particle utilized. The encapsidation process itself is standard in the art. The AAV5 replication machinery, i.e. the rep initiator proteins and other functions required for replication, can be utilized to produce the AAV5 genome that can be packaged in an AAV1, 2, 3, 4, 5 or 6 capsid.

25 The recombinant AAV5 virion containing a vector can also be produced by recombinant methods utilizing multiple plasmids. In one example, the AAV5 rep nucleic acid would be cloned into one plasmid, the AAV5 ITR nucleic acid would be cloned into another plasmid and the AAV1, 2, 3, 4, 5 or 6 capsid nucleic acid would be cloned on another plasmid. These plasmids would then be introduced into cells. The cells that were
30 efficiently transduced by all three plasmids, would exhibit specific integration as well as the ability to produce recombinant AAV5 virion. Additionally, two plasmids could be used where the AAV5 rep nucleic acid would be cloned into one plasmid and the AAV5 ITR and AAV5 capsid would be cloned into another plasmid. These plasmids would then be introduced into cells. The cells that were efficiently transduced by both plasmids, would
35 exhibit specific integration as well as the ability to produce recombinant AAV5 virion.

5 An AAV5 capsid polypeptide encoding the entire VP1, VP2, and VP3 polypeptide can have greater than 56% overall homology to the polypeptide having the amino acid sequence encoded by nucleotides in SEQ ID NOS: 29, 30, 31, as shown in figures 4 and 5. The capsid protein can have about 70% homology, about 75% homology, 80% homology, 85% homology, 90% homology, 95% homology, 98% homology, 99% homology, or even
10 100% homology to the protein having the amino acid sequence encoded by the nucleotides set forth in SEQ ID NOS: 29, 30, or 31. The percent homology used to identify proteins herein, can be based on a nucleotide-by-nucleotide comparison or more preferable is based on a computerized algorithm as described herein. Variations in the amino acid sequence of the AAV5 capsid protein are contemplated herein, as long as the resulting particle
15 comprising an AAV5 capsid protein remains antigenically or immunologically distinct from AAV1, AAV2, AAV3, AAV4 or AAV6 capsid, as can be routinely determined by standard methods. Specifically, for example, ELISA and Western blots can be used to determine whether a viral particle is antigenically or immunologically distinct from AAV2 or the other serotypes. Furthermore, the AAV5 particle preferably retains tissue tropism distinction
20 from AAV2, such as that exemplified in the examples herein. An AAV5 chimeric particle comprising at least one AAV5 coat protein may have a different tissue tropism from that of an AAV5 particle consisting only of AAV5 coat proteins, but is still distinct from the tropism of an AAV2 particle, in that it will infect some cells not infected by AAV2 or an AAV2 particle.

25 The invention further provides a recombinant AAV5 virion, comprising an AAV5 particle containing, i.e., encapsidating, a vector comprising a pair of AAV5 inverted terminal repeats. The recombinant vector can further comprise an AAV5 Rep-encoding nucleic acid. The vector encapsidated in the particle can further comprise an exogenous nucleic acid inserted between the inverted terminal repeats. AAV5 Rep confers targeted
30 integration and efficient replication, thus production of recombinant AAV5, comprising AAV5 Rep, yields more particles than production of recombinant AAV2. Since AAV5 is more efficient at replicating and packaging its genome, the exogenous nucleic acid inserted, or in the AAV5 capsids of the present invention, between the inverted terminal repeats can be packaged in the AAV1, 2, 3, 4, or 6 capsids to achieve the specific tissue tropism
35 conferred by the capsid proteins.

5 The invention further contemplates chimeric recombinant ITRs that contains a rep binding site and a TRS site recognized by that Rep protein. By "Rep protein" is meant all four of the Rep proteins, Rep 40, Rep 78, Rep 52, Rep 68. Alternatively, "Rep protein" could be one or more of the Rep proteins described herein. One example of a chimeric ITR would consist of an AAV5 D region (SEQ ID NO: 45), an AAV5 TRS site (SEQ ID NO: 10 43), an AAV2 hairpin and an AAV2 binding site. Another example would be an AAV5 D region, an AAV5 TRS site, an AAV3 hairpin and an AAV3 binding site. In these chimeric ITRs, the D region can be from AAV1, 2, 3, 4, 5 or 6. The hairpin can be derived from AAV 1,2 3, 4, 5, 6. The binding site can be derived from any of AAV1, 2, 3, 4, 5 or 6. Preferably, the D region and the TRS are from the same serotype.

15 The chimeric ITRs can be combined with AAV5 Rep protein and any of the AAV serotype capsids to obtain recombinant virion. For example, recombinant virion can be produced by an AAV5 D region, an AAV5 TRS site, an AAV2 hairpin, an AAV2 binding site, AAV5 Rep protein and AAV1 capsid. This recombinant virion would possess the cellular tropism conferred by the AAV1 capsid protein and would possess the efficient 20 replication conferred by the AAV5 Rep.

Other examples of the ITR, Rep protein and Capsids that will produce recombinant virion are provided in the list below:

5 ITR + 5Rep + 5Cap=virion
 5 ITR + 5Rep + 1Cap=virion
 25 5 ITR + 5Rep + 2Cap=virion
 5 ITR + 5Rep + 3Cap=virion
 5 ITR + 5Rep + 4Cap=virion
 5 ITR + 5Rep + 6Cap=virion
 1 ITR + 1Rep + 5Cap=virion
 30 2 ITR + 2Rep + 5Cap=virion
 3 ITR + 3Rep + 5Cap=virion
 4 ITR + 4Rep + 5Cap=virion
 6 ITR + 6Rep + 5Cap=virion

In any of the constructs described herein, inclusion of a promoter is preferred. As 35 used in the constructs herein, unless otherwise specified, Cap (capsid) refers to any of AAV5 VP1, AAV5 VP2, AAV5 VP3, combinations thereof, functional fragments of any of

5 VP1, VP2 or VP3, or chimeric capsids as described herein. The ITRs of the constructs described herein, can be chimeric recombinant ITRs as described elsewhere in the application.

Conjugates of recombinant or wild-type AAV5 virions and nucleic acids or proteins can be used to deliver those molecules to a cell. For example, the purified AAV5 can be
10 used as a vehicle for delivering DNA bound to the exterior of the virus. Examples of this are to conjugate the DNA to the virion by a bridge using poly L lysine or other charged molecule. Also contemplated are virosomes that contain AAV5 structural proteins (AAV5 capsid proteins), lipids such as DOTAP, and nucleic acids that are complexed via charge interaction to introduce DNA into cells.

15 Also provided by this invention are conjugates that utilize the AAV5 capsid or a unique region of the AAV5 capsid protein (e.g. VP1, VP2 or VP3 or combinations thereof) to introduce DNA into cells. For example, the type 5 VP3 protein or fragment thereof, can be conjugated to a DNA on a plasmid that is conjugated to a lipid. Cells can be infected using the targeting ability of the VP3 capsid protein to achieve the desired tissue tropism,
20 specific to AAV5. Type 5 VP1 and VP2 proteins can also be utilized to introduce DNA or other molecules into cells. By further incorporating the Rep protein and the AAV TRS into the DNA-containing conjugate, cells can be transduced and targeted integration can be achieved. For example, if AAV5 specific targeted integration is desired, a conjugate composed of the AAV5 VP3 capsid, AAV5 rep or a fragment of AAV5 rep, AAV5 TRS,
25 the rep binding site, the heterologous DNA of interest, and a lipid, can be utilized to achieve AAV5 specific tropism and AAV5 specific targeted integration in the genome.

Further provided by this invention are chimeric viruses where AAV5 can be combined with herpes virus, herpes virus amplicons, baculovirus or other viruses to achieve a desired tropism associated with another virus. For example, the AAV5 ITRs could be
30 inserted in the herpes virus and cells could be infected. Post-infection, the ITRs of AAV5 could be acted on by AAV5 rep provided in the system or in a separate vehicle to rescue AAV5 from the genome. Therefore, the cellular tropism of the herpes simplex virus can be combined with AAV5 rep mediated targeted integration. Other viruses that could be utilized to construct chimeric viruses include, lentivirus, retrovirus, pseudotyped retroviral vectors,
35 and adenoviral vectors.

5 The present invention further provides isolated nucleic acids of AAV5. For example, provided is an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 23 (AAV5 genome). This nucleic acid, or portions thereof, can be inserted into vectors, such as plasmids, yeast artificial chromosomes, or other viral vector (particle), if desired, by standard cloning methods. The present invention also provides an isolated nucleic acid
10 consisting essentially of the nucleotide sequence set forth in SEQ ID NO: 23. The nucleotides of SEQ ID NO: 23 can have minor modifications and still be contemplated by the present invention. For example, modifications that do not alter the amino acid encoded by any given codon (such as by modification of the third, "wobble," position in a codon) can readily be made, and such alterations are known in the art. Furthermore, modifications that
15 cause a resulting neutral (conserved) amino acid substitution of a similar amino acid can be made in a coding region of the genome. Additionally, modifications as described herein for the AAV5 components, such as the ITRs, the p5 promoter, etc. are contemplated in this invention. Furthermore, modifications to regions of SEQ ID NO: 23 other than in the ITR, TRS Rep binding site and hairpin are likely to be tolerated without serious impact on the
20 function of the nucleic acid as a recombinant vector.

 The present invention additionally provides an isolated nucleic acid that selectively hybridizes with any nucleic acid disclosed herein, including the entire AAV5 genome and any unique fragment thereof, including the Rep and capsid encoding sequences (e.g. SEQ ID NOS: 23, 29, 30, 31, 32, 33, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, and 45). Specifically, the
25 nucleic acid can selectively or specifically hybridize to an isolated nucleic acid consisting of the nucleotide sequence set forth in SEQ ID NO: 23 (AAV5 genome). The present invention further provides an isolated nucleic acid that selectively or specifically hybridizes with an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 23 (AAV5 genome). By "selectively hybridizes" as used herein is meant a nucleic acid that
30 hybridizes to one of the disclosed nucleic acids under sufficient stringency conditions without significant hybridization to a nucleic acid encoding an unrelated protein, and particularly, without detectably hybridizing to nucleic acids of AAV2. Thus, a nucleic acid that selectively hybridizes with a nucleic acid of the present invention will not selectively hybridize under stringent conditions with a nucleic acid encoding a different protein or the
35 corresponding protein from a different serotype of the virus, and vice versa. A "specifically hybridizing" nucleic acid is one that hybridizes under stringent conditions to only a nucleic

5 acid found in AAV5. Therefore, nucleic acids for use, for example, as primers and probes to detect or amplify the target nucleic acids are contemplated herein. Nucleic acid fragments that selectively hybridize to any given nucleic acid can be used, e.g., as primers and or probes for further hybridization or for amplification methods (e.g., polymerase chain reaction (PCR), ligase chain reaction (LCR)). Additionally, for example, a primer or probe
10 can be designed that selectively hybridizes with both AAV5 and a gene of interest carried within the AAV5 vector (i.e., a chimeric nucleic acid).

A nucleic acid that selectively hybridizes to any portion of the AAV5 genome is contemplated herein. Therefore, a nucleic acid that selectively hybridizes to AAV5 can be of longer length than the AAV5 genome, it can be about the same length as the AAV5 genome
15 or it can be shorter than the AAV5 genome. The length of the nucleic acid is limited on the shorter end of the size range only by its specificity for hybridization to AAV5, i.e., once it is too short, typically less than about 5 to 7 nucleotides in length, it will no longer bind specifically to AAV5, but rather will hybridize to numerous background nucleic acids. Additionally contemplated by this invention is a nucleic acid that has a portion that
20 specifically hybridizes to AAV5 and a portion that specifically hybridizes to a gene of interest inserted within AAV5.

The present invention further provides an isolated nucleic acid encoding an adeno-associated virus 5 Rep protein. The AAV5 Rep proteins are encoded by open reading frame (ORF) 1 of the AAV5 genome. Examples of the AAV5 Rep genes are shown in the nucleic
25 acid set forth in SEQ ID NO: 23, and include nucleic acids consisting essentially of the nucleotide sequences set forth in SEQ ID NOS: 32 (Rep52), 33 (Rep78), 35 (Rep40), and 37 (Rep68), and nucleic acids comprising the nucleotide sequences set forth in SEQ ID NOS: 32, 33, 35, and 37. However, the present invention contemplates that the Rep nucleic acid can include any one, two, three, or four of the four Rep proteins, in any order, in such a
30 nucleic acid. Furthermore, minor modifications are contemplated in the nucleic acid, such as silent mutations in the coding sequences, mutations that make neutral or conservative changes in the encoded amino acid sequence, and mutations in regulatory regions that do not disrupt the expression of the gene. Examples of other minor modifications are known in the art. Further modifications can be made in the nucleic acid, such as to disrupt or alter
35 expression of one or more of the Rep proteins in order to, for example, determine the effect of such a disruption; such as to mutate one or more of the Rep proteins to determine the

5 resulting effect, etc. However, in general, a modified nucleic acid encoding a Rep protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the Rep nucleic sequences described herein e.g., SEQ ID NOS: ~~11, 13 and 15~~ 32, 33, 35 and 37, and the Rep polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence
10 described herein, e.g., SEQ ID NOS: 24, 25, 34 and 36. Percent homology is determined by the techniques described herein.

The present invention also provides an isolated nucleic acid that selectively or specifically hybridizes with a nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NOS: 32, 33, 35 and 37, and an isolated nucleic acid that selectively
15 hybridizes with a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NOS: 32, 33, 35 and 37. "Selectively hybridizing" and "stringency of hybridization" is defined elsewhere herein.

As described above, provided is the nucleic acid encoding a Rep 40 protein and, in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID
20 NO: 35, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO: 35, and a nucleic acid encoding the adeno-associated virus 5 protein having the amino acid sequence set forth in SEQ ID NO: 34. The present invention also provides the nucleic acid encoding a Rep 52 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 32, an isolated nucleic acid consisting
25 essentially of the nucleotide sequence set forth in SEQ ID NO: 32, and a nucleic acid encoding the adeno-associated virus 5 Rep protein having the amino acid sequence set forth in SEQ ID NO: 24. The present invention further provides the nucleic acid encoding a Rep 68 protein and, in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 37, an isolated nucleic acid consisting essentially of the nucleotide
30 sequence set forth in SEQ ID NO: 37, and a nucleic acid encoding the adeno-associated virus 5 protein having the amino acid sequence set forth in SEQ ID NO: 36. And, further, provided is the nucleic acid encoding a Rep 78 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 33, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO: 33, and a
35 nucleic acid encoding the adeno-associated virus 5 Rep protein having the amino acid sequence set forth in SEQ ID NO: 25. As described elsewhere herein, these nucleic acids

5 can have minor modifications, including silent nucleotide substitutions, mutations causing conservative amino acid substitutions in the encoded proteins, and mutations in control regions that do not or minimally affect the encoded amino acid sequence.

The present invention further provides a nucleic acid encoding the entire AAV5 Capsid polypeptide. Furthermore, provided is a nucleic acid encoding each of the three
10 AAV5 coat proteins, VP1, VP2, and VP3. Thus, provided is a nucleic acid encoding AAV5 VP1, a nucleic acid encoding AAV5 VP2, and a nucleic acid encoding AAV5 VP3. Thus, provided is a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO: 26 (VP1); a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO: 27 (VP2), and a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO: 28 (VP3).
15 The present invention also specifically provides a nucleic acid comprising SEQ ID NO: 29 (VP1 gene); a nucleic acid comprising SEQ ID NO: 30 (VP2 gene); and a nucleic acid comprising SEQ ID NO: 31 (VP3 gene). The present invention also specifically provides a nucleic acid consisting essentially of SEQ ID NO: 29 (VP1 gene), a nucleic acid consisting essentially of SEQ ID NO: 30 (VP2 gene), and a nucleic acid consisting essentially of SEQ
20 ID NO: 31 (VP3 gene). Minor modifications in the nucleotide sequences encoding the capsid, or coat, proteins are contemplated, as described above for other AAV5 nucleic acids. However, in general, a modified nucleic acid encoding a capsid protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the capsid nucleic sequences described herein e.g., SEQ ID NOS: 29, 30 and 31, and the capsid polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about
25 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NOS: 26, 27, and 28. Nucleic acids that selectively hybridize with the nucleic acids of SEQ ID NOS: 29, 30, and 31 under the conditions described above are also provided.

Provided is an isolated AAV5 Rep protein. An AAV5 Rep polypeptide is encoded
30 by ORF1 of AAV5. The present invention also provides each individual AAV5 Rep protein. Thus provided is AAV5 Rep 40 (e.g., SEQ ID NO: 34), or a unique fragment thereof. Provided is AAV5 Rep 52 (e.g., SEQ ID NO: 24), or a unique fragment thereof. Provided is AAV5 Rep 68 (e.g., SEQ ID NO: 36), or a unique fragment thereof. Provided is an example of AAV5 Rep 78 (e.g., SEQ ID NO: 25), or a unique fragment thereof. By "unique fragment
35 thereof" is meant any smaller polypeptide fragment encoded by an AAV5 rep gene that is of sufficient length to be found only in the Rep polypeptide. Substitutions and modifications of

5 the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide.

The present invention further provides an AAV5 Capsid polypeptide or a unique fragment thereof. AAV5 capsid polypeptide is encoded by ORF 2 of AAV5. The present invention further provides the individual AAV5 capsid proteins, VP1, VP2 and VP3 or
10 unique fragments thereof. Thus, provided is an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO: 26 (VP1). The present invention additionally provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO: 27 (VP2). The present invention also provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO: 28 (VP3). By "unique fragment thereof" is meant any smaller
15 polypeptide fragment encoded by any AAV5 capsid gene that is of sufficient length to be found only in the AAV5 capsid protein. Substitutions and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide. However, an AAV5 Capsid polypeptide including all three coat proteins will have greater than about 56% overall
20 homology to the polypeptide encoded by the nucleotides set forth in SEQ ID NOS: 26, 27, or 28. The protein can have about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, 93%, 95%, 97% or even 100% homology to the amino acid sequence encoded by the nucleotides set forth in SEQ ID NOS: 26, 27 or 28. An AAV5 VP1 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or
25 about 100% homology to the amino acid sequence set forth in SEQ ID NO: 26. An AAV5 VP2 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO: 27. An AAV5 VP3 polypeptide can have at least about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set
30 forth in SEQ ID NO: 28.

The AAV ITRs in the vector for the herein described delivery methods can be AAV5 ITRs (SEQ ID NOS: 41 and 42). Furthermore, the AAV ITRs in the vector for the herein described nucleic acid delivery methods can also comprise AAV1, AAV2, AAV3, AAV4, or AAV6 inverted terminal repeats.

5 Compositions and methods for making BAAV vectors

Compositions and methods for making and using BAAV vectors have been previously described in U.S. Patent Application No. 60/526786, filed December 4, 2003, and in International Patent Application No. PCT/US04/40825, filed December 6, 2004, which are hereby incorporated herein by reference for this teaching.

10 Provided is a recombinant bovine adeno-associated virus (BAAV). This virus has one or more of the characteristics described below. The compositions of the present invention do not include wild-type BAAV. The methods of the present invention can use either wild-type BAAV or recombinant BAAV-based delivery.

Provided are novel BAAV particles, recombinant BAAV vectors and recombinant
15 BAAV virions. An BAAV particle is a viral particle comprising an BAAV capsid protein. A recombinant BAAV vector is a nucleic acid construct that comprises at least one unique nucleic acid of BAAV. A recombinant BAAV virion is a particle containing a recombinant BAAV vector, wherein the particle can be either an BAAV particle as described herein or a non-BAAV particle. Alternatively, the recombinant BAAV virion is an BAAV particle
20 containing a recombinant vector, wherein the vector can be either an BAAV vector as described herein or a non-BAAV vector. These vectors, particles, virions, nucleic acids and polypeptides are described below.

Provided is the nucleotide sequence of the BAAV genome and vectors and particles derived therefrom. Specifically, provided is a nucleic acid vector comprising a pair of
25 BAAV inverted terminal repeats (ITRs) and a promoter between the inverted terminal repeats. The rep proteins of AAV5 and BAAV will bind to the BAAV ITR and it can function as an origin of replication for packaging of recombinant AAV particles. The minimum sequence necessary for this activity is the TRS site where Rep cleaves in order to replicate the virus. Minor modifications in an ITR are contemplated and are those that will
30 not interfere with the hairpin structure formed by the ITR as described herein and known in the art. Furthermore, to be considered within the term e.g. it must retain the Rep binding site described herein.

The D- region of the AAV2 ITR, a single stranded region of the ITR, inboard of the TRS site, has been shown to bind a factor which depending on its phosphorylation state
35 correlates with the conversion of the AAV from a single stranded genome to a transcriptionally active form that allows for expression of the viral DNA. This region is

5 conserved between AAV2, 3, 4, and 6 but is divergent in AAV5 and BAAV (SEQ ID NO: 59). The D+ region is the reverse complement of the D- region.

The promoter can be any desired promoter, selected by known considerations, such as the level of expression of a nucleic acid functionally linked to the promoter and the cell type in which the vector is to be used. That is, the promoter can be tissue/cell-specific.

10 Promoters can be prokaryotic, eukaryotic, fungal, nuclear, mitochondrial, viral or plant promoters. Promoters can be exogenous or endogenous to the cell type being transduced by the vector. Promoters can include, for example, bacterial promoters, known strong promoters such as SV40 or the inducible metallothionein promoter, or an AAV promoter, such as an AAV p5 promoter. Additionally, chimeric regulatory promoters for targeted gene

15 expression can be utilized. Examples of these regulatory systems, which are known in the art, include the tetracycline based regulatory system which utilizes the tet transactivator protein (tTA), a chimeric protein containing the VP16 activation domain fused to the tet repressor of *Escherichia coli*, the IPTG based regulatory system, the CID based regulatory system, and the Ecdysone based regulatory system. Other promoters include promoters

20 derived from actin genes, immunoglobulin genes, cytomegalovirus (CMV), adenovirus, bovine papilloma virus, adenoviral promoters, such as the adenoviral major late promoter, an inducible heat shock promoter, respiratory syncytial virus, Rous sarcoma virus (RSV), etc., specifically, the promoter can be AAV2 p5 promoter or AAV5 p5 promoter or BAAV p5 promoter. More specifically, the BAAV p5 promoter can be in about the same location in

25 SEQ ID NO: 47 as the AAV2 p5 promoter, in the corresponding AAV2 published sequence. Additionally, the p5 promoter may be enhanced by nucleotides 1-173 of SEQ ID NO: 47. Furthermore, smaller fragments of p5 promoter that retain promoter activity can readily be determined by standard procedures including, for example, constructing a series of deletions in the p5 promoter, linking the deletion to a reporter gene, and determining whether the

30 reporter gene is expressed, i.e., transcribed and/or translated. The promoter can be the promoter of any of the AAV serotypes, and can be the p19 promoter (SEQ ID NO: 62) or the p40 promoter set forth in the sequence listing as SEQ ID NO: 63.

It should be recognized that any errors in any of the nucleotide sequences disclosed herein can be corrected, for example, by using the hybridization procedure described below

35 with various probes derived from the described sequences such that the coding sequence can be reisolated and resequenced. Rapid screening for point mutations can also be achieved

5 with the use of polymerase chain reaction single strand conformation polymorphism (PCR SSCP). The corresponding amino acid sequence can then be corrected accordingly.

The BAAV-derived vector can include any normally occurring BAAV nucleic acid sequences in addition to an ITR and promoter. The BAAV-derived vector can also include sequences that are at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to
10 the BAAV nucleic acids set forth herein. Examples of vector constructs are provided below.

The present vector or BAAV particle or recombinant BAAV virion can utilize any unique fragment of these present BAAV nucleic acids, including the BAAV nucleic acids set forth in SEQ ID NOS: 47, 48, 50, 52, 54, 56 and 58-63. To be unique, the fragment must be of sufficient size to distinguish it from other known sequences, most readily determined
15 by comparing any nucleic acid fragment to the nucleotide sequences of nucleic acids in computer databases, such as GenBank. Such comparative searches are standard in the art. Typically, a unique fragment useful as a primer or probe will be at least about 8 or 10, preferable at least 20 or 25 nucleotides in length, depending upon the specific nucleotide content of the sequence. Additionally, fragments can be, for example, at least about 30, 40,
20 50, 75, 100, 200 or 500 nucleotides in length and can encode polypeptides or be probes. The nucleic acid can be single or double stranded, depending upon the purpose for which it is intended. Where desired, the nucleic acid can be RNA.

The present invention further provides a BAAV capsid protein to contain the vector. In particular, provided is not only a polypeptide comprising all three BAAV coat proteins,
25 i.e., VP1, VP2 and VP3, but also a polypeptide comprising each BAAV coat protein individually, SEQ ID NOS: 53, 55, and 57, respectively. Thus, an BAAV particle comprising an BAAV capsid protein comprises at least one BAAV coat protein VP1, VP2 or VP3. A BAAV particle comprising an BAAV capsid protein can be utilized to deliver a nucleic acid vector to a cell, tissue or subject. For example, the herein described BAAV
30 vectors can be encapsidated in an AAV5 capsid-derived particle and utilized in a gene delivery method. Furthermore, other viral nucleic acids can be encapsidated in the BAAV particle and utilized in such delivery methods. For example, an AAV1-8 or AAV vector (e.g. AAV1-8 or AAV ITR and nucleic acid of interest) can be encapsidated in an BAAV particle and administered. Furthermore, a BAAV chimeric capsid incorporating both AAV1-
35 8 or AAV capsid and BAAV capsid sequences can be generated, by standard cloning methods, selecting regions from the known sequences of each protein as desired. For

5 example, particularly antigenic regions of the BAAV capsid protein can be replaced with the corresponding region of the BAAV capsid protein. In addition to chimeric capsids incorporating AAV2 capsid sequences, chimeric capsids incorporating AAV1, 3-8, and AAV5 capsid sequences can be generated, by standard cloning methods, selecting regions from the known sequences of each protein as desired. Alternatively a chimeric capsid can be made by the addition of a plasmid that expresses AAV1-8 capsid proteins at a ratio with the BAAV capsid expression plasmid that allows only a few capsid proteins to be incorporated into the BAAV particle. Thus, for example, a chimeric particle may be constructed that contains 6 AAV2 capsid proteins and 54 BAAV capsid proteins if the complete capsid contains 60 capsid proteins.

15 The capsids can also be modified to alter their specific tropism by genetically altering the capsid to encode a specific ligand to a cell surface receptor. Alternatively, the capsid can be chemically modified by conjugating a ligand to a cell surface receptor. By genetically or chemically altering the capsids, the tropism can be modified to direct BAAV to a particular cell or population of cells. The capsids can also be altered immunologically by conjugating the capsid to an antibody that recognizes a specific protein on the target cell or population of cells.

It has been recently reported that insertion of foreign epitopes (RGD motif, LH receptor targeting epitope) in certain regions of AAV2 capsid can redirect viral tropism. However, AAV2 naturally infects a wide variety of cell types and complete retargeting of rAAV2 would be difficult to achieve. Provided are two regions in the capsid of BAAV that are on the virus surface and could tolerate substitution. These two regions are aa 257-264 (GSSNASDT, SEQ ID NO:67) and aa 444-457 (TTSGGTLNQGNSAT, SEQ ID NO:68). Other regions of the BAAV capsid could also accommodate the substitution of amino acids that would allow for epitope presentation on the surface of the virus. All of these regions would have in common 1) Surface exposure 2) able to support a substitution of sequence to insert the epitope 3) still allow for capsid assembly.

Because of the symmetry of the AAV particles, a substitution in one subunit of the capsid will appear multiple times on the capsid surface. For example the capsid is made of approximately 55 VP3 proteins. Therefore an epitope incorporated in the VP3 protein could be expressed 55 times on the surface of each particle increasing the likelihood of the epitope forming a stable interaction with its target. In some cases this may be too high of a ligand

5 density for functional binding or this high density of epitope may interfere with capsid formation. The epitope density could be lowered by introducing another plasmid into the packaging system for production of recombinant particles and the ratio between the packaging plasmid with the modified VP3 protein and the wt VP3 protein altered to balance the epitope density on the virus surface.

10 Epitopes could be incorporated into the virus capsid for the purpose of 1) altering the tropism of the virus 2) blocking an immune response directed at the virus 3) developing a host immune response to the epitope for the purpose of vaccination.

Examples of epitopes that could be added to BAAV capsids include but are not limited to:

15 LH receptor binding epitope
RGD integrin binding epitope
CD13 binding epitope NGRAHA SEQ ID NO:69
The Retanef polyprotein vaccine candidate for HIV-1
single chain antibody fragments directed against tumor cells
20 Endothelial cell binding epitope SIGYPLP SEQ ID NO:70
serpin receptor ligand, KFNKPFVFLI SEQ ID NO:71
protective B-cell epitope hemagglutinin (HA) 91-108 from influenza HA
NDV B-cell immunodominant epitope (IDE) spanning residues 447 to 455
Major immunogenic epitope for parvovirus B19 (NISLDNPLENPSSLFDLVARIK,
25 SEQ ID NO:72) that can elicit protective antibody titers.

The capsids can also be assembled into empty particles by expression in mammalian, bacterial, fungal or insect cells. For example, AAV2 particles are known to be made from VP3 and VP2 capsid proteins in baculovirus. The same basic protocol can produce an empty BAAV particle comprising BAAV capsid proteins and also full particles.

30 The herein described recombinant BAAV nucleic acid derived vector can be encapsidated in an AAV particle. In particular, it can be encapsidated in an AAV1 particle, an AAV2 particle, an AAV3 particle, an AAV4 particle, an AAV5 particle or an AAV6 or AAV7 or an AAV8 or AAV particle, a portion of any of these capsids, or a chimeric capsid particle as described above, by standard methods using the appropriate capsid
35 proteins in the encapsidation process, as long as the nucleic acid vector fits within the size limitation of the particle utilized. The encapsidation process itself is standard in the art. The

5 BAAV replication machinery, i.e. the rep initiator proteins and other functions required for replication, can be utilized to produce the BAAV genome that can be packaged in an AAV1-8 or AAV capsid.

The recombinant BAAV virion containing a vector can also be produced by recombinant methods utilizing multiple plasmids. In one example, the BAAV rep nucleic acid would be cloned into one plasmid, the BAAV ITR nucleic acid would be cloned into
10 another plasmid and the AAV1-8 capsid nucleic acid would be cloned on another plasmid. These plasmids would then be introduced into cells. The cells that were efficiently transduced by all three plasmids, would exhibit specific integration as well as the ability to produce BAAV recombinant virus. Additionally, two plasmids could be used where the
15 BAAV rep nucleic acid would be cloned into one plasmid and the BAAV ITR and BAAV capsid would be cloned into another plasmid. These plasmids would then be introduced into cells. The cells that were efficiently transduced by both plasmids, would exhibit specific integration as well as the ability to produce BAAV recombinant virus.

An BAAV capsid polypeptide encoding the entire VP1, VP2, and VP3 polypeptide
20 can overall have greater than 56% homology to the polypeptide having the amino acid sequence encoded by nucleotides in SEQ ID NOS: 52, 54 and 56. The capsid protein can have about 70% homology, about 75% homology, 80% homology, 85% homology, 90% homology, 95% homology, 98% homology, 99% homology, or even 100% homology to the protein having the amino acid sequence encoded by the nucleotides set forth in SEQ ID
25 NOS: 52, 54 and 56. The percent homology used to identify proteins herein, can be based on a nucleotide-by-nucleotide comparison or more preferable is based on a computerized algorithm as described herein. Variations in the amino acid sequence of the BAAV capsid protein are contemplated herein, as long as the resulting particle comprising an BAAV capsid protein remains antigenically or immunologically distinct from AAV1-8 or AAV
30 capsid, as can be routinely determined by standard methods. Specifically, for example, ELISA and Western blots can be used to determine whether a viral particle is antigenically or immunologically distinct from AAV2 or the other serotypes. Furthermore, the BAAV particle preferably retains tissue tropism distinction from other AAVs, such as that exemplified in the examples herein. A BAAV chimeric particle comprising at least one
35 BAAV coat protein may have a different tissue tropism from that of an BAAV particle

5 consisting only of BAAV coat proteins, but is still distinct from the tropism of an AAV2 particle.

The invention further provides a recombinant BAAV virion, comprising a BAAV particle containing, i.e., encapsidating, a vector comprising a pair of BAAV inverted terminal repeats. The recombinant vector can further comprise a BAAV Rep-encoding
 10 nucleic acid. The vector encapsidated in the particle can further comprise an exogenous nucleic acid inserted between the inverted terminal repeats.

The invention further contemplates chimeric recombinant ITRs that contain a rep binding site and a TRS site recognized by that Rep protein. By "Rep protein" is meant all four of the Rep proteins, Rep 40, Rep 78, Rep 52, Rep 68. Alternatively, "Rep protein"
 15 could be one or more of the Rep proteins described herein. One example of a chimeric ITR would consist of an BAAV D region (SEQ ID NO: 59), an BAAV TRS site (SEQ ID NO: 60), an AAV2 hairpin and an AAV2 Rep binding site. Another example would be a BAAV D region, an BAAV TRS site, an AAV3 hairpin and an AAV3 Rep binding site. In these chimeric ITRs, the D region can be from AAV1-8 or AAV. The hairpin can be derived
 20 from AAV 1-8 or AAV. The binding site can be derived from any of AAV1-8 or AAV. Preferably, the D region and the TRS are from the same serotype.

The chimeric ITRs can be combined with BAAV Rep protein and any of the AAV serotype capsids to obtain recombinant virion. For example, recombinant virion can be produced by a BAAV D region, an BAAV TRS site, an AAV2 hairpin, an AAV2 binding
 25 site, BAAV Rep protein and AAV1 capsid. This recombinant virion would possess the cellular tropism conferred by the AAV1 capsid protein and would possess the efficient replication conferred by the BAAV Rep.

Other examples of the ITR, Rep protein and Capsids that will produce recombinant virus are provided in the list below but not limited to :

30 BAAV ITR + BAAV Rep + BAAV Cap=virus
 AAV5 ITR + BAAV Rep + BAAV Cap=virus
 AAV5 ITR + BAAV Rep + AAV1 Cap=virus
 AAV5 ITR + BAAV Rep + AAV2 Cap=virus
 AAV5 ITR + BAAV Rep + AAV3 Cap=virus
 35 AAV5 ITR + BAAV Rep + AAV4 Cap=virus
 AAV5 ITR + BAAV Rep + AAV5 Cap=virus

- 5 AAV5 ITR + BAAV Rep + AAV6 Cap=virus
 AAV5 ITR + BAAV Rep + AAV7 Cap=virus
 AAV5 ITR + BAAV Rep + AAV8 Cap=virus
 BAAV ITR + AAV5 Rep + BAAV Cap=virus
 BAAV ITR + AAV5 Rep + AAV1 Cap=virus
 10 BAAV ITR + AAV5 Rep + AAV2 Cap=virus
 BAAV ITR + AAV5 Rep + AAV3 Cap=virus
 BAAV ITR + AAV5 Rep + AAV4 Cap=virus
 BAAV ITR + AAV5 Rep + AAV5 Cap=virus
 BAAV ITR + AAV5 Rep + AAV6 Cap=virus
 15 BAAV ITR + AAV5 Rep + AAV7 Cap=virus
 BAAV ITR + AAV5 Rep + AAV8 Cap=virus
 AAV5 ITR + AAV5 Rep + BAAV Cap=virus
 AAV1 ITR + AAV1 Rep + BAAV Cap=virus
 AAV2 ITR + AAV2 Rep + BAAV Cap=virus
 20 AAV3 ITR + AAV3 Rep + BAAV Cap=virus
 AAV4 ITR + AAV4 Rep + BAAV Cap=virus
 AAV5 ITR + AAV5 Rep + BAAV Cap=virus
 AAV6 ITR + AAV6 Rep + BAAV Cap=virus
 AAV7 ITR + AAV7 Rep + BAAV Cap=virus
 25 AAV8 ITR + AAV8 Rep + BAAV Cap=virus

In any of the constructs described herein, inclusion of a promoter is preferred. As used in the constructs herein, unless otherwise specified, Cap (capsid) refers to any of BAAV VP1, BAAV VP2, BAAV VP3, combinations thereof, functional fragments of any of VP1, VP2 or VP3, or chimeric capsids as described herein. The ITRs of the constructs
 30 described herein, can be chimeric recombinant ITRs as described elsewhere in the application.

Conjugates of recombinant or wild-type BAAV virions and nucleic acids or proteins can be used to deliver those molecules to a cell. For example, the purified BAAV can be used as a vehicle for delivering DNA bound to the exterior of the virus. Examples of this are
 35 to conjugate the DNA to the virion by a bridge using poly L lysine or other charged molecule. Also contemplated are virosomes that contain BAAV structural proteins (BAAV

5 capsid proteins), lipids such as DOTAP, and nucleic acids that are complexed via charge interaction to introduce DNA into cells.

Also provided by this invention are conjugates that utilize the BAAV capsid or a unique region of the BAAV capsid protein (e.g. VP1, VP2 or VP3 or combinations thereof) to introduce DNA into cells. For example, the BAAV VP3 protein or fragment thereof, can be conjugated to a DNA on a plasmid that is conjugated to a lipid. Cells can be infected using the targeting ability of the VP3 capsid protein to achieve the desired tissue tropism, specific to BAAV. BAAV VP1 and VP2 proteins can also be utilized to introduce DNA or other molecules into cells. By further incorporating the Rep protein and the AAV TRS into the DNA-containing conjugate, cells can be transduced and targeted integration can be achieved. For example, if BAAV specific targeted integration is desired, a conjugate composed of the BAAV VP3 capsid, BAAV rep or a fragment of BAAV rep, BAAV TRS, the rep binding site, the heterologous DNA of interest, and a lipid, can be utilized to achieve BAAV specific tropism and BAAV specific targeted integration in the genome.

Further provided by this invention are chimeric viruses where BAAV can be combined with herpes virus, baculovirus or other viruses to achieve a desired tropism associated with another virus. For example, the BAAV ITRs could be inserted in the herpes virus and cells could be infected. Post-infection, the ITRs of BAAV could be acted on by BAAV rep provided in the system or in a separate vehicle to rescue BAAV from the genome. Therefore, the cellular tropism of the herpes simplex virus can be combined with BAAV rep mediated targeted integration. Other viruses that could be utilized to construct chimeric viruses include lentivirus, retrovirus, pseudotyped retroviral vectors, and adenoviral vectors.

The present invention further provides isolated nucleic acids of BAAV. For example, provided is an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 47 (BAAV genome). This nucleic acid, or portions thereof, can be inserted into vectors, such as plasmids, yeast artificial chromosomes, or other viral vector (particle), if desired, by standard cloning methods. The present invention also provides an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO: 47. The nucleotides of SEQ ID NO: 47 can have minor modifications and still be contemplated by the present invention. For example, modifications that do not alter the amino acid encoded by any given codon (such as by modification of the third, "wobble," position in a

5 codon) can readily be made, and such alterations are known in the art. Furthermore, modifications that cause a resulting neutral (conserved) amino acid substitution of a similar amino acid can be made in a coding region of the genome. Additionally, modifications as described herein for the BAAV components, such as the ITRs, the p5 promoter, etc. are contemplated in this invention. Furthermore, modifications to regions of SEQ ID NO: 1 47
10 other than in the ITR, TRS, Rep binding site and hairpin are likely to be tolerated without serious impact on the function of the nucleic acid as a recombinant vector.

The present invention additionally provides an isolated nucleic acid that selectively hybridizes with any nucleic acid disclosed herein, including the entire BAAV genome and any unique fragment thereof, including the Rep and capsid encoding sequences (e.g. SEQ ID
15 NOS: 47, 48, 50, 52, 54, 56, 58, 59, 60, 61, 62, 63). Specifically, the nucleic acid can selectively or specifically hybridize to an isolated nucleic acid consisting of the nucleotide sequence set forth in SEQ ID NO: 47 (BAAV genome). The present invention further provides an isolated nucleic acid that selectively or specifically hybridizes with an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 47 (BAAV
20 genome). By "selectively hybridizes" as used herein is meant a nucleic acid that hybridizes to one of the disclosed nucleic acids under sufficient stringency conditions without significant hybridization to a nucleic acid encoding an unrelated protein, and particularly, without detectably hybridizing to nucleic acids of AAV2. Thus, a nucleic acid that selectively hybridizes with a nucleic acid of the present invention will not selectively
25 hybridize under stringent conditions with a nucleic acid encoding a different protein or the corresponding protein from a different serotype of the virus, and vice versa. A "specifically hybridizing" nucleic acid is one that hybridizes under stringent conditions to only a nucleic acid found in BAAV. Therefore, nucleic acids for use, for example, as primers and probes to detect or amplify the target nucleic acids are contemplated herein. Nucleic acid fragments
30 that selectively hybridize to any given nucleic acid can be used, e.g., as primers and or probes for further hybridization or for amplification methods (e.g., polymerase chain reaction (PCR), ligase chain reaction (LCR)). Additionally, for example, a primer or probe can be designed that selectively hybridizes with both BAAV and a gene of interest carried within the BAAV vector (i.e., a chimeric nucleic acid).

35 A nucleic acid that selectively hybridizes to any portion of the BAAV genome is contemplated herein. Therefore, a nucleic acid that selectively hybridizes to BAAV can be

5 of longer length than the BAAV genome, it can be about the same length as the BAAV genome or it can be shorter than the BAAV genome. The length of the nucleic acid is limited on the shorter end of the size range only by its specificity for hybridization to BAAV, i.e., once it is too short, typically less than about 5 to 7 nucleotides in length, it will no longer bind specifically to BAAV, but rather will hybridize to numerous background
10 nucleic acids. Additionally contemplated by this invention is a nucleic acid that has a portion that specifically hybridizes to BAAV and a portion that specifically hybridizes to a gene of interest inserted within BAAV.

The present invention further provides an isolated nucleic acid encoding a bovine adeno-associated virus Rep protein. The BAAV Rep proteins are encoded by open reading
15 frame (ORF) 1 of the BAAV genome. Examples of the BAAV Rep genes are shown in the nucleic acid set forth in SEQ ID NO: 47, and include nucleic acids consisting essentially of the nucleotide sequences set forth in SEQ ID NOS: 48 (rep78), 4(rep52) and nucleic acids comprising the nucleotide sequences set forth in SEQ ID NOS: 48 and 50. However, the present invention contemplates that the Rep nucleic acid can include any one, two, three, or
20 four of the four Rep proteins, in any order, in such a nucleic acid. Furthermore, minor modifications are contemplated in the nucleic acid, such as silent mutations in the coding sequences, mutations that make neutral or conservative changes in the encoded amino acid sequence, and mutations in regulatory regions that do not disrupt the expression of the gene. Examples of other minor modifications are known in the art. Further modifications can be
25 made in the nucleic acid, such as to disrupt or alter expression of one or more of the Rep proteins in order to, for example, determine the effect of such a disruption; such as to mutate one or more of the Rep proteins to determine the resulting effect, etc. However, in general, a modified nucleic acid encoding a Rep protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the Rep nucleic sequences
30 described herein e.g., SEQ ID NOS: 48 and 50, and the Rep polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NOS: 49 and 51. Percent homology is determined by the techniques described herein.

The present invention also provides an isolated nucleic acid that selectively or
35 specifically hybridizes with a nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NOS: 48 and 50, and an isolated nucleic acid that selectively hybridizes

5 with a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NOS: 48 and 50.
"Selectively hybridizing" and "stringency of hybridization" is defined elsewhere herein.

As described above, provided is the nucleic acid encoding a Rep 78 protein and, in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 48, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in
10 SEQ ID NO: 48, and a nucleic acid encoding the bovine adeno-associated virus protein having the amino acid sequence set forth in SEQ ID NO: 49. The present invention also provides the nucleic acid encoding a Rep 52 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 50, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO: 50, and a
15 nucleic acid encoding the bovine adeno-associated virus Rep 52 protein having the amino acid sequence set forth in SEQ ID NO: 51. As described elsewhere herein, these nucleic acids can have minor modifications, including silent nucleotide substitutions, mutations causing conservative amino acid substitutions in the encoded proteins, and mutations in control regions that do not or minimally affect the encoded amino acid sequence.

20 The present invention further provides a nucleic acid encoding the entire BAAV Capsid polypeptide. Furthermore, provided is a nucleic acid encoding each of the three BAAV coat proteins, VP1, VP2, and VP3. Thus, provided is a nucleic acid encoding BAAV VP1, a nucleic acid encoding BAAV VP2, and a nucleic acid encoding BAAV VP3. Thus, provided is a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO: 53
25 (VP1); a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO: 55 (VP2), and a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO: 57 (VP3). The present invention also specifically provides a nucleic acid comprising SEQ ID NO: 52 (VP1 gene); a nucleic acid comprising SEQ ID NO: 54 (VP2 gene); and a nucleic acid comprising SEQ ID NO: 56 (VP3 gene). The present invention also specifically provides a
30 nucleic acid consisting essentially of SEQ ID NO: 52 (VP1 gene), a nucleic acid consisting essentially of SEQ ID NO: 54 (VP2 gene), and a nucleic acid consisting essentially of SEQ ID NO: 56 (VP3 gene). Minor modifications in the nucleotide sequences encoding the capsid, or coat, proteins are contemplated, as described above for other BAAV nucleic acids. However, in general, a modified nucleic acid encoding a capsid protein will have at
35 least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the capsid nucleic sequences described herein e.g., SEQ ID NOS: 52, 54 and 56, and the capsid

5 polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NOS: 53, 55 and 57. Nucleic acids that selectively hybridize with the nucleic acids of SEQ ID NOS: 52, 54 and 56 under the conditions described above are also provided.

10 Provided is an isolated BAAV Rep protein. An BAAV Rep polypeptide is encoded by ORF1 of BAAV. The present invention also provides each individual BAAV Rep protein. Thus provided is BAAV Rep 52 (e.g., SEQ ID NO: 50), or a unique fragment thereof. Provided is BAAV Rep 78 (e.g., SEQ ID NO: 48), or a unique fragment thereof. By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by an BAAV rep gene that is of sufficient length to be found only in the Rep polypeptide. Substitutions
15 and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide.

The present invention further provides a BAAV Capsid polypeptide or a unique fragment thereof. BAAV capsid polypeptide is encoded by ORF 2 of BAAV. The present invention further provides the individual BAAV capsid proteins, VP1, VP2 and VP3 or
20 unique fragments thereof. Thus, provided is an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:52 (VP1). The present invention additionally provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO: 54 (VP2). The present invention also provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:56 (VP3). By "unique fragment thereof" is meant any smaller
25 polypeptide fragment encoded by any BAAV capsid gene that is of sufficient length to be found only in the BAAV capsid protein. Substitutions and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide. However, an BAAV Capsid polypeptide including all three coat proteins will have greater than about 56% overall
30 homology to the polypeptide encoded by the nucleotides set forth in SEQ ID NOS: 52, 54 or 56. The protein can have about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, 93%, 95%, 97% or even 100% homology to the amino acid sequence encoded by the nucleotides set forth in SEQ ID NOS: 52, 54 or 56. An BAAV VP1 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about
35 100% homology to the amino acid sequence set forth in SEQ ID NO: 53. An BAAV VP2 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%,

5 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO: 55. An BAAV VP3 polypeptide can have at least about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO: 57.

10 The present invention also provides a method of producing the BAAV virus by transducing a cell with the nucleic acid encoding the virus.

The present method further provides a method of delivering an exogenous (heterologous) nucleic acid to a cell comprising administering to the cell an BAAV particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the cell.

15 The AAV ITRs in the vector for the herein described delivery methods can be AAV ITRs (SEQ ID NOS: 58). Furthermore, the AAV ITRs in the vector for the herein described nucleic acid delivery methods can also comprise AAV1-8 or AAV inverted terminal repeats.

Compositions and methods for making AAV7 vectors

20 Compositions and methods for making and using AAV7 vectors have been previously described in Gao GP, et al. Proc Natl Acad Sci U S A. 2002 Sep 3;99(18):11854-9; U.S. Patent Application 2003/0228282; and International Patent Application No. PCT/US02/33630, which are hereby incorporated by reference herein for the teaching of compositions and method for making and using AAV7 virions, vectors, and particles.

25 Provided is a recombinant adeno-associated virus-7 (AAV7). This virus has one or more of the characteristics described below. The compositions of the present invention do not include wild-type AAV7. The methods of the present invention can use either wild-type AAV7 or recombinant AAV7-based delivery.

30 Provided are AAV7 particles, recombinant AAV7 vectors and recombinant AAV7 virions. An AAV7 particle is a viral particle comprising an AAV7 capsid protein. A recombinant AAV7 vector is a nucleic acid construct that comprises at least one unique nucleic acid of AAV7. A recombinant AAV7 virion is a particle containing a recombinant AAV7 vector, wherein the particle can be either an AAV7 particle as described herein or a non-AAV7 particle. Alternatively, the recombinant AAV7 virion is an AAV7 particle
35 containing a recombinant vector, wherein the vector can be either an AAV7 vector as

5 described herein or a non-AAV7 vector. These vectors, particles, virions, nucleic acids and polypeptides are described below.

The AAV7-derived vector can include any normally occurring AAV7 nucleic acid sequences. The AAV7-derived vector can also include sequences that are at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the AAV7 nucleic acids set forth
10 herein. Examples of vector constructs are provided below.

The present vector or AAV7 particle or recombinant AAV7 virion can utilize any unique fragment of the present AAV7 nucleic acids, including the AAV7 nucleic acids set forth in SEQ ID NO:64. Fragments can be, for example, at least about 30, 40, 50, 75, 100, 200 or 500 nucleotides in length. The nucleic acid can be single or double stranded,
15 depending upon the purpose for which it is intended.

The present invention further provides an AAV7 capsid protein to contain the vector. In particular, provided is a polypeptide comprising AAV7 capsid protein, SEQ ID NO:66. An AAV7 particle comprising an AAV7 capsid protein can be utilized to deliver a nucleic acid vector to a cell, tissue or subject. For example, the herein described AAV7 vectors can
20 be encapsidated in an AAV5 capsid-derived particle and utilized in a gene delivery method. Furthermore, other viral nucleic acids can be encapsidated in the AAV7 particle and utilized in such delivery methods. For example, an AAV1-6, 8, BAAV or AAV vector (e.g. AAV1-6, 8, BAAV or AAV ITR and nucleic acid of interest) can be encapsidated in an AAV7 particle and administered. Furthermore, a AAV7 chimeric capsid incorporating
25 AAV1-6, 8, BAAV or AAV capsid, and AAV7 capsid sequences can be generated, by standard cloning methods, selecting regions from the known sequences of each protein as desired. For example, particularly antigenic regions of the AAV2 capsid protein can be replaced with the corresponding region of the AAV7 capsid protein. In addition to chimeric capsids incorporating AAV2 capsid sequences, chimeric capsids incorporating AAV1, 3-6, 8, BAAV and AAV5 capsid sequences can be generated, by standard cloning methods,
30 selecting regions from the known sequences of each protein as desired. Alternatively a chimeric capsid can be made by the addition of a plasmid that expresses AAV1, 3-6, 8, BAAV or AAV5 capsid proteins at a ratio with the AAV7 capsid expression plasmid that allows only a few capsid proteins to be incorporated into the AAV7 particle. Thus, for
35 example, a chimeric particle may be constructed that contains 6 AAV2 capsid proteins and 54 AAV7 capsid proteins if the complete capsid contains 60 capsid proteins.

5 The capsids can also be assembled into empty particles by expression in mammalian, bacterial, fungal or insect cells. For example, AAV2 particles are known to be made from VP3 and VP2 capsid proteins in baculovirus. The same basic protocol can produce an empty AAV7 particle comprising AAV7 capsid proteins and also full particles.

 The herein described recombinant AAV7 nucleic acid derived vector can be
10 encapsidated in an AAV particle. In particular, it can be encapsidated in an AAV1 particle, an AAV2 particle, an AAV3 particle, an AAV4 particle, an AAV5 particle, an AAV6, an AAV8, a BAAV particle or AAV particle, a portion of any of these capsids, or a chimeric capsid particle as described above, by standard methods using the appropriate capsid proteins in the encapsidation process, as long as the nucleic acid vector fits within the size
15 limitation of the particle utilized. The encapsidation process itself is standard in the art. The AAV7 replication machinery, i.e. the rep initiator proteins and other functions required for replication, can be utilized to produce the AAV7 genome that can be packaged in an AAV1-6, 8, BAAV or AAV capsid.

 The recombinant AAV7 virion containing a vector can also be produced by
20 recombinant methods utilizing multiple plasmids. In one example, the AAV7 rep nucleic acid would be cloned into one plasmid, the AAV2 ITR nucleic acid would be cloned into another plasmid and the AAV7 capsid nucleic acid would be cloned on another plasmid. These plasmids would then be introduced into cells. The cells that were efficiently transduced by all three plasmids, would exhibit specific integration as well as the ability to
25 produce AAV7 recombinant virus. Additionally, two plasmids could be used where the AAV7 rep nucleic acid would be cloned into one plasmid and the AAV7 ITR and AAV7 capsid would be cloned into another plasmid. These plasmids would then be introduced into cells. The cells that were efficiently transduced by both plasmids, would exhibit specific integration as well as the ability to produce AAV7 recombinant virus.

30 An AAV7 capsid polypeptide encoding the entire VP1 polypeptide can overall have greater than 56% homology to the polypeptide having the amino acid sequence encoded by nucleotides in SEQ ID NO:66. The capsid protein can have about 70% homology, about 75% homology, 80% homology, 85% homology, 90% homology, 95% homology, 98% homology, 99% homology, or even 100% homology to the protein having the amino acid
35 sequence encoded by the nucleotides set forth in SEQ ID NO:66. The percent homology used to identify proteins herein, can be based on a nucleotide-by-nucleotide comparison or

5 more preferable is based on a computerized algorithm as described herein. Variations in the amino acid sequence of the AAV7 capsid protein are contemplated herein, as long as the resulting particle comprising an AAV7 capsid protein remains antigenically or immunologically distinct from AAV1-6, 8, BAAV or AAV capsid, as can be routinely determined by standard methods. Specifically, for example, ELISA and Western blots can
 10 be used to determine whether a viral particle is antigenically or immunologically distinct from AAV2 or the other serotypes. Furthermore, the AAV7 particle preferably retains tissue tropism distinction from other AAVs. An AAV7 chimeric particle comprising at least one AAV7 coat protein may have a different tissue tropism from that of an AAV7 particle consisting only of AAV7 coat proteins, but is still distinct from the tropism of an AAV2
 15 particle.

The invention further provides a recombinant AAV7 virion, comprising an AAV7 particle containing, i.e., encapsidating, a vector comprising a pair of AAV7 inverted terminal repeats. The recombinant vector can further comprise an AAV7 Rep-encoding nucleic acid. The vector encapsidated in the particle can further comprise an exogenous
 20 nucleic acid inserted between the inverted terminal repeats.

For example, recombinant virion can be produced by a AAV2 ITR, AAV2 Rep protein and AAV7 capsid. This recombinant virion would possess the cellular tropism conferred by the AAV7 capsid protein and would possess the efficient replication conferred by the AAV2 Rep.

25 Other examples of the ITR, Rep protein and Capsids that will produce recombinant virus are provided in the list below but not limited to :

AAV5 ITR + AAV7 Rep + AAV1 Cap=virus
 AAV5 ITR + AAV7 Rep + AAV2 Cap=virus
 AAV5 ITR + AAV7 Rep + AAV3 Cap=virus
 30 AAV5 ITR + AAV7 Rep + AAV4 Cap=virus
 AAV5 ITR + AAV7 Rep + AAV5 Cap=virus
 AAV5 ITR + AAV7 Rep + AAV6 Cap=virus
 AAV5 ITR + AAV7 Rep + AAV7 Cap=virus
 AAV5 ITR + AAV7 Rep + AAV8 Cap=virus
 35 AAV5 ITR + AAV7 Rep + BAAV Cap=virus
 AAV5 ITR + AAV7 Rep + AAV Cap=virus

5 AAV1 ITR + AAV1 Rep + AAV7 Cap=virus
 AAV2 ITR + AAV2 Rep + AAV7 Cap=virus
 AAV3 ITR + AAV3 Rep + AAV7 Cap=virus
 AAV4 ITR + AAV4 Rep + AAV7 Cap=virus
 AAV5 ITR + AAV5 Rep + AAV7 Cap=virus
 10 AAV6 ITR + AAV6 Rep + AAV7 Cap=virus
 AAV8 ITR + AAV8 Rep + AAV7 Cap=virus
 BAAV ITR + BAAV Rep + AAV7 Cap=virus
 AAAV ITR + AAAV Rep + AAV7 Cap=virus

15 In any of the constructs described herein, inclusion of a promoter is preferred. As
 used in the constructs herein, unless otherwise specified, Cap (capsid) refers to any of
 AAV7 VP1, AAV7 VP2, AAV7 VP3, combinations thereof, functional fragments of any of
 VP1, VP2 or VP3, or chimeric capsids as described herein. The ITRs of the constructs
 described herein, can be chimeric recombinant ITRs as described elsewhere in the
 20 application.

Conjugates of recombinant or wild-type AAV7 virions and nucleic acids or proteins
 can be used to deliver those molecules to a cell. For example, the purified AAV7 can be
 used as a vehicle for delivering DNA bound to the exterior of the virus. Examples of this are
 to conjugate the DNA to the virion by a bridge using poly L lysine or other charged
 25 molecule. Also contemplated are virosomes that contain AAV7 structural proteins (AAV7
 capsid proteins), lipids such as DOTAP, and nucleic acids that are complexed via charge
 interaction to introduce DNA into cells.

Also provided by this invention are conjugates that utilize the AAV7 capsid or a
 unique region of the AAV7 capsid protein (e.g. VP1, VP2 or VP3 or combinations thereof)
 30 to introduce DNA into cells. By "unique" is meant any smaller polypeptide fragment
 encoded by any AAV7 capsid gene that is of sufficient length to be unique to the AAV7
 Capsid protein. For example, the AAV7 VP1 protein or fragment thereof, can be conjugated
 to a DNA on a plasmid that is conjugated to a lipid. Cells can be infected using the targeting
 ability of the VP1 capsid protein to achieve the desired tissue tropism, specific to AAV7.
 35 AAV7 VP1 proteins can also be utilized to introduce DNA or other molecules into cells. By

5 further incorporating an AAV Rep protein and an AAV TRS into the DNA-containing conjugate, cells can be transduced and targeted integration can be achieved.

The present invention further provides isolated nucleic acids of AAV7. For example, provided is an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:64. This nucleic acid, or portions thereof, can be inserted into vectors, such as plasmids, yeast artificial chromosomes, or other viral vector (particle), if desired, by standard cloning methods. The present invention also provides an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:64. The nucleotides of SEQ ID NO:64 can have minor modifications and still be contemplated by the present invention. For example, modifications that do not alter the amino acid encoded by any given codon (such as by modification of the third, "wobble," position in a codon) can readily be made, and such alterations are known in the art. Furthermore, modifications that cause a resulting neutral (conserved) amino acid substitution of a similar amino acid can be made in a coding region of the genome.

The present invention also provides an isolated nucleic acid that selectively or specifically hybridizes with a nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:64, and an isolated nucleic acid that selectively hybridizes with a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:64. "Selectively hybridizing" and "stringency of hybridization" is defined elsewhere herein.

The present invention further provides an isolated nucleic acid encoding a AAV7 Rep protein. The AAV7 Rep proteins are encoded by open reading frame (ORF) 1 of the AAV7 genome. Examples of the AAV7 Rep genes are shown in the nucleic acid set forth in nucleotides 334-2205 of SEQ ID NO:64, and include nucleic acids consisting essentially of the nucleotide sequences set forth in 334-2205 of SEQ ID NO:64 (rep78). Minor modifications are contemplated in the nucleic acid, such as silent mutations in the coding sequences, mutations that make neutral or conservative changes in the encoded amino acid sequence, and mutations in regulatory regions that do not disrupt the expression of the gene. Examples of other minor modifications are known in the art. Further modifications can be made in the nucleic acid, such as to disrupt or alter expression of one or more of the Rep proteins in order to, for example, determine the effect of such a disruption; such as to mutate one or more of the Rep proteins to determine the resulting effect, etc. However, in general, a modified nucleic acid encoding a Rep protein will have at least about 85%, about 90%,

5 about 93%, about 95%, about 98% or 100% homology to the Rep nucleic sequences described herein e.g., SEQ ID NOS:65, and the Rep polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described in SEQ ID NO:65. Percent homology is determined by the techniques described herein.

10 The present invention further provides a nucleic acid encoding the entire AAV7 Capsid polypeptide. Thus, provided is a nucleic acid encoding the amino acid sequence set forth in nucleotides 2222-4435 of SEQ ID NO:64 (VP1). Minor modifications in the nucleotide sequences encoding the capsid, or coat, proteins are contemplated, as described above for other AAV7 nucleic acids. However, in general, a modified nucleic acid encoding
15 a capsid protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the capsid nucleic sequences described herein e.g., nucleotides 2222-4435 of SEQ ID NO:64, and the capsid polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NO:66.

20 AAV Vector Generation

It is understood that as discussed herein the use of the terms "homology" and "identity" mean the same thing as similarity. Thus, for example, if the use of the word homology is used to refer to two non-natural sequences, it is understood that this is not necessarily indicating an evolutionary relationship between these two sequences, but rather
25 is looking at the similarity or relatedness between their nucleic acid sequences. Many of the methods for determining homology between two evolutionarily related molecules are routinely applied to any two or more nucleic acids or proteins for the purpose of measuring sequence similarity regardless of whether they are evolutionarily related.

In general, it is understood that one way to define any known variants and
30 derivatives or those that might arise, of the disclosed nucleic acids and polypeptides herein, is through defining the variants and derivatives in terms of homology to specific known sequences. In general, variants of nucleic acids and polypeptides herein disclosed typically have at least, about 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent homology to the stated sequence or the
35 native sequence. Those of skill in the art readily understand how to determine the homology

5 of two polypeptides or nucleic acids. For example, the homology can be calculated after aligning the two sequences so that the homology is at its highest level.

Another way of calculating homology can be performed by published algorithms. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman Adv. Appl. Math. 2: 482 (1981), by the homology
10 alignment algorithm of Needleman and Wunsch, J. Mol Biol. 48: 443 (1970), by the search for similarity method of Pearson and Lipman, Proc. Natl. Acad. Sci. U.S.A. 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI; the BLAST algorithm of Tatusova and Madden FEMS
15 Microbiol. Lett. 174: 247-250 (1999) available from the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/blast/bl2seq/bl2.html>)), or by inspection.

The same types of homology can be obtained for nucleic acids by for example the algorithms disclosed in Zuker, M. Science 244:48-52, 1989, Jaeger et al. Proc. Natl. Acad. Sci. USA 86:7706-7710, 1989, Jaeger et al. Methods Enzymol. 183:281-306, 1989 which
20 are herein incorporated by reference for at least material related to nucleic acid alignment. It is understood that any of the methods typically can be used and that in certain instances the results of these various methods may differ, but the skilled artisan understands if identity is found with at least one of these methods, the sequences would be said to have the stated identity.

25 For example, as used herein, a sequence recited as having a particular percent homology to another sequence refers to sequences that have the recited homology as calculated by any one or more of the calculation methods described above. For example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using the Zuker
30 calculation method even if the first sequence does not have 80 percent homology to the second sequence as calculated by any of the other calculation methods. As another example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using both the Zuker calculation method and the Pearson and Lipman calculation method even if the first
35 sequence does not have 80 percent homology to the second sequence as calculated by the Smith and Waterman calculation method, the Needleman and Wunsch calculation method,

5 the Jaeger calculation methods, or any of the other calculation methods. As yet another example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using each of calculation methods (although, in practice, the different calculation methods will often result in different calculated homology percentages).

10 Stringency of hybridization is controlled by both temperature and salt concentration of either or both of the hybridization and washing steps. Typically, the stringency of hybridization to achieve selective hybridization involves hybridization in high ionic strength solution (6X SSC or 6X SSPE) at a temperature that is about 12-25°C below the T_m (the melting temperature at which half of the molecules dissociate from their hybridization
15 partners) followed by washing at a combination of temperature and salt concentration chosen so that the washing temperature is about 5°C to 20°C below the T_m . The temperature and salt conditions are readily determined empirically in preliminary experiments in which samples of reference DNA immobilized on filters are hybridized to a labeled nucleic acid of interest and then washed under conditions of different stringencies.
20 Hybridization temperatures are typically higher for DNA-RNA and RNA-RNA hybridizations. The washing temperatures can be used as described above to achieve selective stringency, as is known in the art. (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989; Kunkel et al. Methods Enzymol. 1987:154:367, 1987). A preferable stringent
25 hybridization condition for a DNA:DNA hybridization can be at about 68°C (in aqueous solution) in 6X SSC or 6X SSPE followed by washing at 68°C. Stringency of hybridization and washing, if desired, can be reduced accordingly as the degree of complementarity desired is decreased, and further, depending upon the G-C or A-T richness of any area wherein variability is searched for. Likewise, stringency of hybridization and washing, if
30 desired, can be increased accordingly as homology desired is increased, and further, depending upon the G-C or A-T richness of any area wherein high homology is desired, all as known in the art.

In vivo administration to a human subject or an animal model can be by any of many standard means for administering viruses, depending upon the target organ, tissue or cell.
35 Virus particles can be administered orally, parenterally (e.g., intravenously), by intramuscular injection, intrarectally, by direct tissue or organ injection, by intraperitoneal

5 injection, topically, transdermally, via aerosol delivery, via the mucosa or the like. Viral nucleic acids (non-encapsidated) can also be administered, e.g., as a complex with cationic liposomes, or encapsulated in anionic liposomes. The present compositions can include various amounts of the selected viral particle or non-encapsidated viral nucleic acid in combination with a pharmaceutically acceptable carrier and, in addition, if desired, may
10 include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, etc. Parental administration, if used, is generally characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Dosages will depend upon the mode of administration, the disease or condition to be treated, and the
15 individual subject's condition, but will be that dosage typical for and used in administration of other AAV vectors, such as AAV2 vectors. Often a single dose can be sufficient; however, the dose can be repeated if desirable.

Administration of a recombinant AAV virion to the cell can be accomplished by any means, including simply contacting the particle, optionally contained in a desired liquid
20 such as tissue culture medium, or a buffered saline solution, with the cells. The virion can be allowed to remain in contact with the cells for any desired length of time, and typically the virion is administered and allowed to remain indefinitely. For such *in vitro* methods, the virion can be administered to the cell by standard viral transduction methods, as known in the art and as exemplified herein. Titers of virus to administer can vary, particularly
25 depending upon the cell type, but will be typical of that used for AAV transduction in general which is well known in the art. Additionally the titers used to transduce the particular cells in the present examples can be utilized.

The cells that can be transduced by the present recombinant AAV virions can include any desired cell, such as the following cells and cells derived from the following
30 tissues, human as well as other mammalian tissues, such as primate, horse, sheep, goat, pig, dog, rat, and mouse and avian species: Adipocytes, Adenocyte, Adrenal cortex, Amnion, Aorta, Ascites, Astrocyte, Bladder, Bone, Bone marrow, Brain, Breast, Bronchus, Cardiac muscle, Cecum, Cervix, Chorion, Cochlear, Colon, Conjunctiva, Connective tissue, Cornea, Dermis, Duodenum, Embryonic stem cells, Endometrium, Endothelium, Endothelial cells,
35 Epithelial tissue, Epithelial cells, Epidermis, Esophagus, Eye, Fascia, Fibroblasts, Foreskin, Gastric, Glial cells, Glioblast, Gonad, Hepatic cells, Histocyte, Hair cells in the inner ear,

5 Ileum, Intestine, small Intestine, Jejunum, Keratinocytes, Kidney, Larynx, Leukocytes,
Lipocyte, Liver, Lung, Lymph node, Lymphoblast, Lymphocytes, Macrophages, Mammary
alveolar nodule, Mammary gland, Mastocyte, Maxilla, Melanocytes, Mesenchymal,
Monocytes, Mouth, Myelin, Myoblasts Nervous tissue, Neuroblast, Neurons, Neuroglia,
Osteoblasts, Osteogenic cells, Ovary, Palate, Pancreas, Papilloma, Peritoneum, Pituicytes,
10 Pharynx, Placenta, Plasma cells, Pleura, Prostate, Rectum, Salivary gland, Skeletal muscle,
Skin, Smooth muscle, Somatic, Spleen, Squamous, Stem cells, Stomach, Submandibular
gland, Submaxillary gland, Synoviocytes, Testis, Thymus, Thyroid, Trabeculae, Trachea,
Turbinate, Umbilical cord, Ureter, Uterus, and vestibular hair cells.

Stringency of hybridization is controlled by both temperature and salt concentration
15 of either or both of the hybridization and washing steps. Typically, the stringency of
hybridization to achieve selective hybridization involves hybridization in high ionic strength
solution (6X SSC or 6X SSPE) at a temperature that is about 12-25°C below the T_m (the
melting temperature at which half of the molecules dissociate from their hybridization
partners) followed by washing at a combination of temperature and salt concentration
20 chosen so that the washing temperature is about 5°C to 20°C below the T_m. The
temperature and salt conditions are readily determined empirically in preliminary
experiments in which samples of reference DNA immobilized on filters are hybridized to a
labeled nucleic acid of interest and then washed under conditions of different stringencies.
Hybridization temperatures are typically higher for DNA-RNA and RNA-RNA
25 hybridizations. The washing temperatures can be used as described above to achieve
selective stringency, as is known in the art. (Sambrook et al., Molecular Cloning: A
Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New
York, 1989; Kunkel et al. Methods Enzymol. 1987:154:367, 1987). A preferable stringent
hybridization condition for a DNA:DNA hybridization can be at about 68°C (in aqueous
30 solution) in 6X SSC or 6X SSPE followed by washing at 68°C. Stringency of hybridization
and washing, if desired, can be reduced accordingly as the degree of complementarity
desired is decreased, and further, depending upon the G-C or A-T richness of any area
wherein variability is searched for. Likewise, stringency of hybridization and washing, if
desired, can be increased accordingly as homology desired is increased, and further,
35 depending upon the G-C or A-T richness of any area wherein high homology is desired, all
as known in the art.

5 By the "suitability of an AAV vector for administration to a subject" is meant a determination of whether the AAV vector will elicit a neutralizing immune response upon administration to a particular subject. A vector that does not elicit a significant immune response is a potentially suitable vector, whereas a vector that elicits a significant, neutralizing immune response (e.g. at least 90%) is thus likely to be unsuitable for use in
10 that subject. Significance of any detectable immune response is a standard parameter understood by the skilled artisan in the field. For example, one can incubate the subject's serum with the virus, then determine whether that virus retains its ability to transduce cells in culture. If such virus cannot transduce cells in culture, the vector likely has elicited a significant immune response.

15 Alternatively, or additionally, one skilled in the art could determine whether or not AAV administration would be suitable for a particular cell type of a subject. For example, the artisan could culture muscle cells *in vitro* and transduce the cells with AAV in the presence or absence of the subject's serum. If there is a reduction in transduction efficiency, this could indicate the presence of a neutralizing antibody or other factors that may inhibit
20 transduction. Normally, greater than 90% inhibition would have to be observed in order to rule out the use of AAV-5 as a vector. However, this limitation could be overcome by treating the subject with an immunosuppressant that could block the factors inhibiting transduction.

EXAMPLES

25 The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices and/or methods claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect
30 to numbers (e.g., amounts, temperature, etc.), but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in °C or is at ambient temperature, and pressure is at or near atmospheric.

Example 1

Previous research had demonstrated that Caco-2 and MDCK cells are model cell
35 lines for the study of macromolecular transport via transcytosis. Furthermore these cell lines

5 have been used to demonstrate transcytosis of both viruses and proteins. Therefore, to test if AAV can spread through tissue by transcytosis, 2×10^8 DNA resistant particles of recombinant AAV2 (rAAV2) AAV4, AAV5, AAV6, BAAV suspended in 50ul of medium were placed in the upper (apical) side of the transwell polycarbonate filter over a monolayer of cells each of the following cells Caco-2, MDCKI, MDCKII, Human primary airways
10 epithelia cells (Airway), Human primary immortalized epithelial endometrial, Bovine brain primary endothelia cells (BBB), or HeLa. All cultures had TERs indicating the formation of tight junctions and polarized phenotype. After 3 hours of incubation the medium in the basal side of the transwell was collected and tested for the presence of transcytosed rAAV DNA. Viral DNA was extracted from 200ul of basal medium and quantified by qPCR.

15 In these cell lines, transcytosis was observed with several AAV serotypes and appeared to be both serotype and tissue-specific (Fig. 1). Three hours after the addition of AAV to the apical surface of the cells, over 800,000 particles of AAV5 were present in the media on the basal lateral side of the trans-well insert of CaCo-2 cells, but not the MDCK, airway epithelia, endometrial, or BBB cells (Fig. 1). Similarly BAAV particles were detected in the media on the
20 basal lateral side of the MDCK, airways epithelia, endometrial, and BBB cells but not the Caco-2 cells. Interestingly, AAV4 was detected in the basal lateral media of all cell types. No virus was detected in the basal lateral media when AAV2 was added to the apical surface in either cell type. AAV6 did not transcytose in any of cell types tested, and was not tested on airway epithelia or BBB. HeLa cells do not form barrier epithelia and were used as a control.

25 Example 2

Previous work has demonstrated that transcytosis is a temperature dependent process than can be inhibited at 4°C. Transcytosis can also be inhibited by the addition of agents that selectively fix the plasma membrane. Recently the addition of tannic acid, a mild fixative agent, to the basal lateral surface blocked the transcytosis of GPI-anchored proteins to the
30 apical surface (Polishchuk R, *Nat Cell Biol.* 2004. 6(4):297-307). Therefore the ability of this agent to block the transcytosis of AAV was tested. Treatment of the basal lateral surface of either Caco-2 or MDCK cells prior to virus addition to the apical surface blocked the accumulation of AAV5 or BAAV in the basal lateral media. Furthermore, quantification of the intracellular virus demonstrated inhibition of exocytosis by tannic acid treatment
35 dramatically increase the amount of AAV DNA in the cell suggesting the viral particles

5 detected in the basal lateral media are the result of an intracellular transport process and not a paracellular route.

Treatment of the basal lateral surface of Human primary airways epithelial cell (HAE) with tannic acid blocked the transcytosis of BAAV or AAV4 vector containing a GFP expression cassette from the apical surface to the basal lateral (Fig. 2). Furthermore
10 transduction dramatically increased when assayed at 24 hrs post inoculation. In contrast no change was observed in AAV2 transduction, which did not demonstrate any transcytosis activity and has limited binding activity on HAE.

Example 3

To confirm the DNA detected in the basal lateral media was indeed extracted from
15 intact virus, the material was tested for DNase resistance after treatment with heat, ionic detergent or protease. The addition of DNase alone or in combination with the ionic detergent deoxycholine had no effect on the viral DNA present in the media suggesting it was not free DNA or complexed in lipid vesicles. However, heating to 95°C prior to treatment with DNAase completely degraded the viral DNA present in the media. This
20 profile is identical to that of the input AAV particles and suggests the viral DNA is still encapsulated. Titration of the DNase resistant virus in the basal lateral media on Cos cells gave a similar particle to infectivity ratio to the input AAV particles.

While it would appear the AAV DNA detected in the basal lateral media is contained in intact particles, its presence on the basal lateral surface could be the result of
25 lyses of the cells or disruption of the monolayer. Therefore the TER was carefully monitored throughout the course of these experiments and was not observed to decrease. To further confirm the integrity of the cell monolayer, mixing experiments were studied in which two viruses with different gene cassettes were added to the apical surface at the same time and three hours post addition the amount of each virus in the basal lateral media was quantified
30 using QPCR specific for each cassette. Both BAAV and AAV5 were able to pass from the apical to the basal lateral surface of MDCK or Caco cells respectively but the AAV2 did not. Therefore the presence of viral particles in the basal lateral media does not appear to be the result of a disruption in the cell monolayer.

Taken together this data suggest that dependoviruses particles are capable of passing
35 through barrier epithelia via transcytosis and the process is both serotype and cell type specific.

5

Example 4

To further characterize the transcytosis activity observed with AAV5 and BAAV, transcytosis was quantified as both a time and concentration dependent event. After the addition of particles to the apical surface, samples were removed from the basal lateral media at different time points and the amount of virus was quantified by QPCR of the extracted DNA. Viral genomes could be detected as soon as 30 minutes after addition and steadily increased with time. By 24 hrs, over 1/3 of the input recombinant AAV5, BAAVvirus added to Caco or MDCK cells respectively had been transported to the basal lateral surface. In contrast, none of the input AAV2 or adenovirus was detected on the basal lateral side after 24 hrs.

15 If transcytosis is an activity used by AAV to spread through tissue, this finding would help explain the lack of transduction of barrier epithelia reported with some isolates of AAV. Primary human bronchial airway epithelia (HAE) are known to transport albumin from the apical to the basal lateral surface by receptor-mediated transcytosis in vivo. While the interaction of BAAV with primary HAE has not been investigated, AAV4, 5 are reported to bind to HAE, however, for AAV4, this interaction does not result in transduction. Because of the interaction of AAV4 with O-link sialic acid, it was proposed, and has been demonstrated, that mucins, which contained large amounts of O-linked sialic acid and are expressed on the apical surface of HAE, can block AAV4 transduction. Alternatively the lack of transduction could be the result of transcytosis of the virus through the tissue.

25 To test this hypothesis, AAV2, 4, 5, BAAV were added to the apical surface of confluent monolayer cultures of primary human bronchial airway and transcytosis to the basal lateral surface was measured by QPCR after 3 hrs. All cultures had high TERs and expressed ciliated structures on their apical surface. Highly differentiated HAE cultures in contrast to immature cultures are resistant to transduction by adenoviral vectors due to a lack of integrin expression that is necessary for adenovirus entry.

30 Of the 4 AAVs tested for transcytosis, AAV4 and BAAV were detected in the basal lateral media. No transport of AAV2 or AAV5 was detected. As a control, adenovirus also was tested for transcytosis activity in the HAE cultures, but no transport was detected.

5

Example 5

Epithelial cells that line the genitourinary tract form an important epithelial barrier layer and can transport proteins by transcytosis. AAV2, 4, 5 or BAAV were therefore tested to determine for the ability to penetrate this barrier epithelial layer by transcytosis. A well-characterized model of endometrial cells has been reported by Kyo et al. Following addition
10 of the 4 AAVs to the apical surface, BAAV and AAV4 could be detected in the basal lateral media when assayed at 3hrs post inoculation (Fig. 1).

Example 6

Most AAVs were identified originally as contaminants of laboratory stocks of adenovirus, thus our understanding of their natural biology, cell tropism, and knowledge the
15 cellular components required for virus entry is limited. For AAV5, in addition to N-linked sialic acid, the platelet derived growth factor (PDGF) receptors were identified as protein receptors for AAV5 (Di Pasquale et al., Nat Med. 2003 Oct;9(10):1306-12). This interaction was confirmed by modulation of PDGFR expression by transfection of expression plasmids, inhibitor treatment, or competition experiments with the extracellular domain of PDGFR α .
20 Likewise AAV5 transduction could be blocked with sialolactosamine conjugates kaludov et al 2001.

Previous research had demonstrated that transcytosis is actin dependent and occurs by a caviolin mediated pathway. Furthermore transcytosis can be blocked by treatment with tannic acid. Therefore to better characterize the transcytosis pathway utilized by AAV5 in
25 Caco cells the cells were treated with a panel of agents known to block either transcytosis in other systems or AAV5 mediated transduction. It was noted that AAV5 transcytosis could be inhibited by filipin and nocozodol as well as treatment with tannic acid.

Caco cells, which actively transcytosis AAV5, are not reported to express PDGFR and are not transduced by AAV5. In agreement, competition experiments with sPDGFR α
30 had little effect on AAV5 transcytosis. Furthermore, competition experiments with 200 ug/ml sialolactosamine or 200 ug/ml heparin did not inhibited AAV5 transcytosis.

Both BSA and transferrin are reported to transcytosis through Caco cells via distinct receptor mediated pathways. However competition with either agent did not inhibit AAV5 transcytosis suggesting the AAV5 could use a distinct pathway.

5 In addition to confirming the intracellular nature of AAV5 transcytosis in Caco cells, the above experiments suggest that AAV5 transcytosis is occurring by a pathway independent of the one described for transduction. To confirm this Caco cells were stably transfected with PDGFRa and assayed for both transcytosis and transduction activity. Caco cells were not permissive for AAV5 transduction, however transduction dramatically increase
10 following stable expression of PDGFRa. In contrast only a minor increase in transcytosis activity was detected in the Caco/PDGFRa cells.

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention
15 pertains.

It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is
20 intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

CLAIMS

What is claimed is:

1. A method of delivering a heterologous nucleic acid across an epithelial barrier comprising delivering to the epithelial barrier an AAV vector comprising the heterologous nucleic acid.
2. The method of claim 1, wherein the epithelial cells are in the gut, lung, genitourinary tract, kidney, blood vessels or brain.
3. The method of claim 1, wherein the epithelial cells can be selected from a group consisting of bronchial, alveolar, tracheal or upper airway epithelial cells; absorptive enterocytes; endometrial or urinary epithelial cells; renal collecting duct or proximal tubule epithelial cells; cerebral microvascular endothelial cells; or Choroidal Plexus epithelial cells .
4. A method of transcytosing epithelial cells of a human subject comprising administering to the subject an AAV vector comprising a heterologous nucleic acid.
5. The method of claim 4, wherein the epithelial cells are selected from a group consisting of bronchial, alveolar, tracheal or upper airway epithelial cells; absorptive enterocytes; endometrial or urinary epithelial cells; renal collecting duct or proximal tubule epithelial cells; cerebral microvascular endothelial cells; or Choroidal Plexus epithelial cells.
6. A method of delivering a heterologous nucleic acid across human airway epithelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.
7. A method of delivering a heterologous nucleic acid across human cerebral microvascular endothelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.
8. A method of delivering a heterologous nucleic acid across human endometrial epithelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.
9. A method of delivering a heterologous nucleic acid across human kidney epithelial cells, comprising delivering to the cells a BAAV vector comprising the nucleic acid.
10. A method of delivering a heterologous nucleic acid across human enterocytes, comprising delivering to the cells a AAV5 vector comprising the nucleic acid.

11. A method of delivering a heterologous nucleic acid across human airway epithelial cells, comprising delivering to the cells a AAV4 vector comprising the nucleic acid.
12. A method of delivering a heterologous nucleic acid across human cerebral microvascular endothelial cells, comprising delivering to the cells a AAV4 vector comprising the nucleic acid.
13. A method of delivering a heterologous nucleic acid across human endometrial epithelial cells, comprising delivering to the cells a AAV4 vector comprising the nucleic acid.
14. A method of delivering a heterologous nucleic acid across human kidney epithelial cells, comprising delivering to the cells a AAV4 vector comprising the nucleic acid.
15. A method of delivering a heterologous nucleic acid across human enterocytes comprising delivering to the cells a AAV4 vector comprising the nucleic acid.
16. A method of delivering a heterologous nucleic acid across human cerebral microvascular endothelial cells, comprising delivering to the cells a AAV7 vector comprising the nucleic acid.
17. A method of delivering a heterologous nucleic acid across an epithelial barrier of the lung, comprising delivering to the lung a BAAV vector comprising the nucleic acid.
18. The method of claim 17, wherein the epithelial barrier comprises human bronchial, alveolar, tracheal or upper airway epithelial cells.
19. A method of delivering a heterologous nucleic acid across an epithelial barrier in the brain, comprising delivering to the brain a BAAV vector comprising the nucleic acid.
20. The method of claim 19, wherein the epithelial barrier comprises human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.
21. A method of delivering a heterologous nucleic acid across the epithelial barrier of blood vessels into the muscle, comprising delivering to the blood stream a BAAV vector comprising the nucleic acid.
22. The method of claim 21, wherein the epithelial barrier comprises human vascular endothelial cells of the blood brain barrier.

23. A method of delivering a heterologous nucleic acid across an epithelial barrier in the genitourinary tract, comprising delivering to the genitourinary tract a BAAV vector comprising the nucleic acid genitourinary tract.
24. The method of claim 23, wherein the epithelial barrier comprises human endometrial or urinary epithelial cells.
25. A method of delivering a heterologous nucleic acid across an epithelial barrier in the kidney, comprising delivering to the genitourinary tract a BAAV vector comprising the nucleic acid genitourinary tract.
26. The method of claim 25, wherein the epithelial barrier comprises human renal collecting ducts or proximal tubules.
27. A method of transcytosing lung epithelial cells of a subject comprising contacting the lung epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid.
28. The method of claim 27, wherein the epithelial cells are human bronchial, tracheal, or upper airway epithelial cells.
29. A method of transcytosing CNS epithelial cells of a subject comprising contacting the CNS epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid.
30. The method of claim 29, wherein the epithelial cells are human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.
31. A method of transcytosing vascular epithelial cells of a subject comprising contacting the vascular epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid.
32. The method of claim 31, wherein the epithelial cells are human vascular endothelial cells of the blood brain barrier.
33. A method of transcytosing genitourinary tract epithelial cells of a subject comprising contacting the genitourinary tract epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid.
34. The method of claim 33, wherein the epithelial cells are human endometrial or urinary tract epithelial cells.

35. A method of transcytosing genitourinary tract epithelial cells of a subject comprising contacting the kidney epithelial cells of the subject with a BAAV vector comprising a heterologous nucleic acid.
36. The method of claim 35, wherein the epithelial cells are human renal collecting ducts or proximal tubules
37. A method of delivering a heterologous nucleic acid across an epithelial barrier in the gut, comprising delivering to the gut an AAV5 vector comprising the nucleic acid.
38. The method of claim 37, wherein the epithelial barrier comprises human absorptive enterocytes.
39. A method of transcytosing gut epithelial cells of a subject comprising contacting the gut epithelial cells of the subject with an AAV5 vector comprising a heterologous nucleic acid.
40. The method of claim 39, wherein the epithelial cells are human absorptive enterocytes.
41. A method of delivering a heterologous nucleic acid across an epithelial barrier in the gut, comprising delivering to the gut an AAV4 vector comprising the nucleic acid.
42. The method of claim 41, wherein the epithelial barrier comprises human absorptive enterocytes.
43. A method of delivering a heterologous nucleic acid across an epithelial barrier in the lung, comprising delivering to the lung an AAV4 vector comprising the nucleic acid.
44. The method of claim 43, wherein the epithelial barrier comprises human bronchial, tracheal, or upper airway epithelial cells.
45. A method of delivering a heterologous nucleic acid across an epithelial barrier in the CNS, comprising delivering to the CNS an AAV4 vector comprising the nucleic acid.
46. The method of claim 45, wherein the epithelial barrier comprises human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.
47. A method of delivering a heterologous nucleic acid across the epithelial barrier of blood vessels into the muscle, comprising delivering to the blood stream an AAV4 vector comprising the nucleic acid.

48. The method of claim 47, wherein the epithelial barrier comprises human vascular endothelial cells of the blood brain barrier.
49. A method of delivering a heterologous nucleic acid across an epithelial barrier in the genitourinary tract, comprising delivering to the genitourinary tract an AAV4 vector comprising the nucleic acid.
50. The method of claim 49, wherein the epithelial barrier comprises human endometrial or urinary epithelial cells.
51. A method of delivering a heterologous nucleic acid across an epithelial barrier in the kidneys, comprising delivering to the kidneys an AAV4 vector comprising the nucleic acid.
52. The method of claim 51, wherein the epithelial barrier comprises human renal collecting ducts or proximal tubules.
53. A method of transcytosing lung epithelial cells of a subject comprising contacting the lung epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid.
54. The method of 53, wherein the epithelial cells are human bronchial, tracheal, or upper airway epithelial cells.
55. A method of transcytosing CNS epithelial cells of a subject comprising contacting the CNS epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid.
56. The method of claim 55, wherein the epithelial cells are human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.
57. A method of transcytosing vascular epithelial cells of a subject comprising contacting the vascular epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid.
58. The method of claim 57, wherein the epithelial cells are vascular endothelial cells of the blood brain barrier.
59. A method of transcytosing genitourinary tract epithelial cells of a subject comprising contacting the genitourinary epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid.

60. The method of claim 59, wherein the epithelial cells are human endometrial or urinary epithelial cells.
61. A method of transcytosing kidney epithelial cells of a subject comprising contacting the kidney epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid.
62. The method of claim 61, wherein the epithelial cells are human renal collecting ducts or proximal tubules
63. A method of transcytosing gut epithelial cells of a subject comprising contacting the CNS epithelial cells of the subject with an AAV4 vector comprising a heterologous nucleic acid.
64. The method of claim 63, wherein the epithelial cells are human absorptive enterocytes.
65. A method of delivering a heterologous nucleic acid across an epithelial barrier in the brain, comprising delivering to the brain a AAV7 vector comprising the nucleic acid.
66. The method of claim 65, wherein the epithelial barrier comprises human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.
67. A method of transcytosing CNS epithelial cells of a subject comprising contacting the CNS epithelial cells of the subject with a AAV7 vector comprising a heterologous nucleic acid.
68. The method of claim 67, wherein the epithelial cells are human cerebral microvascular endothelial cells or Choroidal Plexus epithelial cells of the blood brain barrier.




FIG. 1




FIG. 2




FIG. 3

SEQUENCE LISTING

<110> The Government of the United States of America, as represented by The Secretary, Department of Health and Human Services, National Institutes of Health
Chiorini, John A.
DePasquale, Giovanni

<120> TRANSCYTOSIS OF ADENO-ASSOCIATED VIRUSES

<130> 14014/0427P1

<140> TBA

<141> 2005-09-08

<150> 60/607,854

<151> 2004-09-08

<160> 72

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 4768

<212> DNA

<213> Artificial sequence

<220>

<223> Description of Artificial sequence; note = synthetic construct

<221> misc_feature

<222> (0)...(0)

<223> n=a,t,c, or g

<221> variation

<222> (0)...(0)

<223> Xaa=any amino acid

<400> 1

ttggccactc	cctctatgcg	cgctcgctca	ctcactcggc	cctggagacc	aaaggtctcc	60
agactgccgg	cctctggccg	gcagggccga	gtgagtgagc	gagcgcgcac	agagggagtg	120
gccaactcca	tcattctagg	ttgcccactg	acgtcaatgt	gacgtcctag	ggttagggag	180
gtccctgtat	tagcagtcac	gtgagtgtcg	tatttcgcgg	agcgtagcgg	agcgcatacc	240
aagctgccac	gtcacagcca	cgtggtccgt	ttgcgacagt	ttgcgacacc	atgtggtcag	300
gagggatat	aaccgcgagt	gagccagcga	ggagctccat	tttgcccgcg	aattttgaac	360
gagcagcagc	catgccgggg	ttctacgaga	tcgtgctgaa	ggtgcccagc	gacctggacg	420
agcacctgcc	cggcatttct	gactcttttg	tgagctgggt	ggccgagaag	gaatgggagc	480
tgccgcccga	ttctgacatg	gacttgaatc	tgattgagca	ggcacccttg	accgtggccg	540
aaaagctgca	acgcgagttc	ctggctcgagt	ggcgccgcgt	gagtaaggcc	ccggaggccc	600
tcttctttgt	ccagttcgag	aagggggaca	gctacttcca	cctgcacatc	ctggtggaga	660
ccgtgggctg	caaattccatg	gtgggtgggc	gctacgtgag	ccagattaaa	gagaagctgg	720
tgacccgcac	ctaccgcggg	gtcgagccgc	agcttccgaa	ctggttcgcg	gtgaccaaga	780
cgcgtaattg	cgccggaggc	gggaacaagg	tggtggacga	ctgctacatc	cccaactacc	840
tgctcccca	gacccagccc	gagctccagt	gggcgtggac	taacatggac	cagtatatata	900
gcgcctgttt	gaatctcgcg	gagcgtaa	ggctgggtgg	gcagcatctg	acgcacgtgt	960
cgagcagcga	ggagcagaac	aaggaaaacc	agaaccccaa	ttctgacgcg	ccggtcatca	1020
ggtcaaaaac	ctccgccagg	tacatggagc	tggtcgggtg	gctgggtggac	cgcgggatca	1080
cgtcagaaaa	gcaatggatc	caggaggacc	aggcgtccta	catctccttc	aacgccgcct	1140
ccaactcgcg	gtcacaatc	aaggccgcgc	tggacaatgc	ctccaaaatc	atgagcctga	1200
caaagacggc	tccggactac	ctggtggg	agaacccgcc	ggaggacatt	tccagcaacc	1260

gcattctaccg	aatcctcagag	atgaacgggt	acgatccgca	gtacgcggcc	tccgtcttcc	1320
tgggctgggc	gcaaaagaag	ttcgggaaga	ggaacacccat	ctggctcttt	gggcccggcca	1380
cgacgggtaa	aaccaacatc	gcggaagcca	tcgcccacgc	cgtgcccttc	tacggctgcg	1440
tgaactggac	caatgagaac	tttccgttca	acgattgcgt	cgacaagatg	gtgatctggt	1500
gggaggagg	caagatgacg	gccaaggctc	tagagagcgc	caaggccatc	ctgggcggaa	1560
gcaagggtgc	cgtggaccaa	aagtgcaggt	catcggccca	gatcgaccca	actcccgtga	1620
tcgtcacctc	caacaccaac	atgtgcgcgg	tcacgcagcg	aaactcgacc	accttcgagc	1680
accaacaacc	actccaggac	cggatgttca	agttcgagct	caccaagcgc	ctggagcacg	1740
acttttgcaa	ggtcaccaag	caggaagtca	aagacttttt	ccggtgggcg	tcagatcacg	1800
tgaccgaggt	gactcacgag	ttttacgtca	gaaaggggtg	agctagaaaag	aggcccggccc	1860
ccaatgacgc	agatataagt	gagcccaagc	gggctgtgcc	gtcagttgcg	cagccatcga	1920
cgtcagacgc	ggaagctccg	gtggactacg	cggacaggta	ccaaaacaaa	tgttctcgtc	1980
acgtgggtat	gaatctgatg	ctttttccct	gccggcaatg	cgagagaatg	aatcagaatg	2040
tggacatttg	cttcacgcac	ggggtcatgg	actgtgccga	gtgcttcccc	gtgtcagaat	2100
ctcaacccgt	gtctgtcgtc	agaaagcgga	cgtatcagaa	actgtgtccg	attcatcaca	2160
tcattggggag	ggcgcccagag	gtggcctgct	cggcctgcga	actggccaat	gtggacttgg	2220
atgactgtga	catggaacaa	taaatgactc	aaaccagata	tgactgacgg	ttaccttcca	2280
gattggctag	aggacaacct	ctctgaaggc	gttcgagagt	ggtgggcgct	gcaacctgga	2340
gcccctaaac	ccaaggcaaa	tcaacaacat	caggacaacg	ctcggggtct	tgtgcttccg	2400
ggttacaaat	acctcggacc	cggcaacgga	ctcgacaagg	gggaacccgt	caacgcagcg	2460
gacgcggcag	ccctcgagca	cgacaaggcc	tacgaccagc	agctcaaggc	cggtgacaac	2520
ccctacctca	agtacaacca	cgccgacgcg	gagttccagc	agcggcttca	gggcgacaca	2580
ccgtttgggg	gcaacctcgg	cagagcagtc	ttccaggcca	aaaagagggt	tcttgaacct	2640
cttggctctgg	ttgagcaagc	gggtgagacg	gctcctggaa	agaagagacc	gttgattgaa	2700
tccccccagc	agcccgaactc	ctccacgggt	atcggcaaaa	aaggcaagca	gccggctaaa	2760
aagaagctcg	ttttcgaaga	cgaaactgga	gcaggcgacg	gacccctga	gggatcaact	2820
tccggagcca	tgtctgatga	cagtgaagtg	cgtgcagcag	ctggcggagc	tgcagtcgag	2880
gsgggacaag	gtgccgatgg	agtgggtaat	gcctcgggtg	attggcattg	cgattccacc	2940
tggctctgag	gccacgtcac	gaccaccagc	accagaacct	gggtcttgcc	cacctacaac	3000
aaccacctnt	acaagcgact	cggagagagc	ctgcagtcca	acacctacaa	cggattctcc	3060
accccttggg	gatactttga	cttcaaccgc	ttccactgcc	acttctcacc	acgtgactgg	3120
cagcgactca	tcaacaacaa	ctggggcatg	cgacccaaag	ccatgcgggt	caaaatcttc	3180
aacatccagg	tcaaggaggt	cacgacgtcg	aacggcgaga	caacgggtggc	taataacctt	3240
accagcacgg	ttcagatctt	tgcggactcg	tcgtacgaac	tgccgtacgt	gatggatgcg	3300
ggtcaagagg	gcagcctgcc	tccttttccc	aacgacgtct	ttatgggtgcc	ccagtacggc	3360
tactgtggac	tggtgaccgg	caacacttcg	cagcaacaga	ctgacagaaa	tgccttctac	3420
tgccctggagt	actttccttc	gcagatgctg	cggactggca	acaactttga	aattacgtac	3480
agttttgaga	aggtgccttt	ccactcgatg	tacgcgcaca	gccagagcct	ggaccggctg	3540
atgaaccctc	tcacgcacca	gtacctgtgg	ggactgcaat	cgaccaccac	cggaaccacc	3600
ctgaatgccg	ggactgccac	caccaacttt	accaagctgc	ggcctaccaa	cttttccaac	3660
tttaaaaaga	actggctgcc	cgggccttca	atcaagcagc	agggcttctc	aaagactgcc	3720
aatcaaaaact	acaagatccc	tgccaccggg	tcagacagtc	tcacaaata	cgagacgcac	3780
agcactctgg	acggaagatg	gagtgccttg	accccgccgac	ctccaatggc	cacggctgga	3840
cctgcggaca	gcaagttcag	caacagccag	ctcatctttg	cggggcctaa	acagaacggc	3900
aacacggcca	ccgtacccgg	gactctgatc	ttcacctctg	aggaggagct	ggcagccacc	3960
aacgccaccg	atacggacat	gtggggcaac	ctacctggcg	gtgaccagag	caacagcaac	4020
ctgccgaccg	tggacagact	gacagccttg	ggagccgtgc	ctggaatggg	ctggcaaaac	4080
agagacattt	actaccaggg	tcccattttg	gccaagattc	ctcataccga	tggacacttt	4140
cacccctcac	cgctgatttg	tgggttttgg	ctgaaacacc	cgctctctca	aattttttatc	4200
aagaacaccc	cgggtacctg	gaatcctgca	acgaccttca	gctctactcc	ggtaaaactcc	4260
ttcattactc	agtacagcac	tggccagggtg	tcgggtgcaga	ttgactggga	gatccagaag	4320
gagcgggtcca	aacgctggaa	ccccgagggtc	cagttttacct	ccaactacgg	acagcaaaac	4380
tctctgttgt	gggctcccga	tgcggctggg	aaatacactg	agcctagggc	tatcgggtacc	4440
cgctacctca	cccaccacct	gtaataacct	gttaatacaat	aaaccggttt	attcgtttca	4500
gttgaacttt	ggtctccgtg	tccttcttat	cttatctcgt	ttccatggct	actgcgtaca	4560
taagcagcgg	cctgcggcgc	ttgcgcttcg	cggttttacaa	ctgccgggta	atcagtaact	4620
tctggcaaac	catgatgatg	gagttggcca	ctccctctat	gcgcgctcgc	tcactcactc	4680
ggccctggag	accaaagggtc	tccagactgc	cggcctctgg	ccggcagggc	cgagtgaagtg	4740
agcgagcgcg	catagaggga	gtggccaa				4768

<210> 2
 <211> 623
 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 2

Met	Pro	Gly	Phe	Tyr	Glu	Ile	Val	Leu	Lys	Val	Pro	Ser	Asp	Leu	Asp
1				5					10					15	
Glu	His	Leu	Pro	Gly	Ile	Ser	Asp	Ser	Phe	Val	Ser	Trp	Val	Ala	Glu
			20					25					30		
Lys	Glu	Trp	Glu	Leu	Pro	Pro	Asp	Ser	Asp	Met	Asp	Leu	Asn	Leu	Ile
		35					40					45			
Glu	Gln	Ala	Pro	Leu	Thr	Val	Ala	Glu	Lys	Leu	Gln	Arg	Glu	Phe	Leu
	50					55					60				
Val	Glu	Trp	Arg	Arg	Val	Ser	Lys	Ala	Pro	Glu	Ala	Leu	Phe	Phe	Val
65					70				75						80
Gln	Phe	Glu	Lys	Gly	Asp	Ser	Tyr	Phe	His	Leu	His	Ile	Leu	Val	Glu
				85					90					95	
Thr	Val	Gly	Val	Lys	Ser	Met	Val	Val	Gly	Arg	Tyr	Val	Ser	Gln	Ile
			100					105					110		
Lys	Glu	Lys	Leu	Val	Thr	Arg	Ile	Tyr	Arg	Gly	Val	Glu	Pro	Gln	Leu
		115					120					125			
Pro	Asn	Trp	Phe	Ala	Val	Thr	Lys	Thr	Arg	Asn	Gly	Ala	Gly	Gly	Gly
	130					135					140				
Asn	Lys	Val	Val	Asp	Asp	Cys	Tyr	Ile	Pro	Asn	Tyr	Leu	Leu	Pro	Lys
145					150				155						160
Thr	Gln	Pro	Glu	Leu	Gln	Trp	Ala	Trp	Thr	Asn	Met	Asp	Gln	Tyr	Ile
				165					170					175	
Ser	Ala	Cys	Leu	Asn	Leu	Ala	Glu	Arg	Lys	Arg	Leu	Val	Ala	Gln	His
			180					185					190		
Leu	Thr	His	Val	Ser	Gln	Thr	Gln	Glu	Gln	Asn	Lys	Glu	Asn	Gln	Asn
		195					200					205			
Pro	Asn	Ser	Asp	Ala	Pro	Val	Ile	Arg	Ser	Lys	Thr	Ser	Ala	Arg	Tyr
	210					215					220				
Met	Glu	Leu	Val	Gly	Trp	Leu	Val	Asp	Arg	Gly	Ile	Thr	Ser	Glu	Lys
225					230					235					240
Gln	Trp	Ile	Gln	Glu	Asp	Gln	Ala	Ser	Tyr	Ile	Ser	Phe	Asn	Ala	Ala
				245					250					255	
Ser	Asn	Ser	Arg	Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Ser	Lys
			260					265					270		
Ile	Met	Ser	Leu	Thr	Lys	Thr	Ala	Pro	Asp	Tyr	Leu	Val	Gly	Gln	Asn
		275					280					285			
Pro	Pro	Glu	Asp	Ile	Ser	Ser	Asn	Arg	Ile	Tyr	Arg	Ile	Leu	Glu	Met
	290					295					300				
Asn	Gly	Tyr	Asp	Pro	Gln	Tyr	Ala	Ala	Ser	Val	Phe	Leu	Gly	Trp	Ala
305					310					315					320
Gln	Lys	Lys	Phe	Gly	Lys	Arg	Asn	Thr	Ile	Trp	Leu	Phe	Gly	Pro	Ala
				325					330					335	
Thr	Thr	Gly	Lys	Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Ala	Val	Pro
			340					345					350		
Phe	Tyr	Gly	Cys	Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp
		355					360					365			
Cys	Val	Asp	Lys	Met	Val	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thr	Ala
	370					375					380				
Lys	Val	Val	Glu	Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg
385					390					395					400
Val	Asp	Gln	Lys	Cys	Lys	Ser	Ser	Ala	Gln	Ile	Asp	Pro	Thr	Pro	Val
				405					410					415	
Ile	Val	Thr	Ser	Asn	Thr	Asn	Met	Cys	Ala	Val	Ile	Asp	Gly	Asn	Ser
			420					425					430		
Thr	Thr	Phe	Glu	His	Gln	Gln	Pro	Leu	Gln	Asp	Arg	Met	Phe	Lys	Phe
		435					440								

Glu Leu Thr Lys Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln
 450 455 460
 Glu Val Lys Asp Phe Phe Arg Trp Ala Ser Asp His Val Thr Glu Val
 465 470 475 480
 Thr His Glu Phe Tyr Val Arg Lys Gly Gly Ala Arg Lys Arg Pro Ala
 485 490 495
 Pro Asn Asp Ala Asp Ile Ser Glu Pro Lys Arg Ala Cys Pro Ser Val
 500 505 510
 Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Pro Val Asp Tyr Ala Asp
 515 520 525
 Arg Tyr Gln Asn Lys Cys Ser Arg His Val Gly Met Asn Leu Met Leu
 530 535 540
 Phe Pro Cys Arg Gln Cys Glu Arg Met Asn Gln Asn Val Asp Ile Cys
 545 550 555 560
 Phe Thr His Gly Val Met Asp Cys Ala Glu Cys Phe Pro Val Ser Glu
 565 570 575
 Ser Gln Pro Val Ser Val Val Arg Lys Arg Thr Tyr Gln Lys Leu Cys
 580 585 590
 Pro Ile His His Ile Met Gly Arg Ala Pro Glu Val Ala Cys Ser Ala
 595 600 605
 Cys Glu Leu Ala Asn Val Asp Leu Asp Asp Cys Asp Met Glu Gln
 610 615 620

<210> 3

<211> 2495

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 3

Ala Thr Gly Cys Cys Gly Gly Gly Gly Thr Thr Cys Thr Ala Cys Gly
 1 5 10 15
 Ala Gly Ala Thr Cys Gly Thr Gly Cys Thr Gly Ala Ala Gly Gly Thr
 20 25 30
 Gly Cys Cys Cys Ala Gly Cys Gly Ala Cys Cys Thr Gly Gly Ala Cys
 35 40 45
 Met Pro Gly Phe Tyr Glu Ile Val Leu Lys Val Pro Ser Asp Leu Asp
 50 55 60
 Gly Ala Gly Cys Ala Cys Cys Thr Gly Cys Cys Cys Gly Gly Cys Ala
 65 70 75 80
 Thr Thr Thr Cys Thr Gly Ala Cys Thr Cys Thr Thr Thr Thr Gly Thr
 85 90 95
 Gly Ala Gly Cys Thr Gly Gly Gly Thr Gly Gly Cys Cys Gly Ala Gly
 100 105 110
 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu
 115 120 125
 Ala Ala Gly Gly Ala Ala Thr Gly Gly Gly Ala Gly Cys Thr Gly Cys
 130 135 140
 Cys Gly Cys Cys Gly Gly Ala Thr Thr Cys Thr Gly Ala Cys Ala Thr
 145 150 155 160
 Gly Gly Ala Cys Thr Thr Gly Ala Ala Thr Cys Thr Gly Ala Thr Thr
 165 170 175
 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
 180 185 190
 Gly Ala Gly Cys Ala Gly Gly Cys Ala Cys Cys Cys Cys Thr Gly Ala
 195 200 205
 Cys Cys Gly Thr Gly Gly Cys Cys Gly Ala Ala Ala Ala Gly Cys Thr
 210 215 220
 Gly Cys Ala Ala Cys Gly Cys Gly Ala Gly Thr Thr Cys Cys Thr Gly
 225 230 235 240

Glu	Gln	Ala	Pro	Leu	Thr	Val	Ala	Glu	Lys	Leu	Gln	Arg	Glu	Phe	Leu
				245					250					255	
Gly	Thr	Cys	Gly	Ala	Gly	Thr	Gly	Gly	Cys	Gly	Cys	Cys	Gly	Cys	Gly
			260					265					270		
Thr	Gly	Ala	Gly	Thr	Ala	Ala	Gly	Gly	Cys	Cys	Cys	Cys	Gly	Gly	Ala
		275					280					285			
Gly	Gly	Cys	Cys	Cys	Thr	Cys	Thr	Thr	Cys	Thr	Thr	Thr	Gly	Thr	Cys
	290					295					300				
Val	Glu	Trp	Arg	Arg	Val	Ser	Lys	Ala	Pro	Glu	Ala	Leu	Phe	Phe	Val
305					310					315					320
Cys	Ala	Gly	Thr	Thr	Cys	Gly	Ala	Gly	Ala	Ala	Gly	Gly	Gly	Gly	Gly
				325				330						335	
Ala	Cys	Ala	Gly	Cys	Thr	Ala	Cys	Thr	Thr	Cys	Cys	Ala	Cys	Cys	Thr
			340					345					350		
Gly	Cys	Ala	Cys	Ala	Thr	Cys	Cys	Thr	Gly	Gly	Thr	Gly	Gly	Ala	Gly
		355					360					365			
Gln	Phe	Glu	Lys	Gly	Asp	Ser	Tyr	Phe	His	Leu	His	Ile	Leu	Val	Glu
	370					375					380				
Ala	Cys	Cys	Gly	Thr	Gly	Gly	Gly	Cys	Gly	Thr	Cys	Ala	Ala	Ala	Thr
385					390					395					400
Cys	Cys	Ala	Thr	Gly	Gly	Thr	Gly	Gly	Thr	Gly	Gly	Gly	Cys	Cys	Gly
				405				410					415		
Cys	Thr	Ala	Cys	Gly	Thr	Gly	Ala	Gly	Cys	Cys	Ala	Gly	Ala	Thr	Thr
			420					425				430			
Thr	Val	Gly	Val	Lys	Ser	Met	Val	Val	Gly	Arg	Tyr	Val	Ser	Gln	Ile
		435					440					445			
Ala	Ala	Ala	Gly	Ala	Gly	Ala	Ala	Gly	Cys	Thr	Gly	Gly	Thr	Gly	Ala
	450					455					460				
Cys	Cys	Cys	Gly	Cys	Ala	Thr	Cys	Thr	Ala	Cys	Cys	Gly	Cys	Gly	Gly
465					470					475					480
Gly	Gly	Thr	Cys	Gly	Ala	Gly	Cys	Cys	Gly	Cys	Ala	Gly	Cys	Thr	Thr
				485				490						495	
Lys	Glu	Lys	Leu	Val	Thr	Arg	Ile	Tyr	Arg	Gly	Val	Glu	Pro	Gln	Leu
			500					505					510		
Cys	Cys	Gly	Ala	Ala	Cys	Thr	Gly	Gly	Thr	Thr	Cys	Gly	Cys	Gly	Gly
		515					520					525			
Thr	Gly	Ala	Cys	Cys	Ala	Ala	Gly	Ala	Cys	Gly	Cys	Gly	Thr	Ala	Ala
	530					535				540					
Thr	Gly	Gly	Cys	Gly	Cys	Gly	Gly	Ala	Gly	Gly	Cys	Gly	Gly	Gly	Gly
545					550				555						560
Pro	Asn	Trp	Phe	Ala	Val	Thr	Lys	Thr	Arg	Asn	Gly	Ala	Gly	Gly	Gly
				565					570					575	
Ala	Ala	Cys	Ala	Ala	Gly	Gly	Thr	Gly	Gly	Thr	Gly	Gly	Ala	Cys	Gly
			580					585					590		
Ala	Cys	Thr	Gly	Cys	Thr	Ala	Cys	Ala	Thr	Cys	Cys	Cys	Cys	Ala	Ala
		595					600					605			
Cys	Thr	Ala	Cys	Cys	Thr	Gly	Cys	Thr	Cys	Cys	Cys	Cys	Ala	Ala	Gly
	610					615					620				
Asn	Lys	Val	Val	Asp	Asp	Cys	Tyr	Ile	Pro	Asn	Tyr	Leu	Leu	Pro	Lys
625					630					635					640
Ala	Cys	Cys	Cys	Ala	Gly	Cys	Cys	Cys	Gly	Ala	Gly	Cys	Thr	Cys	Cys
				645					650					655	
Ala	Gly	Thr	Gly	Gly	Gly	Cys	Gly	Thr	Gly	Gly	Ala	Cys	Thr	Ala	Ala
			660					665					670		
Cys	Ala	Thr	Gly	Gly	Ala	Cys	Cys	Ala	Gly	Thr	Ala	Thr	Ala	Thr	Ala
		675					680					685			
Thr	Gln	Pro	Glu	Leu	Gln	Trp	Ala	Trp	Thr	Asn	Met	Asp	Gln	Tyr	Ile
	690					695					700				
Ala	Gly	Cys	Gly	Cys	Cys	Thr	Gly	Thr	Thr	Thr	Gly	Ala	Ala	Thr	Cys
705					710					715					720
Thr	Cys	Gly	Cys	Gly	Gly	Ala	Gly	Cys	Gly	Thr	Ala	Ala	Ala	Cys	Gly
				725					730					735	
Gly	Cys	Thr	Gly	Gly	Thr	Gly	Gly	Cys	Gly	Cys	Ala	Gly	Cys	Ala	Thr

6/60

Cys Thr Thr Cys Cys Thr Gly Gly Gly Cys Thr Gly Gly Gly Cys Gly
1250 1255 1260
Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu Gly Trp Ala
1265 1270 1275 1280
Cys Ala Ala Ala Ala Gly Ala Ala Gly Thr Thr Cys Gly Gly Gly Ala
1285 1290 1295
Ala Gly Ala Gly Gly Ala Ala Cys Ala Cys Cys Ala Thr Cys Thr Gly
1300 1305 1310
Gly Cys Thr Cys Thr Thr Thr Gly Gly Gly Cys Cys Gly Gly Cys Cys
1315 1320 1325
Gln Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala
1330 1335 1340
Ala Cys Gly Ala Cys Gly Gly Gly Thr Ala Ala Ala Cys Cys Ala
1345 1350 1355 1360
Ala Cys Ala Thr Cys Gly Cys Gly Gly Ala Ala Gly Cys Cys Ala Thr
1365 1370 1375
Cys Gly Cys Cys Cys Ala Cys Gly Cys Cys Gly Thr Gly Cys Cys Cys
1380 1385 1390
Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro
1395 1400 1405
Thr Thr Cys Thr Ala Cys Gly Gly Cys Thr Gly Cys Gly Thr Gly Ala
1410 1415 1420
Ala Cys Thr Gly Gly Ala Cys Cys Ala Ala Thr Gly Ala Gly Ala Ala
1425 1430 1435 1440
Cys Thr Thr Thr Cys Cys Gly Thr Thr Cys Ala Ala Cys Gly Ala Thr
1445 1450 1455
Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
1460 1465 1470
Thr Gly Cys Gly Thr Cys Gly Ala Cys Ala Ala Gly Ala Thr Gly Gly
1475 1480 1485
Thr Gly Ala Thr Cys Thr Gly Gly Thr Gly Gly Gly Ala Gly Gly Ala
1490 1495 1500
Gly Gly Gly Cys Ala Ala Gly Ala Thr Gly Ala Cys Gly Gly Cys Cys
1505 1510 1515 1520
Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala
1525 1530 1535
Ala Ala Gly Gly Thr Cys Gly Thr Ala Gly Ala Gly Ala Gly Cys Gly
1540 1545 1550
Cys Cys Ala Ala Gly Gly Cys Cys Ala Thr Cys Cys Thr Gly Gly Gly
1555 1560 1565
Cys Gly Gly Ala Ala Gly Cys Ala Ala Gly Gly Thr Gly Cys Gly Cys
1570 1575 1580
Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
1585 1590 1595 1600
Gly Thr Gly Gly Ala Cys Cys Ala Ala Ala Gly Thr Gly Cys Ala
1605 1610 1615
Ala Gly Thr Cys Ala Thr Cys Gly Gly Cys Cys Cys Ala Gly Ala Thr
1620 1625 1630
Cys Gly Ala Cys Cys Cys Ala Ala Cys Thr Cys Cys Cys Gly Thr Gly
1635 1640 1645
Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
1650 1655 1660
Ala Thr Cys Gly Thr Cys Ala Cys Cys Thr Cys Cys Ala Ala Cys Ala
1665 1670 1675 1680
Cys Cys Ala Ala Cys Ala Thr Gly Thr Gly Cys Gly Cys Gly Gly Thr
1685 1690 1695
Cys Ala Thr Cys Gly Ala Cys Gly Gly Ala Ala Ala Cys Thr Cys Gly
1700 1705 1710
Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
1715 1720 1725
Ala Cys Cys Ala Cys Cys Thr Thr Cys Gly Ala Gly Cys Ala Cys Cys
1730 1735 1740
Ala Ala Cys Ala Ala Cys Cys Ala Cys Thr Cys Cys Ala Gly Gly Ala

1745 Cys Cys Gly Gly Ala Thr Gly Thr Thr Cys Ala Ala Gly Thr Thr Cys 1750 1755 1760
 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 1765 1770 1775
 Gly Ala Gly Cys Thr Cys Ala Cys Cys Ala Ala Gly Cys Gly Cys Cys 1780 1785 1790
 Thr Gly Gly Ala Gly Cys Ala Cys Gly Ala Cys Thr Thr Thr Gly Gly 1795 1800 1805
 Cys Ala Ala Gly Gly Thr Cys Ala Cys Cys Ala Ala Gly Cys Ala Gly 1810 1815 1820
 Glu Leu Thr Lys Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 1825 1830 1835 1840
 Gly Ala Ala Gly Thr Cys Ala Ala Ala Gly Ala Cys Thr Thr Thr Thr 1845 1850 1855
 Thr Cys Cys Gly Gly Thr Gly Gly Gly Cys Gly Thr Cys Ala Gly Ala 1860 1865 1870
 Thr Cys Ala Cys Gly Thr Gly Ala Cys Cys Gly Ala Gly Gly Thr Gly 1875 1880 1885
 Glu Val Lys Asp Phe Phe Arg Trp Ala Ser Asp His Val Thr Glu Val 1890 1895 1900
 Ala Cys Thr Cys Ala Cys Gly Ala Gly Thr Thr Thr Thr Ala Cys Gly 1905 1910 1915 1920
 Thr Cys Ala Gly Ala Ala Ala Gly Gly Gly Thr Gly Gly Ala Gly Cys 1925 1930 1935
 Thr Ala Gly Ala Ala Ala Gly Ala Gly Gly Cys Cys Cys Gly Cys Cys 1940 1945 1950
 Thr His Glu Phe Tyr Val Arg Lys Gly Gly Ala Arg Lys Arg Pro Ala 1955 1960 1965
 Cys Cys Cys Ala Ala Thr Gly Ala Cys Gly Cys Ala Gly Ala Thr Ala 1970 1975 1980
 Thr Ala Ala Gly Thr Gly Ala Gly Cys Cys Cys Ala Ala Gly Cys Gly 1985 1990 1995 2000
 Gly Gly Cys Cys Thr Gly Thr Cys Cys Gly Thr Cys Ala Gly Thr Thr 2005 2010 2015
 Pro Asn Asp Ala Asp Ile Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 2020 2025 2030
 Gly Cys Gly Cys Ala Gly Cys Cys Ala Thr Cys Gly Ala Cys Gly Thr 2035 2040 2045
 Cys Ala Gly Ala Cys Gly Cys Gly Gly Ala Ala Gly Cys Thr Cys Cys 2050 2055 2060
 Gly Gly Thr Gly Gly Ala Cys Thr Ala Cys Gly Cys Gly Gly Ala Cys 2065 2070 2075 2080
 Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Pro Val Asp Tyr Ala Asp 2085 2090 2095
 Ala Gly Gly Thr Ala Cys Cys Ala Ala Ala Ala Cys Ala Ala Ala Thr 2100 2105 2110
 Gly Thr Thr Cys Thr Cys Gly Thr Cys Ala Cys Gly Thr Gly Gly Gly 2115 2120 2125
 Thr Ala Thr Gly Ala Ala Thr Cys Thr Gly Ala Thr Gly Cys Thr Thr 2130 2135 2140
 Arg Tyr Gln Asn Lys Cys Ser Arg His Val Gly Met Asn Leu Met Leu 2145 2150 2155 2160
 Thr Thr Thr Cys Cys Cys Thr Gly Cys Cys Gly Gly Cys Ala Ala Thr 2165 2170 2175
 Gly Cys Gly Ala Gly Ala Gly Ala Ala Thr Gly Ala Ala Thr Cys Ala 2180 2185 2190
 Gly Ala Ala Thr Gly Thr Gly Gly Ala Cys Ala Thr Thr Thr Gly Cys 2195 2200 2205
 Phe Pro Cys Arg Gln Cys Glu Arg Met Asn Gln Asn Val Asp Ile Cys 2210 2215 2220
 Thr Thr Cys Ala Cys Gly Cys Ala Cys Gly Gly Gly Gly Thr Cys Ala 2225 2230 2235 2240
 2245 2250 2255

Thr Gly Gly Ala Cys Thr Gly Thr Gly Cys Cys Gly Ala Gly Thr Gly
 2260 2265 2270
 Cys Thr Thr Cys Cys Cys Cys Gly Thr Gly Thr Cys Ala Gly Ala Ala
 2275 2280 2285
 Phe Thr His Gly Val Met Asp Cys Ala Glu Cys Phe Pro Val Ser Glu
 2290 2295 2300
 Thr Cys Thr Cys Ala Ala Cys Cys Cys Gly Thr Gly Thr Cys Thr Gly
 2305 2310 2315 2320
 Thr Cys Gly Thr Cys Ala Gly Ala Ala Ala Gly Cys Gly Gly Ala Cys
 2325 2330 2335
 Gly Thr Ala Thr Cys Ala Gly Ala Ala Ala Cys Thr Gly Thr Gly Thr
 2340 2345 2350
 Ser Gln Pro Val Ser Val Val Arg Lys Arg Thr Tyr Gln Lys Leu Cys
 2355 2360 2365
 Cys Cys Gly Ala Thr Thr Cys Ala Thr Cys Ala Cys Ala Thr Cys Ala
 2370 2375 2380
 Thr Gly Gly Gly Gly Ala Gly Gly Gly Cys Gly Cys Cys Cys Gly Ala
 2385 2390 2395 2400
 Gly Gly Thr Gly Gly Cys Cys Thr Gly Cys Thr Cys Gly Gly Cys Cys
 2405 2410 2415
 Pro Ile His His Ile Met Gly Arg Ala Pro Glu Val Ala Cys Ser Ala
 2420 2425 2430
 Thr Gly Cys Gly Ala Ala Cys Thr Gly Gly Cys Cys Ala Ala Thr Gly
 2435 2440 2445
 Thr Gly Ala Cys Thr Thr Gly Gly Ala Thr Gly Ala Cys Thr Gly
 2450 2455 2460
 Thr Gly Ala Cys Ala Thr Gly Gly Ala Ala Cys Ala Ala Thr Ala Ala
 2465 2470 2475 2480
 Cys Glu Leu Ala Asn Val Asp Leu Asp Asp Cys Asp Met Glu Gln
 2485 2490 2495

<210> 4
 <211> 734
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 4
 Met Thr Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser Glu
 1 5 10 15
 Gly Val Arg Glu Trp Trp Ala Leu Gln Pro Gly Ala Pro Lys Pro Lys
 20 25 30
 Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro Gly
 35 40 45
 Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro Val
 50 55 60
 Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp Gln
 65 70 75 80
 Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala Asp
 85 90 95
 Ala Glu Phe Gln Gln Arg Leu Gln Gly Asp Thr Ser Phe Gly Gly Asn
 100 105 110
 Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro Leu
 115 120 125
 Gly Leu Val Glu Gln Ala Gly Glu Thr Ala Pro Gly Lys Lys Arg Pro
 130 135 140
 Leu Ile Glu Ser Pro Gln Gln Pro Asp Ser Ser Thr Gly Ile Gly Lys
 145 150 155 160
 Lys Gly Lys Gln Pro Ala Lys Lys Lys Leu Val Phe Glu Asp Glu Thr
 165 170 175

Gly	Ala	Gly	Asp	Gly	Pro	Pro	Glu	Gly	Ser	Thr	Ser	Gly	Ala	Met	Ser
Asp	Asp	Ser	Glu	Met	Arg	Ala	Ala	Ala	Gly	Gly	Ala	Ala	Val	Glu	Gly
Gly	Gln	Gly	Ala	Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly	Asp	Trp	His	Cys
Asp	Ser	Thr	Trp	Ser	Glu	Gly	His	Val	Thr	Thr	Thr	Ser	Thr	Arg	Thr
225	Val	Leu	Pro	Thr	Tyr	Asn	Asn	His	Leu	Tyr	Lys	Arg	Leu	Gly	Glu
Trp	Leu	Gln	Ser	Asn	Thr	Tyr	Asn	Gly	Phe	Ser	Thr	Pro	Trp	Gly	Tyr
Ser	Leu	Gln	Ser	Asn	Arg	Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp
Phe	Asp	Phe	Asn	Asn	Asn	Trp	Gly	Met	Arg	Pro	Lys	Ala	Met	Arg	Val
Arg	Leu	Ile	Asn	Asn	Asn	Trp	Gly	Met	Arg	Pro	Lys	Ala	Met	Arg	Val
Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Thr	Ser	Asn	Gly	Glu
305	Thr	Val	Ala	Asn	Asn	Leu	Thr	Ser	Thr	Val	Gln	Ile	Phe	Ala	Asp
Thr	Thr	Val	Ala	Asn	Asn	Leu	Thr	Ser	Thr	Val	Gln	Ile	Phe	Ala	Asp
Ser	Ser	Tyr	Glu	Leu	Pro	Tyr	Val	Met	Asp	Ala	Gly	Gln	Glu	Gly	Ser
Leu	Pro	Pro	Phe	Pro	Asn	Asp	Val	Phe	Met	Val	Pro	Gln	Tyr	Gly	Tyr
Cys	Gly	Leu	Val	Thr	Gly	Asn	Thr	Ser	Gln	Gln	Gln	Thr	Asp	Arg	Asn
Ala	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe	Pro	Ser	Gln	Met	Leu	Arg	Thr	Gly
385	Asn	Asn	Phe	Glu	Ile	Thr	Tyr	Ser	Phe	Glu	Lys	Val	Pro	Phe	His
Asn	Asn	Phe	Glu	Ile	Thr	Tyr	Ser	Phe	Glu	Lys	Val	Pro	Phe	His	Ser
Met	Tyr	Ala	His	Ser	Gln	Ser	Leu	Asp	Arg	Leu	Met	Asn	Pro	Leu	Ile
Asp	Gln	Tyr	Leu	Trp	Gly	Leu	Gln	Ser	Thr	Thr	Thr	Gly	Thr	Thr	Leu
Asn	Ala	Gly	Thr	Ala	Thr	Thr	Asn	Phe	Thr	Lys	Leu	Arg	Pro	Thr	Asn
450	Phe	Ser	Asn	Phe	Lys	Lys	Asn	Trp	Leu	Pro	Gly	Pro	Ser	Ile	Lys
465	Gln	Gly	Phe	Ser	Lys	Thr	Ala	Asn	Gln	Asn	Tyr	Lys	Ile	Pro	Ala
Gln	Gly	Phe	Ser	Lys	Thr	Ala	Asn	Gln	Asn	Tyr	Lys	Ile	Pro	Ala	Thr
Gly	Ser	Asp	Ser	Leu	Ile	Lys	Tyr	Glu	Thr	His	Ser	Thr	Leu	Asp	Gly
Arg	Trp	Ser	Ala	Leu	Thr	Pro	Gly	Pro	Pro	Met	Ala	Thr	Ala	Gly	Pro
Ala	Asp	Ser	Lys	Phe	Ser	Asn	Ser	Gln	Leu	Ile	Phe	Ala	Gly	Pro	Lys
530	Gln	Asn	Gly	Asn	Thr	Ala	Thr	Val	Pro	Gly	Thr	Leu	Ile	Phe	Thr
545	Glu	Glu	Glu	Leu	Ala	Ala	Thr	Asn	Ala	Thr	Asp	Thr	Asp	Met	Trp
Glu	Glu	Glu	Leu	Ala	Ala	Thr	Asn	Ala	Thr	Asp	Thr	Asp	Met	Trp	Gly
Asn	Leu	Pro	Gly	Gly	Asp	Gln	Ser	Asn	Ser	Asn	Leu	Pro	Thr	Val	Asp
Arg	Leu	Thr	Ala	Leu	Gly	Ala	Val	Pro	Gly	Met	Val	Trp	Gln	Asn	Arg
Asp	Ile	Tyr	Tyr	Gln	Gly	Pro	Ile	Trp	Ala	Lys	Ile	Pro	His	Thr	Asp
610	Gly	His	Phe	His	Pro	Ser	Pro	Leu	Ile	Gly	Gly	Phe	Gly	Leu	Lys
625	Pro	Pro	Pro	Gln	Ile	Phe	Ile	Lys	Asn	Thr	Pro	Val	Pro	Ala	Asn
Pro	Pro	Pro	Gln	Ile	Phe	Ile	Lys	Asn	Thr	Pro	Val	Pro	Ala	Asn	Pro
Ala	Thr	Thr	Phe	Ser	Ser	Thr	Pro	Val	Asn	Ser	Phe	Ile	Thr	Gln	Tyr
Ser	Thr	Gly	Gln	Val	Ser	Val	Gln	Ile	Asp	Trp	Glu	Ile	Gln	Lys	Glu
Ser	Thr	Gly	Gln	Val	Ser	Val	Gln	Ile	Asp	Trp	Glu	Ile	Gln	Lys	Glu

<400> 5						
atgactgacg	gttaccttcc	agattggcta	gaggacaacc	tctctgaagg	cgttcgagag	60
tggtggg'gc	tgcaacctgg	agcccctaaa	cccaaggcaa	atcaacaaca	tcaggacaac	120
gctcggggtc	ttgtgcttcc	gggttacaaa	tacctcggac	ccggcaacgg	actcgacaag	180
ggggaacccg	tcaacgcagc	ggacgcggca	gccctcgagc	acgacaaggc	ctacgaccag	240
cagctcaagg	ccggtgacaa	cccctacctc	aagtacaacc	acgccgacgc	ggagttccag	300
cagcggcttc	agggcgacac	atcgtttggg	ggcaacctcg	gcagagcagt	cttccaggcc	360
aaaaagaggg	ttcttgaacc	tcttggtctg	gttgagcaag	cgggtgagac	ggctcctgga	420
aagaagagac	cgttgattga	atccccccag	cagcccgact	cctccacggg	tatcggcaaa	480
aaaggcaagc	agccggctaa	aaagaagctc	gttttcgaag	acgaaactgg	agcaggcgac	540
ggacccccctg	agggatcaac	ttccggagcc	atgtctgatg	acagtgagat	gcgtgcagca	600
gctggcggag	ctgcagtcga	ggsggacaa	ggtgccgatg	gagtgggtaa	tgctcgggt	660
gattggcatt	gcgattccac	ctggtctgag	ggccacgtca	cgaccaccag	caccagaacc	720
tgggtcttgc	ccacctacaa	caaccacctn	tacaagcgac	tcggagagag	cctgcagtcc	780
aacacctaca	acggattctc	caccccctgg	ggatactttg	acttcaaccg	cttccactgc	840
cacttctcac	cacgtgactg	gcagcgactc	atcaacaaca	actggggcat	gcgacccaaa	900
gccatgcggg	tcaaaatctt	caacatccag	gtcaaggagg	tcacgacgtc	gaacggcgag	960
acaacggtgg	ctaataacct	taccagcacg	gttcagatct	ttgcggactc	gtcgtacgaa	1020
ctgccgtacg	tgatggatgc	gggtcaagag	ggcagcctgc	ctccttttcc	caacgacgtc	1080
tttatggtgc	cccagtacgg	ctactgtgga	ctggtgaccg	gcaacacttc	gcagcaacag	1140
actgacagaa	atgccttcta	ctgcctggag	tactttcctt	cgcagatgct	gcggactggc	1200
aacaactttg	aaattacgta	cagttttgag	aaggtgcctt	tccactcgat	gtacgcgcac	1260
agccagagcc	tggaaccgct	gatgaaccct	ctcatcgacc	agtacctgtg	gggactgcaa	1320
tcgaccacca	ccggaaccac	cctgaatgcc	gggactgcca	ccaccaactt	taccaagctg	1380
cggcctacca	actttttcaa	ctttaaaaaag	aactggctgc	ccgggccttc	aatcaagcag	1440
cagggcttct	caaagactgc	caatcaaaac	tacaagatcc	ctgccaccgg	gtcagacagt	1500
ctcatcaaat	acgagacgca	cagcactctg	gacggaagat	ggagtgcctt	gacccccgga	1560
cctccaatgg	ccacggctgg	acctgcggac	agcaagttca	gcaacagcca	gctcatcttt	1620
gcggggccta	aacagaacgg	caacacggcc	accgtaccgg	ggactctgat	cttcacctct	1680
gaggaggagc	tggcagccac	caacgccacc	gatacggaca	tgtggggcaa	cctacctggc	1740
ggtgaccaga	gcaacagcaa	cctgccgacc	gtggacagac	tgacagcctt	gggagccgtg	1800
cctggaatgg	tctggcaaaa	cagagacatt	tactaccagg	gtcccatttg	ggccaagatt	1860
cctcataccg	atggacactt	tcacccctca	ccgctgattg	gtgggttttg	gctgaaacac	1920
ccgcctcctc	aaatttttat	caagaacacc	ccggtacctg	cgaatcctgc	aacgaccttc	1980
agctctactc	cggtaaactc	cttcattact	cagtacagca	ctggccagggt	gtcgggtgcag	2040
attgactggg	agatccagaa	ggagcgggtc	aaacgctgga	accccgagggt	ccagtttacc	2100
tccaactacg	gacagcaaaa	ctctctgttg	tgggctcccg	atgcggctgg	gaaatacact	2160
gagcctaggg	ctatcgggtac	ccgctacctc	accaccacc	tgtataaa		2208

<210> 6
 <211> 125
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 6
 ttggccactc cctctatgcg cgctcgctca ctactcggc cctggagacc aaaggtctcc 60
 agactgccg cctctggccg gcagggccga gtgagtgagc gagcgcgcat agaggggagtg 120
 gccaa 125

<210> 7
 <211> 245
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 7
 ctccatcatc taggtttgcc cactgacgtc aatgtgacgt cctaggggta gggagggtccc 60
 tgtattagca gtcacgtgag tgtcgtattt cgcgagcgt agcgagcgc ataccaagct 120
 gccacgtcac agccacgtgg tccgtttgcg acagtttgcg acaccatgtg gtcaggaggg 180
 tatataaccg cgagtgagcc agcgaggagc tccattttgc ccgcgaattt tgaacgagca 240
 gcagc 245

<210> 8
 <211> 313
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 8
 Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys
 1 5 10 15
 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala
 20 25 30
 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Ser Lys
 35 40 45
 Ile Met Ser Leu Thr Lys Thr Ala Pro Asp Tyr Leu Val Gly Gln Asn
 50 55 60
 Pro Pro Glu Asp Ile Ser Ser Asn Arg Ile Tyr Arg Ile Leu Glu Met
 65 70 75 80
 Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu Gly Trp Ala
 85 90 95
 Gln Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala
 100 105 110
 Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro
 115 120 125
 Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
 130 135 140
 Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala
 145 150 155 160
 Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
 165 170 175

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
 180 185 190
 Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
 195 200 205
 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
 210 215 220
 Glu Leu Thr Lys Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln
 225 230 235 240
 Glu Val Lys Asp Phe Phe Arg Trp Ala Ser Asp His Val Thr Glu Val
 245 250 255
 Thr His Glu Phe Tyr Val Arg Lys Gly Gly Ala Arg Lys Arg Pro Ala
 260 265 270
 Pro Asn Asp Ala Asp Ile Ser Glu Pro Lys Arg Ala Cys Pro Ser Val
 275 280 285
 Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Pro Val Asp Tyr Ala Asp
 290 295 300
 Arg Leu Ala Arg Gly Gln Pro Leu Xaa
 305 310

<210> 9

<211> 399

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 9

Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys
 1 5 10 15
 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala
 20 25 30
 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Ser Lys
 35 40 45
 Ile Met Ser Leu Thr Lys Thr Ala Pro Asp Tyr Leu Val Gly Gln Asn
 50 55 60
 Pro Pro Glu Asp Ile Ser Ser Asn Arg Ile Tyr Arg Ile Leu Glu Met
 65 70 75 80
 Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu Gly Trp Ala
 85 90 95
 Gln Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala
 100 105 110
 Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro
 115 120 125
 Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
 130 135 140
 Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala
 145 150 155 160
 Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
 165 170 175
 Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
 180 185 190
 Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
 195 200 205
 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
 210 215 220
 Glu Leu Thr Lys Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln
 225 230 235 240
 Glu Val Lys Asp Phe Phe Arg Trp Ala Ser Asp His Val Thr Glu Val
 245 250 255
 Thr His Glu Phe Tyr Val Arg Lys Gly Gly Ala Arg Lys Arg Pro Ala
 260 265 270

Pro Asn Asp Ala Asp Ile Ser Glu Pro Lys Arg Ala Cys Pro Ser Val
 275 280 285
 Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Pro Val Asp Tyr Ala Asp
 290 295 300
 Arg Tyr Gln Asn Lys Cys Ser Arg His Val Gly Met Asn Leu Met Leu
 305 310 315 320
 Phe Pro Cys Arg Gln Cys Glu Arg Met Asn Gln Asn Val Asp Ile Cys
 325 330 335
 Phe Thr His Gly Val Met Asp Cys Ala Glu Cys Phe Pro Val Ser Glu
 340 345 350
 Ser Gln Pro Val Ser Val Val Arg Lys Arg Thr Tyr Gln Lys Leu Cys
 355 360 365
 Pro Ile His His Ile Met Gly Arg Ala Pro Glu Val Ala Cys Ser Ala
 370 375 380
 Cys Glu Leu Ala Asn Val Asp Leu Asp Asp Cys Asp Met Glu Gln
 385 390 395

<210> 10

<211> 537

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 10

Met Pro Gly Phe Tyr Glu Ile Val Leu Lys Val Pro Ser Asp Leu Asp
 1 5 10 15
 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu
 20 25 30
 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
 35 40 45
 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Glu Phe Leu
 50 55 60
 Val Glu Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val
 65 70 75 80
 Gln Phe Glu Lys Gly Asp Ser Tyr Phe His Leu His Ile Leu Val Glu
 85 90 95
 Thr Val Gly Val Lys Ser Met Val Val Gly Arg Tyr Val Ser Gln Ile
 100 105 110
 Lys Glu Lys Leu Val Thr Arg Ile Tyr Arg Gly Val Glu Pro Gln Leu
 115 120 125
 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly
 130 135 140
 Asn Lys Val Val Asp Asp Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys
 145 150 155 160
 Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Asp Gln Tyr Ile
 165 170 175
 Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His
 180 185 190
 Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Gln Asn
 195 200 205
 Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr
 210 215 220
 Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys
 225 230 235 240
 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala
 245 250 255
 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Ser Lys
 260 265 270
 Ile Met Ser Leu Thr Lys Thr Ala Pro Asp Tyr Leu Val Gly Gln Asn
 275 280 285

Pro Pro Glu Asp Ile Ser Ser Asn Arg Ile Tyr Arg Ile Leu Glu Met
 290 295 300
 Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu Gly Trp Ala
 305 310 315 320
 Gln Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala
 325 330 335
 Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro
 340 345 350
 Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
 355 360 365
 Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala
 370 375 380
 Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
 385 390 395 400
 Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
 405 410 415
 Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
 420 425 430
 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
 435 440 445
 Glu Leu Thr Lys Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln
 450 455 460
 Glu Val Lys Asp Phe Phe Arg Trp Ala Ser Asp His Val Thr Glu Val
 465 470 475 480
 Thr His Glu Phe Tyr Val Arg Lys Gly Gly Ala Arg Lys Arg Pro Ala
 485 490 495
 Pro Asn Asp Ala Asp Ile Ser Glu Pro Lys Arg Ala Cys Pro Ser Val
 500 505 510
 Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Pro Val Asp Tyr Ala Asp
 515 520 525
 Arg Leu Ala Arg Gly Gln Pro Leu Xaa
 530 535

<210> 11

<211> 623

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 11

Met Pro Gly Phe Tyr Glu Ile Val Leu Lys Val Pro Ser Asp Leu Asp
 1 5 10 15
 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu
 20 25 30
 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
 35 40 45
 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Glu Phe Leu
 50 55 60
 Val Glu Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val
 65 70 75 80
 Gln Phe Glu Lys Gly Asp Ser Tyr Phe His Leu His Ile Leu Val Glu
 85 90 95
 Thr Val Gly Val Lys Ser Met Val Val Gly Arg Tyr Val Ser Gln Ile
 100 105 110
 Lys Glu Lys Leu Val Thr Arg Ile Tyr Arg Gly Val Glu Pro Gln Leu
 115 120 125
 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly
 130 135 140
 Asn Lys Val Val Asp Asp Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys
 145 150 155 160

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Asp Gln Tyr Ile
 165 170 175
 Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His
 180 185 190
 Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Gln Asn
 195 200 205
 Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr
 210 215 220
 Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys
 225 230 235 240
 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala
 245 250 255
 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Ser Lys
 260 265 270
 Ile Met Ser Leu Thr Lys Thr Ala Pro Asp Tyr Leu Val Gly Gln Asn
 275 280 285
 Pro Pro Glu Asp Ile Ser Ser Asn Arg Ile Tyr Arg Ile Leu Glu Met
 290 295 300
 Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu Gly Trp Ala
 305 310 315 320
 Gln Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala
 325 330 335
 Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro
 340 345 350
 Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
 355 360 365
 Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala
 370 375 380
 Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
 385 390 395 400
 Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
 405 410 415
 Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
 420 425 430
 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
 435 440 445
 Glu Leu Thr Lys Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln
 450 455 460
 Glu Val Lys Asp Phe Phe Arg Trp Ala Ser Asp His Val Thr Glu Val
 465 470 475 480
 Thr His Glu Phe Tyr Val Arg Lys Gly Gly Ala Arg Lys Arg Pro Ala
 485 490 495
 Pro Asn Asp Ala Asp Ile Ser Glu Pro Lys Arg Ala Cys Pro Ser Val
 500 505 510
 Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Pro Val Asp Tyr Ala Asp
 515 520 525
 Arg Tyr Gln Asn Lys Cys Ser Arg His Val Gly Met Asn Leu Met Leu
 530 535 540
 Phe Pro Cys Arg Gln Cys Glu Arg Met Asn Gln Asn Val Asp Ile Cys
 545 550 555 560
 Phe Thr His Gly Val Met Asp Cys Ala Glu Cys Phe Pro Val Ser Glu
 565 570 575
 Ser Gln Pro Val Ser Val Val Arg Lys Arg Thr Tyr Gln Lys Leu Cys
 580 585 590
 Pro Ile His His Ile Met Gly Arg Ala Pro Glu Val Ala Cys Ser Ala
 595 600 605
 Cys Glu Leu Ala Asn Val Asp Leu Asp Asp Cys Asp Met Glu Gln
 610 615 620

<210> 12
 <211> 939
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 12
 atggagctgg tcgggtggct ggtggaccgc gggatcacgt cagaaaagca atggatccag 60
 gaggaccagg cgtcctacat ctcttcaac gccgcctcca actcgcggtc acaaatcaag 120
 gccgcgctgg acaatgcctc caaaatcatg agcctgacaa agacggctcc ggactacctg 180
 gtggggccaga acccgccgga ggacatttcc agcaaccgca tctaccgaat cctcgagatg 240
 aacgggtacg atccgcagta cgcggcctcc gtcttcttgg gctgggcgca aaagaagtgc 300
 gggaagagga acaccatctg gctcttttgg ccggccacga cgggtaaaac caacatcgcg 360
 gaagccatcg cccacgcccgt gcccttctac ggctgcgtga actggaccaa tgagaacttt 420
 ccgttcaacg attgcgtcga caagatgggt atctgggtgg aggagggcaa gatgacggcc 480
 aaggtcgtag agagcgccaa ggccatcctg ggcggaagca aggtgcgcgt ggaccaaag 540
 tgcaagtcat cggcccagat cgacccaact cccgtgatcg tcacctcaa caccaacatg 600
 tgcgcggtca tcgacggaaa ctcgaccacc ttcgagcacc aacaaccact ccaggaccgg 660
 atgttcaagt tcgagctcac caagcgctg gagcacgact ttggcaaggc caccaagcag 720
 gaagtcaaag actttttccg gtgggcgtca gatcacgtga ccgaggtgac tcacgagttt 780
 tacgtcagaa aggggtggagc tagaaagagg cccgccccca atgacgcaga tataagttag 840
 cccaagcggg cctgtccgtc agttgcgcag ccacgcacgt cagacgcgga agctccggtg 900
 gactacgcgg acagattggc tagaggacaa cctctctga 939

<210> 13
 <211> 1197
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 13
 atggagctgg tcgggtggct ggtggaccgc gggatcacgt cagaaaagca atggatccag 60
 gaggaccagg cgtcctacat ctcttcaac gccgcctcca actcgcggtc acaaatcaag 120
 gccgcgctgg acaatgcctc caaaatcatg agcctgacaa agacggctcc ggactacctg 180
 gtggggccaga acccgccgga ggacatttcc agcaaccgca tctaccgaat cctcgagatg 240
 aacgggtacg atccgcagta cgcggcctcc gtcttcttgg gctgggcgca aaagaagtgc 300
 gggaagagga acaccatctg gctcttttgg ccggccacga cgggtaaaac caacatcgcg 360
 gaagccatcg cccacgcccgt gcccttctac ggctgcgtga actggaccaa tgagaacttt 420
 ccgttcaacg attgcgtcga caagatgggt atctgggtgg aggagggcaa gatgacggcc 480
 aaggtcgtag agagcgccaa ggccatcctg ggcggaagca aggtgcgcgt ggaccaaag 540
 tgcaagtcat cggcccagat cgacccaact cccgtgatcg tcacctcaa caccaacatg 600
 tgcgcggtca tcgacggaaa ctcgaccacc ttcgagcacc aacaaccact ccaggaccgg 660
 atgttcaagt tcgagctcac caagcgctg gagcacgact ttggcaaggc caccaagcag 720
 gaagtcaaag actttttccg gtgggcgtca gatcacgtga ccgaggtgac tcacgagttt 780
 tacgtcagaa aggggtggagc tagaaagagg cccgccccca atgacgcaga tataagttag 840
 cccaagcggg cctgtccgtc agttgcgcag ccacgcacgt cagacgcgga agctccggtg 900
 gactacgcgg acaggtacca aaacaaatgt tctcgtcacg tgggtatgaa tctgatgctt 960
 tttccctgcc ggcaatgcga gagaatgaat cagaatgtgg acatttgctt cacgcacggg 1020
 gtcattggact gtgccgagtg cttccccgtg tcagaatctc aaccggtgtc tgcgtcaga 1080
 aagcggacgt atcagaaact gtgtccgatt catcacatca tggggagggc gcccgagggtg 1140
 gcctgctcgg cctgcgaact ggccaatgtg gacttggatg actgtgacat ggaacaa 1197

<210> 14
 <211> 1611
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 14

atgccggggt	tctacgagat	cgtgctgaag	gtgcccagcg	acctggacga	gcacctgccc	60
ggcatttctg	actcttttgt	gagctgggtg	gccgagaagg	aatgggagct	gccgccggat	120
tctgacatgg	acttgaatct	gattgagcag	gcacccctga	ccgtggccga	aaagctgcaa	180
cgcgagttcc	tggtcgagtg	gcgccgcgtg	agtaaggccc	cggaggccct	cttctttgtc	240
cagttcgaga	agggggacag	ctacttccac	ctgcacatcc	tggtggagac	cgtgggcgtc	300
aaatccatgg	tggtgggccc	ctacgtgagc	cagattaaag	agaagctggt	gacccgcac	360
taccgcgggg	tcgagccgca	gcttccgaac	tggttcgcgg	tgaccaagac	gcgtaatggc	420
gccggaggcg	ggaacaaggt	ggtggacgac	tgctacatcc	ccaactacct	gctccccaag	480
accagccccg	agctccagtg	ggcgtggact	aacatggacc	agtatataag	cgcctgtttg	540
aatctcgcgg	agcgtaaacc	gctggtggcg	cagcatctga	cgcacgtgtc	gcagacgcag	600
gagcagaaca	aggaaaacca	gaaccccaat	tctgacgcgc	cggatcatcag	gtcaaaaacc	660
tccgccaggt	acatggagct	ggtcgggtgg	ctggtggacc	gcgggatcac	gtcagaaaag	720
caatggatcc	aggaggacca	ggcgtcctac	atctccttca	acgccgcctc	caactcgcgg	780
tcacaaatca	aggccgcgct	ggacaatgcc	tccaaaatca	tgagcctgac	aaagacggct	840
ccggactacc	tggtgggcca	gaacccgccc	gaggacattt	ccagcaaccg	catctaccga	900
atcctcgaga	tgaacgggta	cgatccgcag	tacgcggcct	ccgtcttcct	gggctgggcg	960
caaaagaagt	tcgggaagag	gaacaccatc	tggtctcttg	ggccggccac	gacgggtaaa	1020
accaacatcg	cgggaagccat	cgcccacgcc	gtgcccttct	acggctgcgt	gaactggacc	1080
aatgagaact	ttccgttcaa	cgattgcgtc	gacaagatgg	tgatctggtg	ggaggagggc	1140
aagatgacgg	ccaaggctcg	agagagcgcc	aaggccatcc	tgggcggaag	caaggtgcgc	1200
gtggaccaaa	agtgcagtc	atcggcccag	atcgacccaa	ctcccgtgat	cgtaacctcc	1260
aacaccaaca	tgtgcgcggt	catcgacgga	aactcgacca	ccttcgagca	ccaacaacca	1320
ctccaggacc	ggatgttcaa	gttcgagctc	accaagcgcc	tgagacacga	ctttggcaag	1380
gtcaccaagc	aggaagtcaa	agactttttc	cgggtgggct	cagatcacgt	gaccgaggtg	1440
actcacgagt	tttacgtcag	aaaggggtga	gctagaaaaga	ggcccgcgcc	caatgacgca	1500
gatataagtg	agcccaagcg	ggcctgtccg	tcagttgcgc	agccatcgac	gtcagacgcg	1560
gaagctccgg	tggactacgc	ggacagattg	gctagaggac	aacctctctg	a	1611

<210> 15

<211> 1872

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 15

atgccggggt	tctacgagat	cgtgctgaag	gtgcccagcg	acctggacga	gcacctgccc	60
ggcatttctg	actcttttgt	gagctgggtg	gccgagaagg	aatgggagct	gccgccggat	120
tctgacatgg	acttgaatct	gattgagcag	gcacccctga	ccgtggccga	aaagctgcaa	180
cgcgagttcc	tggtcgagtg	gcgccgcgtg	agtaaggccc	cggaggccct	cttctttgtc	240
cagttcgaga	agggggacag	ctacttccac	ctgcacatcc	tggtggagac	cgtgggcgtc	300
aaatccatgg	tggtgggccc	ctacgtgagc	cagattaaag	agaagctggt	gacccgcac	360
taccgcgggg	tcgagccgca	gcttccgaac	tggttcgcgg	tgaccaagac	gcgtaatggc	420
gccggaggcg	ggaacaaggt	ggtggacgac	tgctacatcc	ccaactacct	gctccccaag	480
accagccccg	agctccagtg	ggcgtggact	aacatggacc	agtatataag	cgcctgtttg	540
aatctcgcgg	agcgtaaacc	gctggtggcg	cagcatctga	cgcacgtgtc	gcagacgcag	600
gagcagaaca	aggaaaacca	gaaccccaat	tctgacgcgc	cggatcatcag	gtcaaaaacc	660
tccgccaggt	acatggagct	ggtcgggtgg	ctggtggacc	gcgggatcac	gtcagaaaag	720
caatggatcc	aggaggacca	ggcgtcctac	atctccttca	acgccgcctc	caactcgcgg	780
tcacaaatca	aggccgcgct	ggacaatgcc	tccaaaatca	tgagcctgac	aaagacggct	840
ccggactacc	tggtgggcca	gaacccgccc	gaggacattt	ccagcaaccg	catctaccga	900
atcctcgaga	tgaacgggta	cgatccgcag	tacgcggcct	ccgtcttcct	gggctgggcg	960
caaaagaagt	tcgggaagag	gaacaccatc	tggtctcttg	ggccggccac	gacgggtaaa	1020
accaacatcg	cgggaagccat	cgcccacgcc	gtgcccttct	acggctgcgt	gaactggacc	1080
aatgagaact	ttccgttcaa	cgattgcgtc	gacaagatgg	tgatctggtg	ggaggagggc	1140
aagatgacgg	ccaaggctcg	agagagcgcc	aaggccatcc	tgggcggaag	caaggtgcgc	1200
gtggaccaaa	agtgcagtc	atcggcccag	atcgacccaa	ctcccgtgat	cgtaacctcc	1260
aacaccaaca	tgtgcgcggt	catcgacgga	aactcgacca	ccttcgagca	ccaacaacca	1320
ctccaggacc	ggatgttcaa	gttcgagctc	accaagcgcc	tgagacacga	ctttggcaag	1380
gtcaccaagc	aggaagtcaa	agactttttc	cgggtgggct	cagatcacgt	gaccgaggtg	1440

actcacgagt	tttacgtcag	aaaggggtgga	gctagaaaga	ggcccgcccc	caatgacgca	1500
gatataagtg	agcccaagcg	ggcctgtccg	tcagttgcgc	agccatcgac	gtcagacgcg	1560
gaagctccgg	tggaactacgc	ggacaggtac	caaaacaaat	gttctcgtca	cgtgggtatg	1620
aatctgatgc	tttttccctg	ccggcaatgc	gagagaatga	atcagaatgt	ggacatttgc	1680
ttcacgcacg	gggtcatgga	ctgtgccgag	tgcttccccg	tgtcagaatc	tcaacccgtg	1740
tctgtcgtca	gaaagcggac	gtatcagaaa	ctgtgtccga	ttcatcacat	catggggagg	1800
gcgcccagg	tggcctgctc	ggcctgcgaa	ctggccaatg	tggaacttga	tgactgtgac	1860
atggaacaat	aa					1872

<210> 16

<211> 598

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 16

Thr	Ala	Pro	Gly	Lys	Lys	Arg	Pro	Leu	Ile	Glu	Ser	Pro	Gln	Gln	Pro
1				5					10					15	
Asp	Ser	Ser	Thr	Gly	Ile	Gly	Lys	Lys	Gly	Lys	Gln	Pro	Ala	Lys	Lys
			20					25					30		
Lys	Leu	Val	Phe	Glu	Asp	Glu	Thr	Gly	Ala	Gly	Asp	Gly	Pro	Pro	Glu
		35					40					45			
Gly	Ser	Thr	Ser	Gly	Ala	Met	Ser	Asp	Asp	Ser	Glu	Met	Arg	Ala	Ala
	50					55					60				
Ala	Gly	Gly	Ala	Ala	Val	Glu	Gly	Gly	Gln	Gly	Ala	Asp	Gly	Val	Gly
65					70					75					80
Asn	Ala	Ser	Gly	Asp	Trp	His	Cys	Asp	Ser	Thr	Trp	Ser	Glu	Gly	His
			85						90					95	
Val	Thr	Thr	Thr	Ser	Thr	Arg	Thr	Trp	Val	Leu	Pro	Thr	Tyr	Asn	Asn
			100					105					110		
His	Leu	Tyr	Lys	Arg	Leu	Gly	Glu	Ser	Leu	Gln	Ser	Asn	Thr	Tyr	Asn
		115					120					125			
Gly	Phe	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	Arg	Phe	His	Cys
	130					135					140				
His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	Asn	Asn	Trp	Gly
145					150					155					160
Met	Arg	Pro	Lys	Ala	Met	Arg	Val	Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys
			165					170						175	
Glu	Val	Thr	Thr	Ser	Asn	Gly	Glu	Thr	Val	Ala	Asn	Asn	Leu	Thr	
		180						185				190			
Ser	Thr	Val	Gln	Ile	Phe	Ala	Asp	Ser	Ser	Tyr	Glu	Leu	Pro	Tyr	Val
		195					200					205			
Met	Asp	Ala	Gly	Gln	Glu	Gly	Ser	Leu	Pro	Pro	Phe	Pro	Asn	Asp	Val
	210					215					220				
Phe	Met	Val	Pro	Gln	Tyr	Gly	Tyr	Cys	Gly	Leu	Val	Thr	Gly	Asn	Thr
225					230					235					240
Ser	Gln	Gln	Gln	Thr	Asp	Arg	Asn	Ala	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe
				245					250					255	
Pro	Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Glu	Ile	Thr	Tyr	Ser
			260					265					270		
Phe	Glu	Lys	Val	Pro	Phe	His	Ser	Met	Tyr	Ala	His	Ser	Gln	Ser	Leu
		275					280					285			
Asp	Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Trp	Gly	Leu	Gln
	290					295					300				
Ser	Thr	Thr	Thr	Gly	Thr	Thr	Leu	Asn	Ala	Gly	Thr	Ala	Thr	Thr	Asn
305					310					315					320
Phe	Thr	Lys	Leu	Arg	Pro	Thr	Asn	Phe	Ser	Asn	Phe	Lys	Lys	Asn	Trp
			325					330						335	
Leu	Pro	Gly	Pro	Ser	Ile	Lys	Gln	Gln	Gly	Phe	Ser	Lys	Thr	Ala	Asn
			340					345						350	

Gln Asn Tyr Lys Ile Pro Ala Thr Gly Ser Asp Ser Leu Ile Lys Tyr
 355 360 365
 Glu Thr His Ser Thr Leu Asp Gly Arg Trp Ser Ala Leu Thr Pro Gly
 370 375 380
 Pro Pro Met Ala Thr Ala Gly Pro Ala Asp Ser Lys Phe Ser Asn Ser
 385 390 395 400
 Gln Leu Ile Phe Ala Gly Pro Lys Gln Asn Gly Asn Thr Ala Thr Val
 405 410 415
 Pro Gly Thr Leu Ile Phe Thr Ser Glu Glu Glu Leu Ala Ala Thr Asn
 420 425 430
 Ala Thr Asp Thr Asp Met Trp Gly Asn Leu Pro Gly Gly Asp Gln Ser
 435 440 445
 Asn Ser Asn Leu Pro Thr Val Asp Arg Leu Thr Ala Leu Gly Ala Val
 450 455 460
 Pro Gly Met Val Trp Gln Asn Arg Asp Ile Tyr Tyr Gln Gly Pro Ile
 465 470 475 480
 Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro Leu
 485 490 495
 Ile Gly Gly Phe Gly Leu Lys His Pro Pro Pro Gln Ile Phe Ile Lys
 500 505 510
 Asn Thr Pro Val Pro Ala Asn Pro Ala Thr Thr Phe Ser Ser Thr Pro
 515 520 525
 Val Asn Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Gln
 530 535 540
 Ile Asp Trp Glu Ile Gln Lys Glu Arg Ser Lys Arg Trp Asn Pro Glu
 545 550 555 560
 Val Gln Phe Thr Ser Asn Tyr Gly Gln Gln Asn Ser Leu Leu Trp Ala
 565 570 575
 Pro Asp Ala Ala Gly Lys Tyr Thr Glu Pro Arg Ala Ile Gly Thr Arg
 580 585 590
 Tyr Leu Thr His His Leu
 595

<210> 17

<211> 1800

<212> DNA

<213> Artificial sequence

<220>

<223> Description of Artificial sequence; note =
synthetic construct

<221> misc_feature

<222> (0)...(0)

<223> n=a,t,c, or g

<221> variation

<222> (0)...(0)

<223> Xaa = any amino acid

<400> 17

acggctcctg	gaaagaagag	accgttgatt	gaatcccccc	agcagcccga	ctcctccacg	60
ggtatcggca	aaaaaggcaa	gcagccggct	aaaaagaagc	tcgttttcga	agacgaaact	120
ggagcaggcg	acggaccccc	tgagggatca	acttccggag	ccatgtctga	tgacagttag	180
atgcgtgcag	cagctggcgg	agctgcagtc	gagggsggac	aagggtgccga	tgagagtgggt	240
aatgcctcgg	gtgattggca	ttgcgattcc	acctgggtctg	agggccacgt	cacgaccacc	300
agcaccagaa	cctgggtcct	gcccacctac	aacaaccacc	tntacaagcg	actcggagag	360
agcctgcagt	ccaacaccta	caacggattc	tccaccccct	ggggatactt	tgacttcaac	420
cgcttccact	gccacttctc	accacgtgac	tggcagcgac	tcatacaaa	caactggggc	480
atgcgaccca	aagccatgcg	ggtcaaaatc	ttcaacatcc	aggtcaagga	ggtcacgacg	540
tcgaacggcg	agacaacggg	ggctaataac	cttaccagca	cggttcagat	ctttgcggac	600
tcgtcgtacg	aactgccgta	cgtgatggat	gcgggtcaag	agggcagcct	gcctcctttt	660
cccaacgacg	tctttatggg	gccccagtac	ggctactgtg	qactgggtgac	cggcaacact	720

tcgcagcaac	agactgacag	aaatgccttc	tactgcctgg	agtactttcc	ttcgcagatg	780
ctgCGgactg	gcaacaactt	tgaaattacg	tacagttttg	agaagggtgcc	tttccactcg	840
atgtacgcgc	acagccagag	cctggaccgg	ctgatgaacc	ctctcatcga	ccagtacctg	900
tggggactgc	aatcgaccac	caccggaacc	accctgaatg	ccgggactgc	caccaccaac	960
tttaccaagc	tgCGgcctac	caactttttc	aacttttaaaa	agaactggct	gcccgggcct	1020
tcaatcaagc	agcagggtt	ctcaaagact	gccaatcaaa	actacaagat	ccctgccacc	1080
gggtcagaca	gtctcatcaa	atacgagacg	cacagcactc	tggacggaag	atggagtgcc	1140
ctgacccccg	gacctccaat	ggccacggct	ggacctgcgg	acagcaagtt	cagcaacagc	1200
cagctcatct	ttgcggggcc	taaacagaac	ggcaacacgg	ccaccgtacc	cgggactctg	1260
atcttcacct	ctgaggagga	gctggcagcc	accaacgcca	ccgatacgga	catgtggggc	1320
aacctacctg	gCGgtgacca	gagcaacagc	aacctgccga	ccgtggacag	actgacagcc	1380
ttgggagccg	tgcctggaat	ggtctggcaa	aacagagaca	tttactacca	gggtcccatt	1440
tgggccaaga	ttcctcatac	cgatggacac	tttcaccctt	caccgctgat	tgggtgggtt	1500
gggctgaaac	acccgcctcc	tcaaattttt	atcaagaaca	ccccgggtacc	tgcgaatcct	1560
gcaacgacct	tcagctctac	tccggtaaac	tccttcatta	ctcagtacag	cactggccag	1620
gtgtcggtgc	agattgactg	ggagatccag	aaggagcggt	ccaaacgctg	gaaccccgag	1680
gtccagttta	cctccaacta	cggacagcaa	aactctctgt	tgtgggctcc	cgatgcggct	1740
gggaataaca	ctgagcctag	ggctatcggt	acccgctacc	tcaccaccca	cctgtaataa	1800

<210> 18

<211> 544

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 18

Met	Ser	Asp	Asp	Ser	Glu	Met	Arg	Ala	Ala	Ala	Gly	Gly	Ala	Ala	Val
1				5					10				15		
Glu	Gly	Gly	Gln	Gly	Ala	Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly	Asp	Trp
			20				25					30			
His	Cys	Asp	Ser	Thr	Trp	Ser	Glu	Gly	His	Val	Thr	Thr	Thr	Ser	Thr
		35				40					45				
Arg	Thr	Trp	Val	Leu	Pro	Thr	Tyr	Asn	Asn	His	Leu	Tyr	Lys	Arg	Leu
	50				55					60					
Gly	Glu	Ser	Leu	Gln	Ser	Asn	Thr	Tyr	Asn	Gly	Phe	Ser	Thr	Pro	Trp
65					70				75					80	
Gly	Tyr	Phe	Asp	Phe	Asn	Arg	Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp
			85					90					95		
Trp	Gln	Arg	Leu	Ile	Asn	Asn	Asn	Trp	Gly	Met	Arg	Pro	Lys	Ala	Met
			100					105					110		
Arg	Val	Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Thr	Ser	Asn
		115					120					125			
Gly	Glu	Thr	Thr	Val	Ala	Asn	Asn	Leu	Thr	Ser	Thr	Val	Gln	Ile	Phe
	130					135					140				
Ala	Asp	Ser	Ser	Tyr	Glu	Leu	Pro	Tyr	Val	Met	Asp	Ala	Gly	Gln	Glu
145					150				155					160	
Gly	Ser	Leu	Pro	Pro	Phe	Pro	Asn	Asp	Val	Phe	Met	Val	Pro	Gln	Tyr
			165					170					175		
Gly	Tyr	Cys	Gly	Leu	Val	Thr	Gly	Asn	Thr	Ser	Gln	Gln	Gln	Thr	Asp
		180						185				190			
Arg	Asn	Ala	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe	Pro	Ser	Gln	Met	Leu	Arg
		195				200						205			
Thr	Gly	Asn	Asn	Phe	Glu	Ile	Thr	Tyr	Ser	Phe	Glu	Lys	Val	Pro	Phe
	210					215					220				
His	Ser	Met	Tyr	Ala	His	Ser	Gln	Ser	Leu	Asp	Arg	Leu	Met	Asn	Pro
225					230					235				240	
Leu	Ile	Asp	Gln	Tyr	Leu	Trp	Gly	Leu	Gln	Ser	Thr	Thr	Thr	Gly	Thr
			245					250						255	
Thr	Leu	Asn	Ala	Gly	Thr	Ala	Thr	Thr	Asn	Phe	Thr	Lys	Leu	Arg	Pro
		260					265						270		

Thr Asn Phe Ser Asn Phe Lys Lys Asn Trp Leu Pro Gly Pro Ser Ile
 275 280 285
 Lys Gln Gln Gly Phe Ser Lys Thr Ala Asn Gln Asn Tyr Lys Ile Pro
 290 295 300
 Ala Thr Gly Ser Asp Ser Leu Ile Lys Tyr Glu Thr His Ser Thr Leu
 305 310 315 320
 Asp Gly Arg Trp Ser Ala Leu Thr Pro Gly Pro Pro Met Ala Thr Ala
 325 330 335
 Gly Pro Ala Asp Ser Lys Phe Ser Asn Ser Gln Leu Ile Phe Ala Gly
 340 345 350
 Pro Lys Gln Asn Gly Asn Thr Ala Thr Val Pro Gly Thr Leu Ile Phe
 355 360 365
 Thr Ser Glu Glu Glu Leu Ala Ala Thr Asn Ala Thr Asp Thr Asp Met
 370 375 380
 Trp Gly Asn Leu Pro Gly Gly Asp Gln Ser Asn Ser Asn Leu Pro Thr
 385 390 395 400
 Val Asp Arg Leu Thr Ala Leu Gly Ala Val Pro Gly Met Val Trp Gln
 405 410 415
 Asn Arg Asp Ile Tyr Tyr Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 420 425 430
 Thr Asp Gly His Phe His Pro Ser Pro Leu Ile Gly Gly Phe Gly Leu
 435 440 445
 Lys His Pro Pro Pro Gln Ile Phe Ile Lys Asn Thr Pro Val Pro Ala
 450 455 460
 Asn Pro Ala Thr Thr Phe Ser Ser Thr Pro Val Asn Ser Phe Ile Thr
 465 470 475 480
 Gln Tyr Ser Thr Gly Gln Val Ser Val Gln Ile Asp Trp Glu Ile Gln
 485 490 495
 Lys Glu Arg Ser Lys Arg Trp Asn Pro Glu Val Gln Phe Thr Ser Asn
 500 505 510
 Tyr Gly Gln Gln Asn Ser Leu Leu Trp Ala Pro Asp Ala Ala Gly Lys
 515 520 525
 Tyr Thr Glu Pro Arg Ala Ile Gly Thr Arg Tyr Leu Thr His His Leu
 530 535 540

<210> 19

<211> 1617

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<221> misc_feature

<222> (0)...(0)

<223> n=a,t,c, or g

<221> variation

<222> (0)...(0)

<223> Xaa = any amino acid

<400> 19

atgcgtgcag	cagctggcgg	agctgcagtc	gagggsggac	aaggtgccga	tggagtgggt	60
aatgcctcgg	gtgattggca	ttgcgattcc	acctggtctg	agggccacgt	cacgaccacc	120
agcaccagaa	cctgggtctt	gccacacctac	aacaaccacc	tntacaagcg	actcggagag	180
agcctgcagt	ccaacaccta	caacggattc	tccaccccct	ggggatactt	tgacttcaac	240
cgcttccact	gccacttctc	accacgtgac	tggcagcgac	tcatcaacaa	caactggggc	300
atgcgaccca	aagccatgcg	ggtcaaaaatc	ttcaacatcc	aggtcaagga	ggtcacgacg	360
tcgaacggcg	agacaacggt	ggctaataac	cttaccagca	cggttcagat	ctttgcggac	420
tcgtcgtacg	aactgccgta	cgtgatggat	gcgggtcaag	agggcagcct	gcctcctttt	480
cccaacgacg	tctttatggt	gccccagtac	ggctactgtg	gactgggtgac	cggcaacact	540
tcgcagcaac	agactgacag	aatgccttc	tactgcctgg	agtactttcc	ttcgcagatg	600

ctgcggactg	gcaacaactt	tgaaattacg	tacagrrrrg	agaagggtgcc	tttccactcg	660
atgtacgcgc	acagccagag	cctggaccgg	ctgatgaacc	ctctcatcga	ccagtacctg	720
tggggactgc	aatcgaccac	caccggaacc	accctgaatg	ccgggactgc	caccaccaac	780
tttaccaagc	tgcggcctac	caacttttcc	aacttttaaaa	agaactggct	gcccgggcct	840
tcaatcaagc	agcagggtt	ctcaaagact	gccaatcaaa	actacaagat	ccctgccacc	900
gggtcagaca	gtctcatcaa	atacgagacg	cacagcactc	tggacggaag	atggagtgcc	960
ctgacccccg	gacctccaat	ggccacggct	ggacctgcgg	acagcaagtt	cagcaacagc	1020
cagctcatct	ttgcggggcc	taaacagaac	ggcaacacgg	ccaccgtacc	cgggactctg	1080
atcttcacct	ctgaggagga	gctggcagcc	accaacgcca	ccgatacggg	catgtggggc	1140
aacctacctg	gcggtgacca	gagcaacagc	aacctgccga	ccgtggacag	actgacagcc	1200
ttgggagccg	tgcttggaa	ggtctggcaa	aacagagaca	tttactacca	gggtcccatt	1260
tggggccaaga	ttcctcatac	cgatggacac	tttcacccct	caccgctgat	tgggtgggttt	1320
gggctgaaac	acccgcctcc	tcaaattttt	atcaagaaca	ccccggtacc	tgcgaatcct	1380
gcaacgacct	tcagctctac	tccggtaaac	tccttcatta	ctcagtacag	cactggccag	1440
gtgtcggtgc	agattgactg	ggagatccag	aaggagcggt	ccaaacgctg	gaaccccgag	1500
gtccagttta	cctccaacta	cggacagcaa	aactctctgt	tgtgggctcc	cgatgcggct	1560
gggaaataca	ctgagcctag	ggctatcggt	acccgctacc	tcacccacca	cctgtaa	1617

<210> 20

<211> 129

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 20

ttggccactc	cctctatgcg	cgctcgctca	ctcactcggc	cctgcggcca	gaggccggca	60
gtctggagac	ctttggtgtc	cagggcaggg	ccgagtgagt	gagcgagcgc	gcatagaggg	120
agtggccaa						129

<210> 21

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 21

tctagtctag	acttgccac	tccctctctg	cgcg	35
------------	-----------	------------	------	----

<210> 22

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 22

aggccttaag	agcagtcgtc	caccaccttg	ttcc	34
------------	------------	------------	------	----

<210> 23

<211> 4652

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =

synthetic construct

<400> 23

tggaactctc	ccccctgtcg	cgttcgctcg	ctcgctggct	cgtttggggg	ggtggcagct	60
caaagagctg	ccagacgacg	gccctctggc	cgtcgcccc	ccaaacgagc	cagcgagcga	120
gcgaacgcga	caggggggag	agtgccacac	tctcaagcaa	gggggttttg	taagcagtga	180
tgtcataatg	atgtaatgct	tattgtcacg	cgatagttaa	tgattaacag	tcatgtgatg	240
tgttttatcc	aataggaaga	aagcgcgcgt	atgagttctc	gcgagacttc	cggggtataa	300
aagaccgagt	gaacgagccc	gccgccattc	tttgctctgg	actgctagag	gacctctgct	360
gccatggcta	ccttctatga	agtcattgtt	cgcgccccat	ttgacgtgga	ggaacatctg	420
cctggaattt	ctgacagctt	tgtggactgg	gtaactgggtc	aaatttgga	gctgcctcca	480
gagtcagatt	taaatttgac	tctggttgaa	cagcctcagt	tgacgggtggc	tgatagaatt	540
cgccgcgtgt	tcctgtacga	gtggaacaaa	ttttccaagc	aggagtccaa	attctttgtg	600
cagtttgaaa	agggatctga	atattttcat	ctgcacacgc	ttgtggagac	ctccggcatc	660
tcttccatgg	tcctcggccg	ctacgtgagt	cagattcgcg	cccagctggg	gaaagtgggtc	720
ttccagggaa	ttgaacccca	gatcaacgac	tgggtcgcca	tcaccaaggt	aaagaagggc	780
ggagccaata	aggtgggtgga	ttctgggtat	attcccgcct	acctgctgcc	gaagggtccaa	840
ccggagcttc	agtgggcgtg	gacaaacctg	gacgagtata	aattggccgc	cctgaatctg	900
gaggagcgca	aacggctcgt	cgcgcgagttt	ctggcagaat	cctcgcagcg	ctcgcaggag	960
gcggcttcgc	agcgtgagtt	ctcggctgac	ccggtcatca	aaagcaagac	ttcccagaaa	1020
tacatggcgc	tcgtcaactg	gctcgtggag	cacggcatca	cttccgagaa	gcagtggatc	1080
caggaaaatc	aggagagcta	cctctccttc	aactccaccg	gcaactctcg	gagccagatc	1140
aaggccgcgc	tcgacaacgc	gaccaaattt	atgagtctga	caaaaagcgc	ggtggactac	1200
ctcgtgggga	gctccgttcc	cgaggacatt	tcaaaaaaca	gaatctggca	aattttttgag	1260
atgaatggct	acgacccggc	ctacgcggga	tccatcctct	acggctgggtg	tcagcgctcc	1320
ttcaacaaga	ggaacaccgt	ctggctctac	ggacccgcca	cgaccggcaa	gaccaacatc	1380
gcggaggcca	tcgcccacac	tgtgcccttt	tacggctgcg	tgaactggac	caatgaaaac	1440
tttcccttta	atgactgtgt	ggacaaaatg	ctcatttggt	gggaggaggg	aaagatgacc	1500
aacaagggtg	ttgaatccgc	caaggccatc	ctgggggggct	caaagggtgcg	ggtcgatcag	1560
aaatgtaaat	cctctgttca	aattgattct	acccctgtca	ttgtaacttc	caatacaaac	1620
atgtgtgtgg	tgttgatgg	gaattccacg	acctttgaac	accagcagcc	gctggaggac	1680
cgcatgttca	aatttgaact	gactaagcgg	ctcccgcag	attttggcaa	gattactaag	1740
caggaagtca	aggacttttt	tgcttgggca	aagggtcaatc	aggtgcccgt	gactcacgag	1800
tttaaagttc	ccagggaatt	ggcgggaact	aaaggggagg	agaaatctct	aaaacgccc	1860
ctgggtgacg	tcaccaatac	tagctataaa	agtctggaga	agcgggccag	gctctcattt	1920
gttcccagga	cgctcgcag	ttcagacgtg	actgttgatc	ccgctcctct	gcgaccgctc	1980
aattggaatt	caaggatatg	ttgcaaatgt	gactatcatg	ctcaatttga	caacatttct	2040
aacaaatgtg	atgaatgtga	atatttgaat	cggggcaaaa	atggatgtat	ctgtcacaat	2100
gtaactcact	gtcaaatttg	tcatgggatt	ccccctggg	aaaaggaaaa	cttgtcagat	2160
tttggggatt	ttgacgatgc	caataaagaa	cagtaataaa	agcgagtagt	catgtctttt	2220
gttgatcacc	ctccagattg	gttgggaagaa	gttgggtgaag	gtcttcgcga	gtttttgggc	2280
cttgaagcgg	gcccaccgaa	accaaaaccc	aatcagcagc	atcaagatca	agcccgtggg	2340
cttgtgctgc	ctgggtataa	ctatctcgga	cccggaaacg	gtctcgatcg	aggagagcct	2400
gtcaacaggg	cagacgaggt	cgcgcgagag	cacgacatct	cgtacaacga	gcagcttgag	2460
gcgggagaca	acccttacct	caagtacaac	cacgcggacg	ccgagtttca	ggagaagctc	2520
gccgacgaca	catccttcgg	gggaaacctc	ggaaaggcag	tctttcaggc	caagaaaagg	2580
gttctcgaac	cttttggcct	ggttgaagag	ggtgctaaga	cggcccctac	cggaaagcgg	2640
atagacgacc	actttccaaa	aagaaagaag	gctcggaccg	aagaggactc	caagccttcc	2700
acctcgtcag	acgccgaagc	tggacccagc	ggatcccagc	agctgcaaat	cccagcccaa	2760
ccagcctcaa	gtttgggagc	tgatacaatg	tctgcgggag	gtggcggccc	attgggcgac	2820
aataaccaag	gtgccgatgg	agtgggcaat	gcctcgggag	attggcattg	cgattccacg	2880
tggatggggg	acagagtcgt	caccaagtcc	acccgaacct	gggtgctgcc	cagctacaac	2940
aaccaccagt	accgagagat	caaaagcggc	tccgtcgacg	gaagcaacgc	caacgcctac	3000
tttgataaca	gcacccccctg	gggttacttt	gactttaacc	gcttccacag	ccactggagc	3060
ccccgagact	ggcaaagact	catcaacaac	tactggggct	tcagaccccc	gtccctcaga	3120
gtcaaaaatct	tcaacattca	agtcaaagag	gtcacgggtg	aggactccac	caccaccatc	3180
gccaacaacc	tcacctccac	cgtccaagtg	tttacggacg	acgactacca	gctgccctac	3240
gtcgtcggca	acgggaccga	gggatgcctg	ccggccttcc	ctccgcaggt	ctttacgctg	3300
ccgcagtagc	gttacgcgac	gctgaaccgc	gacaacacag	aaaatcccac	cgagaggagc	3360
agcttcttct	gcctagagta	ctttcccagc	aagatgctga	gaacgggcaa	caactttgag	3420
tttacctaca	actttgagga	ggtgcccttc	cactccagct	tcgctcccag	tcagaacctg	3480
ttcaagctgg	ccaacccgct	ggtggaccag	tacttgtacc	gcttcgtgag	cacaaataac	3540
actggcggag	tccagttcaa	caagaacctg	gccgggagat	acgccaacac	ctacaaaaac	3600

tggttcccgg	ggcccatggg	ccgaacccag	ggctggaacc	tgggctccgg	ggtcaaccgc	3660
gccagtgtca	gcgccttcgc	cacgaccaat	aggatggagc	tcgagggcgc	gagttaccag	3720
gtgccccgc	agccgaacgg	catgaccaac	aacctccagg	gcagcaacac	ctatgccctg	3780
gagaacacta	tgatcttcaa	cagccagccg	gcgaacccgg	gcaccaccgc	cacgtacctc	3840
gagggcaaca	tgctcatcac	cagcgagagc	gagacgcagc	cggtgaaccg	cgtggcgtag	3900
aacgtcggcg	ggcagatggc	caccaacaac	cagagctcca	ccactgcccc	cgcgaccggc	3960
acgtacaacc	tccaggaaat	cgtgcccggc	agcgtgtgga	tggagagggg	cgtgtacctc	4020
caaggaccca	tctgggcca	gatcccagag	acggggggcg	actttcaccc	ctctccggcc	4080
atgggaggat	tcggactcaa	acacccaccg	cccatgatgc	tcataaagaa	cacgcctgtg	4140
cccggaaata	tcaccagctt	ctcggacgtg	cccgtcagca	gcttcatac	ccagtacagc	4200
accgggcagg	tcaccgtgga	gatggagtgg	gagctcaaga	aggaaaactc	caagagggtg	4260
aacccagaga	tccagtacac	aaacaactac	aacgaccccc	agtttggtga	ctttgccccg	4320
gacagcaccg	gggaatacag	aaccaccaga	cctatcggaa	cccgatacct	tacccgaccc	4380
ctttaaccca	ttcatgtcgc	ataccctcaa	taaaccgtgt	attcgtgtca	gtaaaatact	4440
gcctcttggtg	gtcattcaat	gaataacagc	ttacaacatc	tacaaaacct	ccttgcttga	4500
gagtggtggca	ctctcccccc	tgctcgcgtt	gctcgcctcg	tggctcgttt	gggggggtgg	4560
cagctcaaag	agctgccaga	cgacggccct	ctggccgtcg	cccccccaa	cgagccagcg	4620
agcgagcgaa	cgcgacaggg	gggagagtgc	ca			4652

<210> 24

<211> 390

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 24

Met	Ala	Leu	Val	Asn	Trp	Leu	Val	Glu	His	Gly	Ile	Thr	Ser	Glu	Lys
1				5				10						15	
Gln	Trp	Ile	Gln	Glu	Asn	Gln	Glu	Ser	Tyr	Leu	Ser	Phe	Asn	Ser	Thr
		20						25					30		
Gly	Asn	Ser	Arg	Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Thr	Lys
		35					40					45			
Ile	Met	Ser	Leu	Thr	Lys	Ser	Ala	Val	Asp	Tyr	Leu	Val	Gly	Ser	Ser
	50					55					60				
Val	Pro	Glu	Asp	Ile	Ser	Lys	Asn	Arg	Ile	Trp	Gln	Ile	Phe	Glu	Met
65				70					75					80	
Asn	Gly	Tyr	Asp	Pro	Ala	Tyr	Ala	Gly	Ser	Ile	Leu	Tyr	Gly	Trp	Cys
			85					90					95		
Gln	Arg	Ser	Phe	Asn	Lys	Arg	Asn	Thr	Val	Trp	Leu	Tyr	Gly	Pro	Ala
		100						105					110		
Thr	Thr	Gly	Lys	Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Thr	Val	Pro
		115					120					125			
Phe	Tyr	Gly	Cys	Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp
	130					135					140				
Cys	Val	Asp	Lys	Met	Leu	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thr	Asn
145					150					155				160	
Lys	Val	Val	Glu	Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg
			165					170						175	
Val	Asp	Gln	Lys	Cys	Lys	Ser	Ser	Val	Gln	Ile	Asp	Ser	Thr	Pro	Val
		180						185					190		
Ile	Val	Thr	Ser	Asn	Thr	Asn	Met	Cys	Val	Val	Val	Asp	Gly	Asn	Ser
		195				200						205			
Thr	Thr	Phe	Glu	His	Gln	Gln	Pro	Leu	Glu	Asp	Arg	Met	Phe	Lys	Phe
	210				215					220					
Glu	Leu	Thr	Lys	Arg	Leu	Pro	Pro	Asp	Phe	Gly	Lys	Ile	Thr	Lys	Gln
225					230					235				240	
Glu	Val	Lys	Asp	Phe	Phe	Ala	Trp	Ala	Lys	Val	Asn	Gln	Val	Pro	Val
			245					250						255	
Thr	His	Glu	Phe	Lys	Val	Pro	Arg	Glu	Leu	Ala	Gly	Thr	Lys	Gly	Ala
		260						265					270		

Glu Lys Ser Leu Lys Arg Pro Leu Gly Asp Val Thr Asn Thr Ser Tyr
 275 280 285
 Lys Ser Leu Glu Lys Arg Ala Arg Leu Ser Phe Val Pro Glu Thr Pro
 290 295 300
 Arg Ser Ser Asp Val Thr Val Asp Pro Ala Pro Leu Arg Pro Leu Asn
 305 310 315 320
 Trp Asn Ser Arg Tyr Asp Cys Lys Cys Asp Tyr His Ala Gln Phe Asp
 325 330 335
 Asn Ile Ser Asn Lys Cys Asp Glu Cys Glu Tyr Leu Asn Arg Gly Lys
 340 345 350
 Asn Gly Cys Ile Cys His Asn Val Thr His Cys Gln Ile Cys His Gly
 355 360 365
 Ile Pro Pro Trp Glu Lys Glu Asn Leu Ser Asp Phe Gly Asp Phe Asp
 370 375 380
 Asp Ala Asn Lys Glu Gln
 385 390

<210> 25

<211> 594

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 25

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Asp Trp Val Thr Gly
 1 5 10 15
 Gln Ile Trp Glu Leu Pro Pro Glu Ser Asp Leu Asn Leu Thr Leu Val
 20 25 30
 Glu Gln Pro Gln Leu Thr Val Ala Asp Arg Ile Arg Arg Val Phe Leu
 35 40 45
 Tyr Glu Trp Asn Lys Phe Ser Lys Gln Glu Ser Lys Phe Phe Val Gln
 50 55 60
 Phe Glu Lys Gly Ser Glu Tyr Phe His Leu His Thr Leu Val Glu Thr
 65 70 75 80
 Ser Gly Ile Ser Ser Met Val Leu Gly Arg Tyr Val Ser Gln Ile Arg
 85 90 95
 Ala Gln Leu Val Lys Val Val Phe Gln Gly Ile Glu Pro Gln Ile Asn
 100 105 110
 Asp Trp Val Ala Ile Thr Lys Val Lys Lys Gly Gly Ala Asn Lys Val
 115 120 125
 Val Asp Ser Gly Tyr Ile Pro Ala Tyr Leu Leu Pro Lys Val Gln Pro
 130 135 140
 Glu Leu Gln Trp Ala Trp Thr Asn Leu Asp Glu Tyr Lys Leu Ala Ala
 145 150 155 160
 Leu Asn Leu Glu Glu Arg Lys Arg Leu Val Ala Gln Phe Leu Ala Glu
 165 170 175
 Ser Ser Gln Arg Ser Gln Glu Ala Ala Ser Gln Arg Glu Phe Ser Ala
 180 185 190
 Asp Pro Val Ile Lys Ser Lys Thr Ser Gln Lys Tyr Met Ala Leu Val
 195 200 205
 Asn Trp Leu Val Glu His Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln
 210 215 220
 Glu Asn Gln Glu Ser Tyr Leu Ser Phe Asn Ser Thr Gly Asn Ser Arg
 225 230 235 240
 Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Thr Lys Ile Met Ser Leu
 245 250 255
 Thr Lys Ser Ala Val Asp Tyr Leu Val Gly Ser Ser Val Pro Glu Asp
 260 265 270
 Ile Ser Lys Asn Arg Ile Trp Gln Ile Phe Glu Met Asn Gly Tyr Asp
 275 280 285

Pro Ala Tyr Ala Gly Ser Ile Leu Tyr Gly Trp Cys Gln Arg Ser Phe
 290 295 300
 Asn Lys Arg Asn Thr Val Trp Leu Tyr Gly Pro Ala Thr Thr Gly Lys
 305 310 315 320
 Thr Asn Ile Ala Glu Ala Ile Ala His Thr Val Pro Phe Tyr Gly Cys
 325 330 335
 Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp Cys Val Asp Lys
 340 345 350
 Met Leu Ile Trp Trp Glu Glu Gly Lys Met Thr Asn Lys Val Val Glu
 355 360 365
 Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg Val Asp Gln Lys
 370 375 380
 Cys Lys Ser Ser Val Gln Ile Asp Ser Thr Pro Val Ile Val Thr Ser
 385 390 395 400
 Asn Thr Asn Met Cys Val Val Val Asp Gly Asn Ser Thr Thr Phe Glu
 405 410 415
 His Gln Gln Pro Leu Glu Asp Arg Met Phe Lys Phe Glu Leu Thr Lys
 420 425 430
 Arg Leu Pro Pro Asp Phe Gly Lys Ile Thr Lys Gln Glu Val Lys Asp
 435 440 445
 Phe Phe Ala Trp Ala Lys Val Asn Gln Val Pro Val Thr His Glu Phe
 450 455 460
 Lys Val Pro Arg Glu Leu Ala Gly Thr Lys Gly Ala Glu Lys Ser Leu
 465 470 475 480
 Lys Arg Pro Leu Gly Asp Val Thr Asn Thr Ser Tyr Lys Ser Leu Glu
 485 490 495
 Lys Arg Ala Arg Leu Ser Phe Val Pro Glu Thr Pro Arg Ser Ser Asp
 500 505 510
 Val Thr Val Asp Pro Ala Pro Leu Arg Pro Leu Asn Trp Asn Ser Arg
 515 520 525
 Tyr Asp Cys Lys Cys Asp Tyr His Ala Gln Phe Asp Asn Ile Ser Asn
 530 535 540
 Lys Cys Asp Glu Cys Glu Tyr Leu Asn Arg Gly Lys Asn Gly Cys Ile
 545 550 555 560
 Cys His Asn Val Thr His Cys Gln Ile Cys His Gly Ile Pro Pro Trp
 565 570 575
 Glu Lys Glu Asn Leu Ser Asp Phe Gly Asp Phe Asp Asp Ala Asn Lys
 580 585 590
 Glu Gln

<210> 26
 <211> 724
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 26
 Met Ser Phe Val Asp His Pro Pro Asp Trp Leu Glu Glu Val Gly Glu
 1 5 10 15
 Gly Leu Arg Glu Phe Leu Gly Leu Glu Ala Gly Pro Pro Lys Pro Lys
 20 25 30
 Pro Asn Gln Gln His Gln Asp Gln Ala Arg Gly Leu Val Leu Pro Gly
 35 40 45
 Tyr Asn Tyr Leu Gly Pro Gly Asn Gly Leu Asp Arg Gly Glu Pro Val
 50 55 60
 Asn Arg Ala Asp Glu Val Ala Arg Glu His Asp Ile Ser Tyr Asn Glu
 65 70 75 80
 Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala Asp
 85 90 95

Ala	Glu	Phe	Gln	Glu	Lys	Leu	Ala	Asp	Asp	Thr	Ser	Phe	Gly	Gly	Asn
			100					105					110		
Leu	Gly	Lys	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Val	Leu	Glu	Pro	Phe
		115					120					125			
Gly	Leu	Val	Glu	Glu	Gly	Ala	Lys	Thr	Ala	Pro	Thr	Gly	Lys	Arg	Ile
	130					135					140				
Asp	Asp	His	Phe	Pro	Lys	Arg	Lys	Lys	Ala	Arg	Thr	Glu	Glu	Asp	Ser
145					150					155					160
Lys	Pro	Ser	Thr	Ser	Ser	Asp	Ala	Glu	Ala	Gly	Pro	Ser	Gly	Ser	Gln
				165					170					175	
Gln	Leu	Gln	Ile	Pro	Ala	Gln	Pro	Ala	Ser	Ser	Leu	Gly	Ala	Asp	Thr
			180					185					190		
Met	Ser	Ala	Gly	Gly	Gly	Gly	Pro	Leu	Gly	Asp	Asn	Asn	Gln	Gly	Ala
		195					200					205			
Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly	Asp	Trp	His	Cys	Asp	Ser	Thr	Trp
	210					215					220				
Met	Gly	Asp	Arg	Val	Val	Thr	Lys	Ser	Thr	Arg	Thr	Trp	Val	Leu	Pro
225					230					235					240
Ser	Tyr	Asn	Asn	His	Gln	Tyr	Arg	Glu	Ile	Lys	Ser	Gly	Ser	Val	Asp
				245					250					255	
Gly	Ser	Asn	Ala	Asn	Ala	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr
			260					265					270		
Phe	Asp	Phe	Asn	Arg	Phe	His	Ser	His	Trp	Ser	Pro	Arg	Asp	Trp	Gln
		275					280					285			
Arg	Leu	Ile	Asn	Asn	Tyr	Trp	Gly	Phe	Arg	Pro	Arg	Ser	Leu	Arg	Val
	290					295					300				
Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Val	Gln	Asp	Ser	Thr
305					310					315					320
Thr	Thr	Ile	Ala	Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr	Asp	
				325				330					335		
Asp	Asp	Tyr	Gln	Leu	Pro	Tyr	Val	Val	Gly	Asn	Gly	Thr	Glu	Gly	Cys
			340					345					350		
Leu	Pro	Ala	Phe	Pro	Pro	Gln	Val	Phe	Thr	Leu	Pro	Gln	Tyr	Gly	Tyr
		355				360						365			
Ala	Thr	Leu	Asn	Arg	Asp	Asn	Thr	Glu	Asn	Pro	Thr	Glu	Arg	Ser	Ser
	370					375					380				
Phe	Phe	Cys	Leu	Glu	Tyr	Phe	Pro	Ser	Lys	Met	Leu	Arg	Thr	Gly	Asn
385					390					395					400
Asn	Phe	Glu	Phe	Thr	Tyr	Asn	Phe	Glu	Glu	Val	Pro	Phe	His	Ser	Ser
				405				410					415		
Phe	Ala	Pro	Ser	Gln	Asn	Leu	Phe	Lys	Leu	Ala	Asn	Pro	Leu	Val	Asp
			420					425					430		
Gln	Tyr	Leu	Tyr	Arg	Phe	Val	Ser	Thr	Asn	Asn	Thr	Gly	Gly	Val	Gln
		435				440						445			
Phe	Asn	Lys	Asn	Leu	Ala	Gly	Arg	Tyr	Ala	Asn	Thr	Tyr	Lys	Asn	Trp
	450					455					460				
Phe	Pro	Gly	Pro	Met	Gly	Arg	Thr	Gln	Gly	Trp	Asn	Leu	Gly	Ser	Gly
465					470				475						480
Val	Asn	Arg	Ala	Ser	Val	Ser	Ala	Phe	Ala	Thr	Thr	Asn	Arg	Met	Glu
				485				490					495		
Leu	Glu	Gly	Ala	Ser	Tyr	Gln	Val	Pro	Pro	Gln	Pro	Asn	Gly	Met	Thr
			500					505					510		
Asn	Asn	Leu	Gln	Gly	Ser	Asn	Thr	Tyr	Ala	Leu	Glu	Asn	Thr	Met	Ile
		515					520					525			
Phe	Asn	Ser	Gln	Pro	Ala	Asn	Pro	Gly	Thr	Thr	Ala	Thr	Tyr	Leu	Glu
	530					535					540				
Gly	Asn	Met	Leu	Ile	Thr	Ser	Glu	Ser	Glu	Thr	Gln	Pro	Val	Asn	Arg
545					550					555					560
Val	Ala	Tyr	Asn	Val	Gly	Gly	Gln	Met	Ala	Thr	Asn	Asn	Gln	Ser	Ser
				565					570					575	
Thr	Thr	Ala	Pro	Ala	Thr	Gly	Thr	Tyr	Asn	Leu	Gln	Glu	Ile	Val	Pro
			580					585					590		
Gly	Ser	Val	Trp	Met	Glu	Arg	Asp	Val	Tyr	Leu	Gln	Gly	Pro	Ile	Trp

595 600 605
 Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala Met
 610 615 620
 Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys Asn
 625 630 635 640
 Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val Ser
 645 650 655
 Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met Glu
 660 665 670
 Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln
 675 680 685
 Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro Asp
 690 695 700
 Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr Leu
 705 710 715 720
 Thr Arg Pro Leu

<210> 27

<211> 588

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 27

Thr Ala Pro Thr Gly Lys Arg Ile Asp Asp His Phe Pro Lys Arg Lys
 1 5 10 15
 Lys Ala Arg Thr Glu Glu Asp Ser Lys Pro Ser Thr Ser Ser Asp Ala
 20 25 30
 Glu Ala Gly Pro Ser Gly Ser Gln Gln Leu Gln Ile Pro Ala Gln Pro
 35 40 45
 Ala Ser Ser Leu Gly Ala Asp Thr Met Ser Ala Gly Gly Gly Pro
 50 55 60
 Leu Gly Asp Asn Asn Gln Gly Ala Asp Gly Val Gly Asn Ala Ser Gly
 65 70 75 80
 Asp Trp His Cys Asp Ser Thr Trp Met Gly Asp Arg Val Val Thr Lys
 85 90 95
 Ser Thr Arg Thr Trp Val Leu Pro Ser Tyr Asn Asn His Gln Tyr Arg
 100 105 110
 Glu Ile Lys Ser Gly Ser Val Asp Gly Ser Asn Ala Asn Ala Tyr Phe
 115 120 125
 Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe His Ser
 130 135 140
 His Trp Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Tyr Trp Gly
 145 150 155 160
 Phe Arg Pro Arg Ser Leu Arg Val Lys Ile Phe Asn Ile Gln Val Lys
 165 170 175
 Glu Val Thr Val Gln Asp Ser Thr Thr Thr Ile Ala Asn Asn Leu Thr
 180 185 190
 Ser Thr Val Gln Val Phe Thr Asp Asp Asp Tyr Gln Leu Pro Tyr Val
 195 200 205
 Val Gly Asn Gly Thr Glu Gly Cys Leu Pro Ala Phe Pro Pro Gln Val
 210 215 220
 Phe Thr Leu Pro Gln Tyr Gly Tyr Ala Thr Leu Asn Arg Asp Asn Thr
 225 230 235 240
 Glu Asn Pro Thr Glu Arg Ser Ser Phe Phe Cys Leu Glu Tyr Phe Pro
 245 250 255
 Ser Lys Met Leu Arg Thr Gly Asn Asn Phe Glu Phe Thr Tyr Asn Phe
 260 265 270
 Glu Glu Val Pro Phe His Ser Ser Phe Ala Pro Ser Gln Asn Leu Phe

275 280 285
 Lys Leu Ala Asn Pro Leu Val Asp Gln Tyr Leu Tyr Arg Phe Val Ser
 290 295 300
 Thr Asn Asn Thr Gly Gly Val Gln Phe Asn Lys Asn Leu Ala Gly Arg
 305 310 315 320
 Tyr Ala Asn Thr Tyr Lys Asn Trp Phe Pro Gly Pro Met Gly Arg Thr
 325 330 335
 Gln Gly Trp Asn Leu Gly Ser Gly Val Asn Arg Ala Ser Val Ser Ala
 340 345 350
 Phe Ala Thr Thr Asn Arg Met Glu Leu Glu Gly Ala Ser Tyr Gln Val
 355 360 365
 Pro Pro Gln Pro Asn Gly Met Thr Asn Asn Leu Gln Gly Ser Asn Thr
 370 375 380
 Tyr Ala Leu Glu Asn Thr Met Ile Phe Asn Ser Gln Pro Ala Asn Pro
 385 390 395 400
 Gly Thr Thr Ala Thr Tyr Leu Glu Gly Asn Met Leu Ile Thr Ser Glu
 405 410 415
 Ser Glu Thr Gln Pro Val Asn Arg Val Ala Tyr Asn Val Gly Gly Gln
 420 425 430
 Met Ala Thr Asn Asn Gln Ser Ser Thr Thr Ala Pro Ala Thr Gly Thr
 435 440 445
 Tyr Asn Leu Gln Glu Ile Val Pro Gly Ser Val Trp Met Glu Arg Asp
 450 455 460
 Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro Glu Thr Gly Ala
 465 470 475 480
 His Phe His Pro Ser Pro Ala Met Gly Gly Phe Gly Leu Lys His Pro
 485 490 495
 Pro Pro Met Met Leu Ile Lys Asn Thr Pro Val Pro Gly Asn Ile Thr
 500 505 510
 Ser Phe Ser Asp Val Pro Val Ser Ser Phe Ile Thr Gln Tyr Ser Thr
 515 520 525
 Gly Gln Val Thr Val Glu Met Glu Trp Glu Leu Lys Lys Glu Asn Ser
 530 535 540
 Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Asn Asn Tyr Asn Asp Pro
 545 550 555 560
 Gln Phe Val Asp Phe Ala Pro Asp Ser Thr Gly Glu Tyr Arg Thr Thr
 565 570 575
 Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
 580 585

<210> 28
 <211> 532
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 28
 Met Ser Ala Gly Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly Ala
 1 5 10 15
 Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr Trp
 20 25 30
 Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val Leu Pro
 35 40 45
 Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser Val Asp
 50 55 60
 Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr
 65 70 75 80
 Phe Asp Phe Asn Arg Phe His Ser His Trp Ser Pro Arg Asp Trp Gln
 85 90 95
 Arg Leu Ile Asn Asn Tyr Trp Gly Phe Arg Pro Arg Ser Leu Arg Val

Lys	Ile	Phe	100	Asn	Ile	Gln	Val	Lys	105	Glu	Val	Thr	Val	Gln	110	Asp	Ser	Thr
Thr	Thr	115	Ile	Ala	Asn	Asn	Leu	Thr	120	Ser	Thr	Val	Gln	125	Val	Phe	Thr	Asp
Asp	Asp	130	Tyr	Gln	Leu	Pro	Tyr	Val	135	Val	Gly	Asn	Gly	140	Thr	Glu	Gly	Cys
145	Leu	Pro	Ala	Phe	Pro	Pro	Gln	Val	150	Phe	Thr	Leu	Pro	155	Gln	Tyr	Gly	160
Ala	Thr	Leu	Asn	Arg	Asp	Asn	Thr	Glu	165	Asn	Pro	Thr	Glu	170	Arg	Ser	Ser	175
Phe	Phe	Cys	180	Leu	Glu	Tyr	Phe	Pro	185	Ser	Lys	Met	Leu	Arg	Thr	Gly	Asn	190
Asn	Phe	Glu	195	Phe	Thr	Tyr	Asn	Phe	200	Glu	Glu	Val	Pro	205	Phe	His	Ser	Ser
210	Phe	Ala	Pro	Ser	Gln	Asn	Leu	Phe	215	Lys	Leu	Ala	Asn	220	Pro	Leu	Val	Asp
225	Gln	Tyr	Leu	Tyr	Arg	Phe	Val	Ser	230	Thr	Asn	Asn	Thr	235	Gly	Gly	Val	Gln
Phe	Asn	Lys	Asn	Leu	Ala	Gly	Arg	Tyr	245	Ala	Asn	Thr	Tyr	250	Lys	Asn	Trp	255
Phe	Pro	Gly	260	Pro	Met	Gly	Arg	Thr	265	Gln	Gly	Trp	Asn	270	Leu	Gly	Ser	Gly
Val	Asn	Arg	Ala	Ser	Val	Ser	280	Ala	285	Phe	Ala	Thr	Thr	290	Asn	Arg	Met	Glu
Leu	Glu	Gly	Ala	Ser	Tyr	Gln	Val	Pro	295	Pro	Pro	Gln	Pro	300	Asn	Gly	Met	Thr
305	Asn	Asn	Leu	Gln	Gly	Ser	Asn	Thr	310	Tyr	Ala	Leu	Glu	315	Asn	Thr	Met	Ile
Phe	Asn	Ser	Gln	Pro	Ala	Asn	Pro	Gly	325	Thr	Thr	Ala	Thr	330	Tyr	Leu	Glu	335
Gly	Asn	Met	340	Leu	Ile	Thr	Ser	Glu	345	Ser	Glu	Thr	Gln	350	Pro	Val	Asn	Arg
Val	Ala	Tyr	355	Asn	Val	Gly	Gly	Gln	360	Met	Ala	Thr	Asn	365	Asn	Gln	Ser	Ser
Thr	Thr	Ala	370	Pro	Ala	Thr	Gly	Thr	375	Tyr	Asn	Leu	Gln	380	Glu	Ile	Val	Pro
385	Gly	Ser	Val	Trp	Met	Glu	Arg	Asp	390	Val	Tyr	Leu	Gln	395	Gly	Pro	Ile	Trp
Ala	Lys	Ile	405	Pro	Glu	Thr	Gly	Ala	410	His	Phe	His	Pro	415	Ser	Pro	Ala	Met
Gly	Gly	Phe	420	Gly	Leu	Lys	His	Pro	425	Pro	Pro	Met	Met	430	Leu	Ile	Lys	Asn
Thr	Pro	Val	435	Pro	Gly	Asn	Ile	Thr	440	Ser	Phe	Ser	Asp	445	Val	Pro	Val	Ser
Ser	Phe	Ile	450	Thr	Gln	Tyr	Ser	Thr	455	Gly	Gln	Val	Thr	460	Val	Glu	Met	Glu
465	Trp	Glu	Leu	Lys	Lys	Glu	Asn	Ser	470	Lys	Arg	Trp	Asn	475	Pro	Glu	Ile	Gln
Tyr	Thr	Asn	485	Asn	Tyr	Asn	Asp	Pro	490	Gln	Phe	Val	Asp	495	Phe	Ala	Pro	Asp
Ser	Thr	Gly	500	Glu	Tyr	Arg	Thr	Thr	505	Arg	Pro	Ile	Gly	510	Thr	Arg	Tyr	Leu
Thr	Arg	Pro	515	Leu					520					525				
530																		

<210> 29

<211> 2307

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =

synthetic construct

<400> 29
aggctctcat ttgttcccga gacgcctcgc agttcagacg tgactgttga tcccgtcct 60
ctgcgaccgc tcaattggaa ttcaagtaaa taaagcgagt agtcatgtct tttgttgatc 120
accctccaga ttggttggaa gaagttggtg aaggtcttcg cgagtttttg ggccttgaag 180
cgggccacc gaaacaaaa cccaatcagc agcatcaaga tcaagcccgt ggtcttgtgc 240
tgcctggtta taactatctc ggacccggaa acggtctcga tcgaggagag cctgtcaaca 300
gggcagacga ggtcgcgcga gagcacgaca tctcgtacaa cgagcagctt gaggcgggag 360
acaacccta cctcaagtac aaccacgcgg acgccgagtt tcaggagaag ctgcccagc 420
acacatcctt cgggggaaac ctcggaaagg cagtctttca ggccaagaaa agggttctcg 480
aaccttttgg cctggttgaa gaggttgcta agacggcccc taccggaaag cggatagacg 540
accactttcc aaaaagaaag aaggctcgga ccgaagagga ctccaagcct tccacctcgt 600
cagacgccga agctggaccc agcggatccc agcagctgca aatcccagcc caaccagcct 660
caagtttggg agctgataca atgtctgcgg gaggtggcgg cccattgggg gacaataacc 720
aaggtgccga tggagtgggc aatgcctcgg gagattggca ttgcgattcc acgtggatgg 780
gggacagagt cgtcaccaag tccacccgaa cctgggtgct gccagctac aacaaccacc 840
agtaccgaga gatcaaaagc ggctccgtcg acggaagcaa cgccaacgcc tactttggat 900
acagcacccc ctgggggtac tttgacttta accgcttcca cagccactgg agccccgag 960
actggcaaag actcatcaac aactactggg gcttcagacc ccggtccctc agagtcaaaa 1020
tcttcaacat tcaagtcaaa gaggtcacgg tgcaggactc caccaccacc atcgccaaca 1080
acctcacctc caccgtccaa gtgtttacgg acgacgacta ccagctgccc tacgtcgtcg 1140
gcaacgggac cgagggatgc ctgccggcct tccctccgca ggtctttacg ctgccgcagt 1200
acggttacgc gacgctgaac cgcgacaaca cagaaaatcc caccgagagg agcagcttct 1260
tctgcctaga gtactttccc agcaagatgc tgagaacggg caacaacttt gagtttacct 1320
acaactttga ggaggtgccc ttccactcca gcttcgctcc cagtcagaac ctgttcaagc 1380
tggccaaccc gctggtggac cagtacttgt accgcttcgt gagcacaat aacactggcg 1440
gagtccagtt caacaagaac ctggccggga gatacgccaa cacctacaaa aactggttcc 1500
cggggcccat gggccgaacc cagggttgga acctgggctc cggggtcaac cgcgccagt 1560
tcagcgcctt cgccacgacc aataggatgg agctcgagg cgcgagttac caggtgcccc 1620
cgcagccgaa cggcatgacc aacaacctcc agggcagcaa cacctatgcc ctggagaaca 1680
ctatgatctt caacagccag ccggcgaacc cgggcaccac cgccacgtac ctgagggca 1740
acatgctcat caccagcgag agcgagacgc agccggtgaa ccgctggtcg tacaacgtcg 1800
gcgggcagat ggccaccaac aaccagagct ccaccactgc ccccgcgacc ggcacgtaca 1860
acctccagga aatcgtgccc ggcagcgtgt ggatggagag ggacgtgtac ctccaaggac 1920
ccatctgggc caagatccca gagacggggg cgcactttca cccctctccg gccatgggcg 1980
gattcggact caaacaccca ccgcccata tgctcatcaa gaacacgcct gtgcccggaa 2040
atatcaccag cttctcggac gtgcccgtca gcagcttcat caccagtag agcaccgggc 2100
aggtcaccgt ggagatggag tgggagctca agaaggaaaa ctccaagagg tgaacccag 2160
agatccagta cacaacaac tacaacgacc cccagtttgt ggactttgcc ccggacagca 2220
ccggggaata cagaaccacc agacctatcg gaacccgata ccttaccgga cccctttaac 2280
ccattcatgt cgcataccct caataaa 2307

<210> 30
<211> 2264
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 30
aggctctcat ttgttcccga gacgcctcgc agttcagacg tgactgttga tcccgtcct 60
ctgcgaccgc tcaattggaa ttcaagattg gttggaagaa gttggtgaag gtcttcgcga 120
gtttttgggc cttgaagcgg gccaccgaa accaaaaccc aatcagcagc atcaagatca 180
agcccgtggt cttgtgctgc ctggttataa ctatctcggg cccggaaacg gtctcgatcg 240
aggagagcct gtcaacaggg cagacgaggt cgcgcgagag cacgacatct cgtacaacga 300
gcagcttgag gcgggagaca acccctacct caagtacaac cacgcggacg ccgagtttca 360
ggagaagctc gccgacgaca catccttcgg gggaaacctc ggaaaggcag tctttcaggc 420
caagaaaagg gttctcgaac cttttggcct ggttgaagag ggtgctaaga cggcccctac 480
cggaaagcgg atagacgacc actttccaaa aagaaagaag gctcggaccg aagaggactc 540
caagccttcc acctcgtcag acgccgaagc tggacccagc ggatcccagc agctgcaaat 600

cccagcccaa	ccagcctcaa	gtttgggagc	tgatacaatg	tctgcgggag	gtggcggccc	660
attgggcgac	aataaccaag	gtgccgatgg	agtgggcaat	gcctcgggag	attggcattg	720
cgattccacg	tggatggggg	acagagtcgt	caccaagtcc	acccgaacct	gggtgctgcc	780
cagctacaac	aaccaccagt	accgagagat	caaaagcggc	tccgtcgacg	gaagcaacgc	840
caacgcctac	tttggataca	gcacccccctg	gggggtacttt	gactttaacc	gcttccacag	900
ccactggagc	ccccgagact	ggcaaagact	catcaacaac	tactggggct	tcagacccccg	960
gtccctcaga	gtcaaaatct	tcaacattca	agtcaaagag	gtcacgggtgc	aggactccac	1020
caccaccatc	gccaacaacc	tcacctccac	cgtccaagtg	tttacggacg	acgactacca	1080
gctgccctac	gtcgtcggca	acgggaccga	gggatgcctg	ccggccttcc	ctccgcaggt	1140
ctttacgctg	ccgcagtacg	gttacgcgac	gctgaaccgc	gacaacacag	aaaatcccac	1200
cgagaggagc	agcttcttct	gcctagagta	ctttcccagc	aagatgctga	gaacgggcaa	1260
caactttgag	tttacctaca	actttgagga	gggtgcccttc	cactccagct	tcgctcccag	1320
tcagaacctg	ttcaagctgg	ccaacccgct	gggtggaccag	tacttggtacc	gcttcgtgag	1380
cacaaataac	actggcggag	tccagttcaa	caagaacctg	gccggggagat	acgccaacac	1440
ctacaaaaac	tggttcccgg	ggcccatggg	ccgaaccacg	ggctggaacc	tgggctccgg	1500
gggtcaaccgc	gccagtgtca	gcgcccttcgc	cacgaccaat	aggatggagc	tcgagggcgc	1560
gagttaccag	gtgccccgc	agccgaacgg	catgaccaac	aacctccagg	gcagcaacac	1620
ctatgccctg	gagaacacta	tgatcttcaa	cagccagccg	gcgaaccggg	gcaccaccgc	1680
cacgtacctc	gagggcaaca	tgctcatcac	cagcgagagc	gagacgcagc	cgggtgaaccg	1740
cgtggcgtac	aacgtcggcg	ggcagatggc	caccaacaac	cagagctcca	ccactgcccc	1800
cgcgaccggc	acgtacaacc	tccaggaaat	cgtgcccggc	agcgtgtgga	tggagagggga	1860
cgtgtacctc	caaggaccca	tctggggcaa	gatcccagag	acgggggcgc	actttcaccc	1920
ctctccggcc	atgggcggat	tcggactcaa	acaccaccg	cccatgatgc	tcatacaaga	1980
cacgcctgtg	cccggaaata	tcaccagctt	ctcggacgtg	cccgctagca	gcttcatcac	2040
ccagtacagc	accgggcagg	tcaccgtgga	gatggagtgg	gagctcaaga	aggaaaactc	2100
caagagggtg	aaccagaga	tccagtacac	aaacaactac	aacgaccccc	agtttgtgga	2160
ctttgccccg	gacagcaccg	gggaatacag	aaccaccaga	cctatcggaa	cccgatacct	2220
tacccgaccc	ctttaaccca	ttcatgtcgc	ataccctcaa	taaa		2264

<210> 31

<211> 2264

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 31

aggctctcat	ttgttcccga	gacgcctcgc	agttcagacg	tgactgttga	tcccgtctct	60
ctgcgaccgc	tcaattggaa	ttcaagattg	gttggaaaga	gttgggtgaag	gtcttcgcga	120
gtttttgggc	cttgaagcgg	gcccaccgaa	acaaaaaccc	aatcagcagc	atcaagatca	180
agccccgtgt	cttgtgctgc	ctgggtataa	ctatctcgga	cccggaaacg	gtctcgatcg	240
aggagagcct	gtcaacaggg	cagacgaggt	cgcgcgagag	cacgacatct	cgtacaacga	300
gcagcttgag	gcgggagaca	acccctacct	caagtacaac	cacgcggacg	ccgagtttca	360
ggagaagctc	gccgacgaca	catccttcgg	gggaaacctc	ggaaaggcag	tctttcaggc	420
caagaaaagg	gttctcgaac	cttttggcct	ggttgaagag	ggtgctaaga	cggccccctac	480
cggaaagcgg	atagacgacc	actttccaaa	aagaaagaag	gctcggaccg	aagaggactc	540
caagccttcc	acctcgtcag	acgccgaagc	tggaccacag	ggatcccagc	agctgcaaat	600
cccagcccaa	ccagcctcaa	gtttgggagc	tgatacaatg	tctgcgggag	gtggcggccc	660
attgggcgac	aataaccaag	gtgccgatgg	agtgggcaat	gcctcgggag	attggcattg	720
cgattccacg	tggatggggg	acagagtcgt	caccaagtcc	acccgaacct	gggtgctgcc	780
cagctacaac	aaccaccagt	accgagagat	caaaagcggc	tccgtcgacg	gaagcaacgc	840
caacgcctac	tttggataca	gcacccccctg	gggggtacttt	gactttaacc	gcttccacag	900
ccactggagc	ccccgagact	ggcaaagact	catcaacaac	tactggggct	tcagacccccg	960
gtccctcaga	gtcaaaatct	tcaacattca	agtcaaagag	gtcacgggtgc	aggactccac	1020
caccaccatc	gccaacaacc	tcacctccac	cgtccaagtg	tttacggacg	acgactacca	1080
gctgccctac	gtcgtcggca	acgggaccga	gggatgcctg	ccggccttcc	ctccgcaggt	1140
ctttacgctg	ccgcagtacg	gttacgcgac	gctgaaccgc	gacaacacag	aaaatcccac	1200
cgagaggagc	agcttcttct	gcctagagta	ctttcccagc	aagatgctga	gaacgggcaa	1260
caactttgag	tttacctaca	actttgagga	gggtgcccttc	cactccagct	tcgctcccag	1320
tcagaacctg	ttcaagctgg	ccaacccgct	gggtggaccag	tacttggtacc	gcttcgtgag	1380
cacaaataac	actggcggag	tccagttcaa	caagaacctg	qccgggagat	acgccaacac	1440

ctacaaaaac	tggttcccgg	ggcccatggg	ccgaacccag	ggctggaacc	tgggctccgg	1500
ggtcaaccgc	gccagtgtca	gcgccttcgc	cacgaccaat	aggatggagc	tcgagggcgc	1560
gagttaccag	gtgccccgc	agccgaacgg	catgaccaac	aacctccagg	gcagcaacac	1620
ctatgccctg	gagaacacta	tgatcttcaa	cagccagccg	gcgaacccgg	gcaccaccgc	1680
cacgtacctc	gagggcaaca	tgctcatcac	cagcgagagc	gagacgcagc	cggtgaaccg	1740
cgtggcgtac	aacgtcggcg	ggcagatggc	caccaacaac	cagagctcca	ccactgcccc	1800
cgcgaccggc	acgtacaacc	tccaggaaat	cgtgcccggc	agcgtgtgga	tggagagggg	1860
cgtgtacctc	caaggaccca	tctggggcaa	gatcccagag	acggggggcg	actttcaccc	1920
ctctccggcc	atgggcggat	tcggactcaa	acaccaccg	cccatgatgc	tcatacaaga	1980
cacgcctgtg	cccggaaata	tcaccagctt	ctcggacgtg	cccgtcagca	gcttcatcac	2040
ccagtacagc	accgggcagg	tcaccgtgga	gatggagtgg	gagctcaaga	aggaaaactc	2100
caagaggtgg	aaccagaga	tccagtacac	aaacaactac	aacgaccccc	agtttgtgga	2160
ctttgccccg	gacagcaccg	gggaatacag	aaccaccaga	cctatcgga	cccgatacct	2220
tacccgaccc	ctttaaccca	ttcatgtcgc	ataccctcaa	taaa		2264

<210> 32

<211> 1292

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 32

agcgcaaacg	gctcgtcgcg	cagtttcttg	cagaatcctc	gcagcgctcg	caggaggcgg	60
cttcgcagcg	tgagttctcg	gctgaccggg	tcatacaaaag	caagacttcc	cagaaataca	120
tggcgctcgt	caactggctc	gtggagcacg	gcatcacttc	cgagaagcag	tggatccagg	180
aaaatcagga	gagctacctc	tccttcaact	ccaccggcaa	ctctcggagc	cagatcaagg	240
ccgcgctcga	caacgcgacc	aaaattatga	gtctgacaaa	aagcgcggtg	gactacctcg	300
tggggagctc	cgttcccag	gacatttcaa	aaaacagaat	ctggcaaatt	tttgagatga	360
atggctacga	cccggcctac	gcgggatcca	tcctctacgg	ctggtgtcag	cgctccttca	420
acaagaggaa	caccgtcttg	ctctacggac	ccgccacgac	cggcaagacc	aacatcgcg	480
aggccatcgc	ccacactgtg	cccttttacg	gctgcgtgaa	ctggaccaat	gaaaactttc	540
cctttaatga	ctgtgtggac	aaaatgctca	tttggtggga	ggagggaaag	atgaccaaca	600
aggtggttga	atccgccaaag	gccatcctgg	ggggctcaaa	ggtgcgggtc	gatcagaaat	660
gtaaatcctc	tgttcaaatt	gattctaccc	ctgtcattgt	aacttccaat	acaaacatgt	720
gtgtggtggt	ggatgggaat	tccacgacct	ttgaacacca	gcagccgctg	gaggaccgca	780
tgttcaaatt	tgaactgact	aagcggctcc	cgccagattt	tggcaagatt	actaagcagg	840
aagtcaagga	cttttttgct	tgggcaaagg	tcaatcagg	gccggtgact	cacgagttta	900
aagttcccag	ggaattggcg	ggaactaaag	gggcggagaa	atctctaaaa	cgcccactgg	960
gtgacgtcac	caatactagc	tataaaagtc	tggagaagcg	ggccaggctc	tcatttggtc	1020
ccgagacgcc	tcgcagttca	gacgtgactg	ttgatcccgc	tcctctgcga	ccgctcaatt	1080
ggaattcaag	gtatgattgc	aatgtgact	atcatgctca	atttgacaac	atttctaaca	1140
aatgtgatga	atgtgaatat	ttgaatcggg	gcaaaaatgg	atgtatctgt	cacaatgtaa	1200
ctcactgtca	aatttgtcat	gggattcccc	cctgggaaaa	ggaaaacttg	tcagattttg	1260
gggattttga	cgatgccaat	aaagaacagt	aa			1292

<210> 33

<211> 1870

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 33

attctttgct	ctggactgct	agaggaccct	cgctgccatg	gctaccttct	atgaagtcac	60
tgttcgcgtc	ccatttgacg	tggaggaaca	tctgcctgga	atttctgaca	gctttgtgga	120
ctgggtaact	ggtcaaattt	gggagctgcc	tccagagtca	gatttaaatt	tgactctggt	180
tgaacagcct	cagttgacgg	tggctgatag	aattcgccgc	gtgttcctgt	acgagtggaa	240
caaattttcc	aagcaggagt	ccaaattctt	tgtgcagttt	gaaaagggat	ctgaatattt	300

tcattctgcac	acgcttgtgg	agacctccgg	catctcttcc	atggtcctcg	gccgctacgt	360
gagtcagatt	cgcgcccagc	tggtgaaagt	ggtcttccag	ggaattgaac	cccagatcaa	420
cgactgggtc	gccatcacca	aggtaaagaa	gggcggagcc	aataagggtg	tggtattctgg	480
gtatatctcc	gcctacctgc	tgccgaaggt	ccaaccggag	cttcagtggg	cgtggacaaa	540
cctggacgag	tataaattgg	ccgccctgaa	tctggaggag	cgcaaacggc	tcgtcgcgca	600
gtttctggca	gaatcctcgc	agcgctcgc	ggaggcggct	tcgcagcgtg	agttctcggc	660
tgacccggtc	atcaaaagca	agacttccca	gaaatacatg	gcgctcgtca	actggctcgt	720
ggagcacggc	atcacttccg	agaagcagtg	gatccaggaa	aatcaggaga	gctacctctc	780
cttcaactcc	accggcaact	ctcggagcca	gatcaaggcc	gcgctcgaca	acgcgaccaa	840
aattatgagt	ctgacaaaaa	gcgcggtgga	ctacctcgtg	gggagctccg	ttcccagagga	900
catttcaaaa	aacagaatct	ggcaaatttt	tgagatgaat	ggctacgacc	cggcctacgc	960
gggatccatc	ctctacggct	ggtgtcagcg	ctccttcaac	aagaggaaca	ccgtctggct	1020
ctacggaccc	gccacgaccg	gcaagaccaa	catcgcgag	gccatcgccc	acactgtgcc	1080
cttttacggc	tgcgtgaact	ggaccaatga	aaactttccc	tttaatgact	gtgtggacaa	1140
aatgctcatt	tggtgggagg	agggaaagat	gaccaacaag	gtggttgaat	ccgccaaggc	1200
catcctgggg	ggctcaaagg	tgcggtcgca	tcagaaatgt	aaatcctctg	ttcaaattga	1260
ttctaccctt	gtcattgtaa	cttccaatac	aaacatgtgt	gtggtggtgg	atgggaattc	1320
cacgaccttt	gaacaccagc	agccgctgga	ggaccgcatg	ttcaaatttg	aactgactaa	1380
gcggctcccc	ccagattttg	gcaagattac	taagcaggaa	gtcaaggact	tttttgcttg	1440
ggcaaaggtc	aatcagggtgc	cggtgactca	cgagttttaa	gttcccaggg	aattggcggg	1500
aactaaaggg	gcggagaaat	ctctaaaacg	cccactgggt	gacgtcacca	atactagcta	1560
taaaagtctg	gagaagcggg	ccaggctctc	atttgttccc	gagacgcctc	gcagttcaga	1620
cgtgactgtt	gatcccgtct	ctctgcgacc	gctcaattgg	aattcaaggt	atgattgcaa	1680
atgtgactat	catgctcaat	ttgacaacat	ttctaacaaa	tgtgatgaat	gtgaatatct	1740
gaatcggggc	aaaaatggat	gtatctgtca	caatgtaact	cactgtcaaa	tttgtcatgg	1800
gattcccccc	tgggaaaagg	aaaacttgtc	agattttggg	gattttgacg	atgccaataa	1860
agaacagtaa						1870

<210> 34

<211> 330

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 34

Met	Ala	Leu	Val	Asn	Trp	Leu	Val	Glu	His	Gly	Ile	Thr	Ser	Glu	Lys
1				5				10						15	
Gln	Trp	Ile	Gln	Glu	Asn	Gln	Glu	Ser	Tyr	Leu	Ser	Phe	Asn	Ser	Thr
		20						25					30		
Gly	Asn	Ser	Arg	Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Thr	Lys
		35					40					45			
Ile	Met	Ser	Leu	Thr	Lys	Ser	Ala	Val	Asp	Tyr	Leu	Val	Gly	Ser	Ser
	50				55				60						
Val	Pro	Glu	Asp	Ile	Ser	Lys	Asn	Arg	Ile	Trp	Gln	Ile	Phe	Glu	Met
65				70				75					80		
Asn	Gly	Tyr	Asp	Pro	Ala	Tyr	Ala	Gly	Ser	Ile	Leu	Tyr	Gly	Trp	Cys
			85					90					95		
Gln	Arg	Ser	Phe	Asn	Lys	Arg	Asn	Thr	Val	Trp	Leu	Tyr	Gly	Pro	Ala
		100						105					110		
Thr	Thr	Gly	Lys	Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Thr	Val	Pro
		115					120					125			
Phe	Tyr	Gly	Cys	Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp
	130					135					140				
Cys	Val	Asp	Lys	Met	Leu	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thr	Asn
145				150					155				160		
Lys	Val	Val	Glu	Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg
			165					170					175		
Val	Asp	Gln	Lys	Cys	Lys	Ser	Ser	Val	Gln	Ile	Asp	Ser	Thr	Pro	Val
		180						185					190		
Ile	Val	Thr	Ser	Asn	Thr	Asn	Met	Cys	Val	Val	Val	Asp	Gly	Asn	Ser

```

<400> 36
Met Ala Thr Phe Tyr Glu Val Ile Val Arg Val Pro Phe Asp Val Glu
1      5      10      15
Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Asp Trp Val Thr Gly
20      25      30
Gln Ile Trp Glu Leu Pro Pro Glu Ser Asp Leu Asn Leu Thr Leu Val

```


		35					40				45				
Glu	Gln	Pro	Gln	Leu	Thr	Val	Ala	Asp	Arg	Ile	Arg	Arg	Val	Phe	Leu
	50					55					60				
Tyr	Glu	Trp	Asn	Lys	Phe	Ser	Lys	Gln	Glu	Ser	Lys	Phe	Phe	Val	Gln
65					70					75					80
Phe	Glu	Lys	Gly	Ser	Glu	Tyr	Phe	His	Leu	His	Thr	Leu	Val	Glu	Thr
			85						90					95	
Ser	Gly	Ile	Ser	Ser	Met	Val	Leu	Gly	Arg	Tyr	Val	Ser	Gln	Ile	Arg
			100					105					110		
Ala	Gln	Leu	Val	Lys	Val	Val	Phe	Gln	Gly	Ile	Glu	Pro	Gln	Ile	Asn
		115					120					125			
Asp	Trp	Val	Ala	Ile	Thr	Lys	Val	Lys	Lys	Gly	Gly	Ala	Asn	Lys	Val
	130					135					140				
Val	Asp	Ser	Gly	Tyr	Ile	Pro	Ala	Tyr	Leu	Leu	Pro	Lys	Val	Gln	Pro
145					150					155					160
Glu	Leu	Gln	Trp	Ala	Trp	Thr	Asn	Leu	Asp	Glu	Tyr	Lys	Leu	Ala	Ala
			165						170					175	
Leu	Asn	Leu	Glu	Glu	Arg	Lys	Arg	Leu	Val	Ala	Gln	Phe	Leu	Ala	Glu
			180					185					190		
Ser	Ser	Gln	Arg	Ser	Gln	Glu	Ala	Ala	Ser	Gln	Arg	Glu	Phe	Ser	Ala
		195					200					205			
Asp	Pro	Val	Ile	Lys	Ser	Lys	Thr	Ser	Gln	Lys	Tyr	Met	Ala	Leu	Val
	210					215					220				
Asn	Trp	Leu	Val	Glu	His	Gly	Ile	Thr	Ser	Glu	Lys	Gln	Trp	Ile	Gln
225					230					235					240
Glu	Asn	Gln	Glu	Ser	Tyr	Leu	Ser	Phe	Asn	Ser	Thr	Gly	Asn	Ser	Arg
			245						250					255	
Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Thr	Lys	Ile	Met	Ser	Leu
			260					265					270		
Thr	Lys	Ser	Ala	Val	Asp	Tyr	Leu	Val	Gly	Ser	Ser	Val	Pro	Glu	Asp
		275					280					285			
Ile	Ser	Lys	Asn	Arg	Ile	Trp	Gln	Ile	Phe	Glu	Met	Asn	Gly	Tyr	Asp
	290					295					300				
Pro	Ala	Tyr	Ala	Gly	Ser	Ile	Leu	Tyr	Gly	Trp	Cys	Gln	Arg	Ser	Phe
305					310					315					320
Asn	Lys	Arg	Asn	Thr	Val	Trp	Leu	Tyr	Gly	Pro	Ala	Thr	Thr	Gly	Lys
			325						330					335	
Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Thr	Val	Pro	Phe	Tyr	Gly	Cys
			340					345					350		
Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp	Cys	Val	Asp	Lys
		355					360					365			
Met	Leu	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thr	Asn	Lys	Val	Val	Glu
	370					375					380				
Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg	Val	Asp	Gln	Lys
385					390					395					400
Cys	Lys	Ser	Ser	Val	Gln	Ile	Asp	Ser	Thr	Pro	Val	Ile	Val	Thr	Ser
			405						410					415	
Asn	Thr	Asn	Met	Cys	Val	Val	Val	Asp	Gly	Asn	Ser	Thr	Thr	Phe	Glu
			420					425					430		
His	Gln	Gln	Pro	Leu	Glu	Asp	Arg	Met	Phe	Lys	Phe	Glu	Leu	Thr	Lys
		435					440					445			
Arg	Leu	Pro	Pro	Asp	Phe	Gly	Lys	Ile	Thr	Lys	Gln	Glu	Val	Lys	Asp
	450					455					460				
Phe	Phe	Ala	Trp	Ala	Lys	Val	Asn	Gln	Val	Pro	Val	Thr	His	Glu	Phe
465					470					475					480
Lys	Val	Pro	Arg	Glu	Leu	Ala	Gly	Thr	Lys	Gly	Ala	Glu	Lys	Ser	Leu
			485						490					495	
Lys	Arg	Pro	Leu	Gly	Asp	Val	Thr	Asn	Thr	Ser	Tyr	Lys	Ser	Leu	Glu
		500						505					510		
Lys	Arg	Ala	Arg	Leu	Ser	Phe	Val	Pro	Glu	Thr	Pro	Arg	Ser	Ser	Asp
		515					520					525			
Val	Thr	Val	Asp	Pro	Ala	Pro	Leu	Arg	Pro	Leu	Asn	Trp	Asn	Ser	Arg
	530					535					540				

Leu Val Gly Arg Ser Trp
545 550

<210> 37
<211> 1690
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 37
attctttgct ctggactgct agaggaccct cgctgccatg gctaccttct atgaagtcac 60
tgttcgcgct ccatttgacg tggaggaaca tctgcctgga atttctgaca gctttgtgga 120
ctgggtaact ggtcaaattt gggagctgcc tccagagtca gatttaaatt tgactctggt 180
tgaacagcct cagttgacgg tggctgatag aattcgccgc gtgttcctgt acgagtggaa 240
caaattttcc aagcaggagt ccaaattctt tgtgcagttt gaaaagggat ctgaatattt 300
tcatctgcac acgcttgtgg agacctccgg catctcttcc atggctcctc gccgctacgt 360
gagtcagatt cgcgcccagc tggtgaaagt ggtcttccag ggaattgaac cccagatcaa 420
cgactgggtc gccatcacca aggtaaagaa gggcggagcc aataagggtg tggattctgg 480
gtatatccc gcctacctgc tgccgaaggt ccaaccggag cttcagtggg cgtggacaaa 540
cctggacgag tataaattgg ccgccctgaa tctggaggag cgcaaacggc tcgtcgcgca 600
gtttctggca gaatcctcgc agcgcctcga ggaggcggct tcgcagcgtg agttctcggc 660
tgaccgggtc atcaaaagca agacttccca gaaatacatg gcgctcgtca actggctcgt 720
ggagcacggc atcacttccg agaagcagtg gatccaggaa aatcaggaga gctacctctc 780
cttcaactcc accggcaact ctccggagcca gatcaaggcc gcgctcgaca acgcgaccaa 840
aattatgagt ctgacaaaaa gcgcggtgga ctacctcgtg gggagctccg tttccgagga 900
catttcaaaa aacagaatct ggcaaatttt tgagatgaat ggctacgacc cggcctacgc 960
gggatccatc ctctacggct ggtgtcagcg ctcttcaac aagaggaaca ccgtctggct 1020
ctacggaccc gccacgaccg gcaagaccaa catcgcgag gccatcgccc acactgtgcc 1080
cttttacggc tgcgtgaact ggaccaatga aaactttccc tttaatgact gtgtggacaa 1140
aatgctcatt tgggtgggag agggaaagat gaccaacaag gtggttgaat ccgccaaggc 1200
catcctgggg ggctcaaagg tgcgggtcga tcagaaatgt aaatcctctg ttcaaattga 1260
ttctaccctt gtcattgtaa ctccaatac aaacatgtgt gtggtggtgg atgggaattc 1320
cacgaccttt gaacaccagc agccgctgga ggaccgcatg ttcaaatttg aactgactaa 1380
gcggctcccc ccagattttg gcaagattac taagcaggaa gtcaaggact tttttgcttg 1440
ggcaaagggtc aatcagggtc cggtgactca cgagttaaag gttcccaggg aattggcggg 1500
aactaaaggg gcggagaaat ctctaaaacg cccactgggt gacgtcacca atactagcta 1560
taaaagtctg gagaagcggg ccaggctctc atttgttccc gagacgcctc gcagttcaga 1620
cgtgactgtt gatcccgctc ctctgcgacc gctcaattgg aattcaagat tggttggaag 1680
aagttggtga 1690

<210> 38
<211> 145
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 38
ccatcaccaa ggtaaagaag ggcggagcca ataagggtgt ggattctggg tatattcccc 60
cctacctgct gccgaagggt caaccggagc ttcagtgggc gtggacaaac ctggacgagt 120
ataaattggc cgccctgaat ctgga 145

<210> 39
<211> 174
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 39
taagcaggaa gtcaaggact tttttgcttg ggcaaagggtc aatcagggtgc cgggtgactca 60
cgagttttaa gttcccagggt aattggcggg aactaaagggt gcggagaaat ctctaaaacg 120
cccactgggt gacgtcacca atactagcta taaaagtctg gagaagcggg ccag 174

<210> 40
<211> 187
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial sequence; note =
synthetic construct

<400> 40
cactctcaag caaggggggtt ttgtaagcag tgatgtcata atgatgtaat gcttattgtc 60
acgcgatagt taatgattaa cagtcagtgt atgtgtttta tccaatagga agaaagcgcg 120
cgtatgagtt ctgcgagac ttccggggta taaaagaccg agtgaacgag cccgccgcca 180
ttctttg 187

<210> 41
<211> 168
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial sequence; note =
synthetic construct

<400> 41
aaacctcctt gcttgagagt gtggcactct cccccctgtc gcgttcgctc gctcgctggc 60
tcgtttgggg ggggtggcagc tcaaagagct gccagacgac ggccctctgg ccgtcgcccc 120
cccaaacgag ccagcgagcg agcgaacgcg acagggggga gagtgcc 168

<210> 42
<211> 168
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 42
aaacctcctt gcttgagagt gtggcactct cccccctgtc gcgttcgctc gctcgctggc 60
tcgtttgggg gggcgacggc cagagggccg tcgtctgccg gctctttgag ctgccacccc 120
cccaaacgag ccagcgagcg agcgaacgcg acagggggga gagtgcc 168

<210> 43
<211> 8
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 43
cggtgtga 8

<210> 44
 <211> 8
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 44
 cggttgag

8

<210> 45
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 45
 caaacctcc ttgcttgaga g

21

<210> 46
 <211> 4675
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 46
 ttggccactc cctctctgcg cgctcgctcg ctcactgagg ccgggcgacc aaaggctcgcc 60
 cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc gagcgcgagc agagggagtg 120
 gccaaactcca tcactagggg ttcctggagg ggtggagtcg tgacgtgaat tacgtcatag 180
 ggtaggggag gtcctgtatt agaggtcacg tgagtgtttt gcgacatttt gcgacacccat 240
 gtggtcacgc tgggtattta agcccagagt agcacgcagg gtctccattt tgaagcggga 300
 ggtttgaacg cgacgcccgc atgccggggt tttacgagat tgtgattaag gtccccagcg 360
 accttgacgg gcatctgccc ggcatttctg acagctttgt gaactgggtg gccgagaagg 420
 aatgggagtt gccgccagat tctgacatgg atctgaatct gattgagcag gcacccctga 480
 ccgtggccga gaagctgcag cgcgactttc tgacggaatg gcgccgtgtg agtaaggccc 540
 cggaggccct tttctttgtg caatttgaga agggagagag ctacttccac atgcacgtgc 600
 tcgtggaaac caccgggggtg aaatccatgg ttttgggacg tttcctgagt cagattcgcg 660
 aaaaactgat tcagagaatt taccgcggga tcgagccgac tttgccaaac tggttcgcg 720
 tcacaaagac cagaaatggc gccggaggcg ggaacaaggt ggtggatgag tgctacatcc 780
 ccaattactt gtcccccaaa acccagcctg agctccagtg ggcgtggact aatatggaac 840
 agtatattaag cgctgttttg aatctcacgg agcgtaaacg gttggtggcg cagcatctga 900
 cgacgtgtc gcagacgcag gagcagaaca aagagaatca gaatcccaat tctgatgcgc 960
 cgtgatcag atcaaaaact tcagccaggt acatggagct ggtcgggtgg ctcgtggaca 1020
 aggggattac ctcggagaag cagtggatcc aggaggacca ggcctcatal atctccttca 1080
 atgcggcctc caactcgcg tcccaaatca aggctgcctt ggacaatgcg ggaaagatta 1140
 tgagcctgac taaaaccgcc cccgactacc tgggtgggcca gcagcccgtg gaggacattt 1200
 ccagcaatcg gatttataaa attttggaa taaacgggta cgatcccca tatgaggctt 1260
 ccgtctttct gggatgggcc acgaaaaagt tcggcaagag gaacaccatc tggctgtttg 1320
 ggcctgcaac taccgggaag accaacatcg cggaggccat agcccacact gtgcccttct 1380
 acgggtgcgt aaactggacc aatgagaact ttcccttcaa cgactgtgtc gacaagatgg 1440
 tgatctggtg ggaggagggg aagatgaccg ccaaggtcgt ggagtcggcc aaagccattc 1500
 tcggaggaag caaggtgcgc gtggaccaga aatgcaagtc ctcggcccag atagaccgga 1560
 ctcccgtgat cgtcacctcc aacaccaaca tgtgcgccgt gattgacggg aactcaacga 1620
 ctttcgaaca ccagcagccg ttgcaagacc ggatgttcaa atttgaactc acccgccgtc 1680
 tggatcatga ctttgggaag gtcaccaagc aggaagtcaa agactttttc cgggtgggcaa 1740

aggatcacgt	ggttgagggtg	gagcatgaat	tctacgtcaa	aaaggggtgga	gccaagaaaa	1800
gacccgcccc	cagtgcgca	gatataagt	agcccaaacg	ggtgcgcgag	tcagttgcgc	1860
agccatcgac	gtcagacgcg	gaagcttcga	tcaactacgc	agacaggtac	caaaacaaat	1920
gttctcgtca	cgtgggcatg	aatctgatgc	tgtttcctcg	cagacaatgc	gagagaatga	1980
atcagaattc	aaatatctgc	ttcactcacg	gacagaaaga	ctgttttagag	tgctttcccg	2040
tgtcagaatc	tcaacccggtt	tctgtcgtca	aaaaggcgta	tcagaaactg	tgctacattc	2100
atcatatcat	gggaaagggtg	ccagacgctt	gcactgcctg	cgatctgggtc	aatgtggatt	2160
tggtgactg	catctttgaa	caataaatga	tttaaatacag	gtatggctgc	cgatgggttat	2220
cttceagatt	ggctcgagga	cactctctct	gaaggaataa	gacagtgggtg	gaagctcaaa	2280
cctggccac	caccaccaa	gcccgcagag	cggcataagg	acgacagcag	gggtcttggtg	2340
cttcctgggt	acaagtacct	cggacccttc	aacggactcg	acaagggaga	gccggtcaac	2400
gaggcagacg	ccgcggccct	cgagcacgta	caaagcctac	gaccggcagc	tcgacagcgg	2460
agacaacccg	tacctcaagt	acaaccacgc	cgacgcggag	tttcaggagc	gccttaaaga	2520
agatacgtct	tttgggggca	acctcggacg	agcagtcttc	caggcgaaaa	agagggttct	2580
tgaacctctg	ggcctggttg	aggaacctgt	taagacggct	ccgggaaaaa	agaggccggt	2640
agagcactct	cctgtggagc	cagactcctc	ctcgggaacc	ggaaaggcgg	gccagcagcc	2700
tgcaagaaaa	agattgaatt	ttggtcagac	tggagacgca	gactcagtag	ctgaccccca	2760
gcctctcggg	cagccaccag	cagccccctc	tggtctggga	actaatacga	tggtacagg	2820
cagtggcgca	ccaatggcag	acaataacga	gggcgccgac	ggagtgggta	attcctccgg	2880
aaattggcat	tgcgattcca	catggatggg	cgacagagtc	atcaccacca	gcaccgaac	2940
ctggggccctg	cccacctaca	acaaccacct	ctacaaacaa	atttccagcc	aatcaggagc	3000
ctcgaacgac	aatcactact	ttggctacag	cacccttggtg	gggtattttg	acttcaacag	3060
attccactgc	cacttttcac	cacgtgactg	gcaaagactc	atcaacaaca	actggggatt	3120
ccgacccaag	agactcaact	tcaagctctt	taacattcaa	gtcaaagagg	tcacgcagaa	3180
tgacggtagc	acgacgattg	ccaataacct	taccagcacg	gttcagggtg	ttactgactc	3240
ggagtaccag	ctcccgtacg	tcctcggctc	ggcgcaccaa	ggatgcctcc	cgccgttccc	3300
agcagacgtc	ttcatgggtg	cacagtatgg	atacctcacc	ctgaacaacg	ggagtccaggc	3360
agtaggacgc	tcttcatttt	actgcctgga	gtactttcct	tctcagatgc	tgcgtagcgg	3420
aaacaacttt	accttcagct	acacttttga	ggacgttcct	ttccacagca	gctacgctca	3480
cagccagagt	ctggaccgtc	tcatgaatcc	tctcatcgac	cagtacctgt	attacttgag	3540
cagaacaaac	actccaagt	gaaccaccac	gcagtcaagg	cttcagtttt	ctcaggccgg	3600
agcgagtgc	attcgggacc	agtctaggaa	ctggcttcct	ggaccctgtt	accgccagca	3660
gcgagtatca	aagacatctg	cggataacaa	caacagtga	tactcgtgga	ctggagctac	3720
caagtaccac	ctcaatggca	gagactctct	ggtgaatccg	gccatggcaa	gccacaagga	3780
cgatgaagaa	aagttttttc	ctcagagcgg	ggttctcatc	tttgggaagc	aaggctcaga	3840
gaaaacaaat	gtgaacattg	aaaaggctcat	gattacagac	gaagaggaaa	tcggaacaac	3900
caatcccgtg	gctacggagc	agtatgggtc	tgtatctacc	aacctccaga	gaggcaacag	3960
acaagcagct	accgcagatg	tcaacacaca	aggcgttcct	ccaggcatgg	tctggcagga	4020
cagagatgtg	taccttcagg	ggcccatctg	ggcaaagatt	ccacacacgg	acggacattt	4080
tcacccctct	cccctcatgg	gtggattcgg	acttaaacac	cctcctccac	agattctcat	4140
caagaacacc	ccggtacctg	cgaatccttc	gaccaccttc	agtgcggcaa	agtttgcttc	4200
cttcatcaca	cagtactcca	cgggacacgg	tcagcgtgga	gatcgagtgg	gagctgcaga	4260
aggaaaacag	caaacgctgg	aatcccga	ttcagtacac	ttccaactac	aacaagtctg	4320
ttaatcgtgg	acttaccgtg	gataactaat	gcgtgtattc	agagcctcgc	cccattggca	4380
ccagatacct	gactcgtaat	ctgtaattgc	ttgttaatca	ataaaccgtt	taattcgttt	4440
cagttgaact	ttggtctctg	cgtatttctt	tcttatctag	tttccatggc	tacgtagata	4500
agtagcatgg	cggtttaatc	attaactaca	aggaaccctt	agtgatggag	ttggccactc	4560
cctctctgcg	cgctcgtcgc	ctcactgagg	ccgggcgacc	aaaggtcgcc	cgacgcccgg	4620
gctttgcccc	ggcggcctca	gtgagcgcg	gagcgcgcg	agagggagtg	gcca	4675

<210> 47

<211> 4694

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 47

gtggcactcc	ccccctgtc	gcgttcgctc	gttcgctggc	tcgattgggg	gggtggcagc	60
tcaaagagct	gccagacgac	ggccctctgg	gccgtcgccc	ccccaatcga	gccagcgaac	120
gagcgaacgc	gacagggggg	ggagtgccac	actctctagc	aaggggggtt	tgtagggtgt	180

gatgtcattg	ttgatgtcat	tatagttgtc	acgcgatagt	taatgattaa	cagtcattgtg	240
atgtgtgtta	tccaatagga	tgaaagcgcg	cgaatgagat	ctcgcgagac	ttccggggta	300
taaaaggggt	gagtgaacga	gcccgcgcgc	attctctgct	ctggactgct	agaggaccct	360
cgctgccatg	gctaccttct	atgaagtcac	tgttcgcggt	ccatttgatg	tggagagca	420
cctgcctgga	atttctgaca	actttgtaga	ctgggtaact	ggtcaaattt	gggagctgcc	480
tcccagagtca	gatttgaatt	tgactctgat	tgagcagcct	cagctgacgg	tggctgacag	540
aattcgccgc	gtgttcctgt	acgagtggaa	caaattttcc	aagcaggaga	gcaaatttctt	600
tgtgcagttt	gaaaagggat	ctgaatat	tcactctgcac	acgctcgtgg	agacctccgg	660
catctcttct	atggctcctg	gccgctacgt	gagtcagatt	cgcgcccagc	tgggtgaagg	720
ggtgttccag	aacattgagc	cgcggtat	cgactgggtc	gccatcacca	aggtaaagaa	780
gggcggagcc	aataaggtgg	tggattctgg	gtatatctcc	gcctacctgc	tgccgaagg	840
ccaaccagag	cttcagtggg	cgtggactaa	cctcgaagag	tataaattgg	ccgccctcaa	900
tctggaggag	cgcaaacggc	tcgtcgtca	gtttcagctt	gagtcctcgc	agcgctcgca	960
agaggcatct	tcccagaggg	acgtttcggc	tgaccgggtc	atcaagagca	agacttccca	1020
gaaatacatg	gcgctggtaa	gctggctggg	ggaacatggc	atcacttccg	agaagcagtg	1080
gattcaggag	aatcaggaga	gctacctgtc	cttcaactcc	acgggaaact	ctcggagcca	1140
gattaaagcc	gcgcttgaca	acgcgtcaaa	aattatgagt	ctgaccaaat	ctgcctcaga	1200
ctatctcgtg	ggacagactg	ttccagagga	catttctgaa	aacagaatct	ggcagatttt	1260
tgatctcaac	ggctacgacc	cggcatacgc	gggctctgtt	ctctacggct	ggtgcactcg	1320
cgcttttga	aagaggaaca	ccgtctggct	gtatggacc	gcgaccaccg	gaaagaccaa	1380
catcgcgga	gccatctctc	acaccgtgcc	cttttatggc	tgtgtgaact	ggactaatga	1440
gaactttccc	tttaatgact	gtgtggaaaa	aatgttgatc	tgggtggagg	agggaaagat	1500
gaccagcaag	gtggtggaac	ccgccaaagg	catcttgggg	gggtctagag	tacgagtggg	1560
tcaaaaatgt	aaatcctctg	tacaagtaga	ctctaccccg	gtgattatca	cctccaatac	1620
taacatgtgt	gtggtgggtg	atgggaactc	cacgaccttt	gaacaccagc	agccgctgga	1680
agaccgcatg	ttcagatttg	aactcatgcg	gcggctcccg	ccagattttg	gcaagattac	1740
caagcaggaa	gtcaaagact	tttttgcttg	ggcaaagggtc	aaccagggtg	cggtgactca	1800
cgagtttatg	gttcccaaga	aagtggcggg	aactgagagg	gcggagactt	ctagaaaacg	1860
cccactggat	gacgtcacca	ataccaacta	taaaagtccg	gagaagcggg	cccggctctc	1920
agttgttctt	gagacgcctc	gcagttcaga	cgtgcctgta	gagcccgtc	ctctgcgacc	1980
tctcaactgg	tcttccagg	atgaatgcag	atgtgactat	catgctaaat	ttgactctgt	2040
aacgggggaa	tgtgacgagt	gtgaatat	gaatcggggc	aaaaatggct	gtatctttca	2100
taatgtctaca	cattgtcaaa	tttgtcacgc	tgttcctcca	tgggaaaagg	aaaatgtgtc	2160
agattttaat	gattttgatg	actgtaataa	agagcagtaa	ataaagttag	tagtcatgtc	2220
ttttgttgac	caccctccag	attggttgga	atcgatcggc	gacggctttc	gtgaatttct	2280
cggccttgag	gcgggtcccc	cgaaacccaa	ggccaatcaa	cagaagcaag	ataacgctcg	2340
aggtcttgtg	cttcttggtg	acaagtatct	tggtcctggg	aacggccttg	ataagggcga	2400
tcctgtcaat	tttgctgacg	aggttgcccc	agagcacgac	ctctcctacc	agaaacagct	2460
tgaggcgggc	gataaccctt	acctcaagta	caaccacgcg	gacgcagagt	ttcaggagaa	2520
actcgcttct	gacacttctt	ttgggggaaa	ccttggggaag	gctgttttcc	aggctaataa	2580
gaggattctc	gaacctcttg	gcctggttga	gacgccggat	aaaacggcgc	ctgcggcaaa	2640
aaagaggcct	ctagagcaga	gtcctcaaga	gccagactcc	tcgagcggag	ttggcaagaa	2700
aggcaaacag	cctgccagaa	agagactcaa	ctttgacgac	gaacctggag	ccggagacgg	2760
gcctccccc	gaaggaccat	cttccggagc	tatgtctact	gagactgaaa	tgctgtcagc	2820
agctggcgga	aatggtggcg	atgcgggaca	agggtccgag	ggagtgggta	atgcctccgg	2880
tgattggcat	tgcgattcca	cttgggtcaga	gagccacgtc	accaccacct	caacccgcac	2940
ctgggtcctg	ccgacctaca	acaaccacct	gtacctgcgg	ctcggctcga	gcaacgccag	3000
cgacaccttc	aacggattct	ccaccccttg	gggatacttt	gactttaacc	gcttccactg	3060
ccacttctcg	ccaagagact	ggcaaaggct	catcaacaac	cactggggac	tgcgcccca	3120
aagcatgcaa	gtccgcatct	tcaacatcca	agttaaggag	gtcacgacgt	ctaacgggga	3180
gacgaccgta	tccaacaacc	tcaccagcac	ggtccagatc	tttgcgga	gcacgtacga	3240
gctcccgtac	gtgatggatg	caggtcagga	gggcagcttg	cctcctttcc	ccaacgacgt	3300
gttcatgggtg	cctcagtacg	ggtactgcgg	actggtaacc	ggaggcagct	ctcaaaacca	3360
gacagacaga	aatgccttct	actgtctgga	gtactttccc	agccagatgc	tgagaaccgg	3420
aaacaacttt	gagatgggtg	acaagtgtga	aaacgtgcc	ttccactcca	tgtacgtctca	3480
cagccagagc	ctggataggc	tgatgaaccc	gctgctggac	cagtacctgt	gggagctcca	3540
gtctaccacc	tctggaggaa	ctctcaacca	gggcaattca	gccaccaact	ttgccaagct	3600
gacaaaaca	aacttttctg	gctaccgcaa	aaactggctc	ccggggccca	tgatgaagca	3660
gcagagattc	tccaagactg	ccagtcaaaa	ctacaagatt	ccccagggaa	gaaacaacag	3720
tctgtccat	tatgagacca	gaactaccct	cgacggaaga	tggagcaatt	ttgccccggg	3780
aacggccatg	gcaaccgcag	ccaacgacgc	caccgacttc	tctcaggccc	agctcatctt	3840
tgcggggccc	aacatcaccg	gcaacaccac	cacagatgcc	aataacctga	tgttcacttc	3900
agaagatgaa	cttagggcca	ccaacccccg	ggacactgac	ctgtttggcc	acctggcaac	3960

caaccagcaa	aacgccacca	ccgttcctac	cgtagacgac	gtggacggag	tcggcgtgta	4020
cccgggaatg	gtgtggcagg	acagagacat	ttactaccaa	gggcccattt	gggccaaaat	4080
tccacacacg	gatggacact	ttcacccgtc	tcctctcatt	ggcggatttg	gactgaaaag	4140
cccgctcca	caaattattca	tcaaaaacac	tcctgtaccc	gccaatcccg	caacgacctt	4200
ctctccggcc	agaatcaaca	gcttcatcac	ccagtacagc	accggacagg	tggctgtcaa	4260
aatagaatgg	gaaatccaga	aggagcggtc	caagagatgg	aaccagagg	tccagttcac	4320
gtccaactac	ggagcacagg	actcgcttct	ctgggctccc	gacaacgccg	gagcctacaa	4380
agagcccagg	gccattggat	cccgatacct	caccaaccac	ctctagccca	attctgttgc	4440
ataccctcaa	taaaccgtgt	attcgtttca	gtaaaatact	gcctcttggtg	gtcattcggc	4500
gtacaacagc	ttacaacaac	aacaaaaccc	ccttgctaga	gagtgtggca	ctcccccccc	4560
tgtcgcgttc	gctcgttcgc	tggctcgatt	gggggggtgg	cagctcaaag	agctgccaga	4620
cgacggccct	ctgggccgtc	gcccccccaa	tcgagccagc	gaacgagcga	acgcgacagg	4680
ggggggagtg	ccac					4694

<210> 48

<211> 1833

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 48

atggctacct	tctatgaagt	cattgttcgc	gttccatttg	atgtggaaga	gcacctgcct	60
ggaatttctg	acaactttgt	agactgggta	actggtcaaa	tttgggagct	gcctcccag	120
tcagatttga	atttgactct	gattgagcag	cctcagctga	cgggtggctga	cagaattcgc	180
cgcggtgtcc	tgtacgagtg	gaacaaattt	tccaagcagg	agagcaaatt	ctttgtgcag	240
tttgaaaagg	gatctgaata	ttttcatctg	cacacgctcg	tggagacctc	cgcatctct	300
tctatggtcc	ttggccgcta	cgtgagtcag	attcgcgccc	agctggtgaa	ggtggtgttc	360
cagaacattg	agccgcggat	taacgactgg	gtcgccatca	ccaaggtaaa	gaaggcgga	420
gccaaataagg	tgggtggattc	tgggtatatt	cccgcctacc	tgtgcccga	ggtccaacca	480
gagcttcagt	gggcgtggac	taacctcgaa	gagtataaat	tggccgccct	caatctggag	540
gagcgcaaac	ggctcgtcgc	tcagtttcag	cttgagtcct	cgcagcgctc	gcaagaggca	600
tcttcccaga	gggacgtttc	ggctgacccg	gtcatcaaga	gcaagacttc	ccagaaatac	660
atggcgctgg	taagctggct	ggtggaacat	ggcatcactt	ccgagaagca	gtggattcag	720
gagaatcagg	agagctacct	gtccttcaac	tccacgggaa	actctcggag	ccagattaaa	780
gccgcgcttg	acaacgcgtc	aaaaattatg	agtctgacca	aatctgcctc	agactatctc	840
gtgggacaga	ctgttccaga	ggacatttct	gaaaacagaa	tctggcagat	ttttgatctc	900
aacggctacg	acccggcata	cgcgggctct	gttctctacg	gctggtgcac	tcgcgccttt	960
ggaaagagga	acaccgtctg	gctgtatgga	cccgcgacca	ccggaaagac	caacatcgcg	1020
gaagccatct	ctcacaccgt	gcccttttat	ggctgtgtga	actggactaa	tgagaacttt	1080
cccttttaatg	actgtgtgga	aaaaatgttg	atctggtggg	aggaggga	gatgaccagc	1140
aaggtggtgg	aaccgcgcaa	ggccatcttg	ggggggtcta	gagtacgagt	ggatcaaaaa	1200
tgtaaatcct	ctgtacaagt	agactctacc	ccggtgatta	tcacctcaa	tactaacatg	1260
tgtgtggtgg	tggatgggaa	ctccacgacc	tttgaacacc	agcagccgct	ggaagaccgc	1320
atgttcagat	ttgaactcat	gcggcggtc	ccgccagatt	ttggcaagat	taccaagcag	1380
gaagtcaaag	acttttttgc	ttgggcaaag	gtcaaccagg	tgccggtgac	tcacgagttt	1440
atggttccca	agaaagtggc	gggaactgag	agggcggaga	cttctagaaa	acgcccactg	1500
gatgacgtca	ccaataccaa	ctataaaaagt	ccggagaagc	gggcccggct	ctcagttgtt	1560
cctgagacgc	ctcgcagttc	agacgtgcct	gtagagcccc	ctcctctgcg	acctctcaac	1620
tggcttcca	ggtatgaatg	cagatgtgac	tatcatgcta	aatttgactc	tgtaacgggg	1680
gaatgtgacg	agtgatgaata	tttgaatcgg	ggcaaaaatg	gctgtatctt	tcataatgct	1740
acacattgtc	aaatttgtca	cgctgttcct	ccatgggaaa	aggaaaatgt	gtcagatttt	1800
aatgattttg	atgactgtaa	taaagagcag	taa			1833

<210> 49

<211> 610

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =

synthetic construct

<400> 49
 Met Ala Thr Phe Tyr Glu Val Ile Val Arg Val Pro Phe Asp Val Glu
 1 5 10 15
 Glu His Leu Pro Gly Ile Ser Asp Asn Phe Val Asp Trp Val Thr Gly
 20 25 30
 Gln Ile Trp Glu Leu Pro Pro Glu Ser Asp Leu Asn Leu Thr Leu Ile
 35 40 45
 Glu Gln Pro Gln Leu Thr Val Ala Asp Arg Ile Arg Arg Val Phe Leu
 50 55 60
 Tyr Glu Trp Asn Lys Phe Ser Lys Gln Glu Ser Lys Phe Phe Val Gln
 65 70 75 80
 Phe Glu Lys Gly Ser Glu Tyr Phe His Leu His Thr Leu Val Glu Thr
 85 90 95
 Ser Gly Ile Ser Ser Met Val Leu Gly Arg Tyr Val Ser Gln Ile Arg
 100 105 110
 Ala Gln Leu Val Lys Val Val Phe Gln Asn Ile Glu Pro Arg Ile Asn
 115 120 125
 Asp Trp Val Ala Ile Thr Lys Val Lys Lys Gly Gly Ala Asn Lys Val
 130 135 140
 Val Asp Ser Gly Tyr Ile Pro Ala Tyr Leu Leu Pro Lys Val Gln Pro
 145 150 155 160
 Glu Leu Gln Trp Ala Trp Thr Asn Leu Glu Tyr Lys Leu Ala Ala
 165 170 175
 Leu Asn Leu Glu Arg Lys Arg Leu Val Ala Gln Phe Gln Leu Glu
 180 185 190
 Ser Ser Gln Arg Ser Gln Glu Ala Ser Ser Gln Arg Asp Val Ser Ala
 195 200 205
 Asp Pro Val Ile Lys Ser Lys Thr Ser Gln Lys Tyr Met Ala Leu Val
 210 215 220
 Ser Trp Leu Val Glu His Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln
 225 230 235 240
 Glu Asn Gln Glu Ser Tyr Leu Ser Phe Asn Ser Thr Gly Asn Ser Arg
 245 250 255
 Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Ser Lys Ile Met Ser Leu
 260 265 270
 Thr Lys Ser Ala Ser Asp Tyr Leu Val Gly Gln Thr Val Pro Glu Asp
 275 280 285
 Ile Ser Glu Asn Arg Ile Trp Gln Ile Phe Asp Leu Asn Gly Tyr Asp
 290 295 300
 Pro Ala Tyr Ala Gly Ser Val Leu Tyr Gly Trp Cys Thr Arg Ala Phe
 305 310 315 320
 Gly Lys Arg Asn Thr Val Trp Leu Tyr Gly Pro Ala Thr Thr Gly Lys
 325 330 335
 Thr Asn Ile Ala Glu Ala Ile Ser His Thr Val Pro Phe Tyr Gly Cys
 340 345 350
 Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp Cys Val Glu Lys
 355 360 365
 Met Leu Ile Trp Trp Glu Glu Gly Lys Met Thr Ser Lys Val Val Glu
 370 375 380
 Pro Ala Lys Ala Ile Leu Gly Gly Ser Arg Val Arg Val Asp Gln Lys
 385 390 395 400
 Cys Lys Ser Ser Val Gln Val Asp Ser Thr Pro Val Ile Ile Thr Ser
 405 410 415
 Asn Thr Asn Met Cys Val Val Val Asp Gly Asn Ser Thr Thr Phe Glu
 420 425 430
 His Gln Gln Pro Leu Glu Asp Arg Met Phe Arg Phe Glu Leu Met Arg
 435 440 445
 Arg Leu Pro Pro Asp Phe Gly Lys Ile Thr Lys Gln Glu Val Lys Asp
 450 455 460
 Phe Phe Ala Trp Ala Lys Val Asn Gln Val Pro Val Thr His Glu Phe
 465 470 475 480

Met Val Pro Lys Lys Val Ala Gly Thr Glu Arg Ala Glu Thr Ser Arg
 485 490 495
 Lys Arg Pro Leu Asp Asp Val Thr Asn Thr Asn Tyr Lys Ser Pro Glu
 500 505 510
 Lys Arg Ala Arg Leu Ser Val Val Pro Glu Thr Pro Arg Ser Ser Asp
 515 520 525
 Val Pro Val Glu Pro Ala Pro Leu Arg Pro Leu Asn Trp Ser Ser Arg
 530 535 540
 Tyr Glu Cys Arg Cys Asp Tyr His Ala Lys Phe Asp Ser Val Thr Gly
 545 550 555 560
 Glu Cys Asp Glu Cys Glu Tyr Leu Asn Arg Gly Lys Asn Gly Cys Ile
 565 570 575
 Phe His Asn Ala Thr His Cys Gln Ile Cys His Ala Val Pro Pro Trp
 580 585 590
 Glu Lys Glu Asn Val Ser Asp Phe Asn Asp Phe Asp Asp Cys Asn Lys
 595 600 605
 Glu Gln
 610

<210> 50
 <211> 1173
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 50
 atggcgctgg taagctggct ggtggaacat ggcatacctt ccgagaagca gtggattcag 60
 gagaatcagg agagctacct gtccttcaac tccacgggaa actctcggag ccagattaaa 120
 gccgcgcttg acaacgcgtc aaaaattatg agtctgacca aatctgcctc agactatctc 180
 gtgggacaga ctgttccaga ggacatttct gaaaacagaa tctggcagat ttttgatctc 240
 aacggctacg acccggcata cgcgggctct gttctctacg gctggtgcac tcgcgccttt 300
 ggaaagagga acaccgtctg gctgtatgga cccgcgacca ccggaagac caacatcgcg 360
 gaagccatct ctacacaccgt gcccttttat ggctgtgtga actggactaa tgagaacttt 420
 ccttttaatg actgtgtgga aaaaatgttg atctggtggg aggagggaaa gatgaccagc 480
 aaggtggtgg aaccgcgcaa ggccatcttg ggggggtcta gactacgagt ggatcaaaaa 540
 tgtaaatacct ctgtacaagt agactctacc ccggtgatta tcacctcaa tactaacatg 600
 tgtgtggtgg tggatgggaa ctccacgacc tttgaacacc agcagccgct ggaagaccgc 660
 atgttcagat ttgaactcat gcggcggtc ccgccagatt ttggcaagat taccaagcag 720
 gaagtcaaag acttttttgc ttgggcaaag gtcaaccagg tgccggtgac tcacgagttt 780
 atggttccca agaaagtggc gggaactgag agggcgagga cttctagaaa acgcccactg 840
 gatgacgtca ccaataccaa ctataaaaagt ccggagaagc gggcccggct ctcagttggt 900
 cctgagacgc ctgcagttc agacgtgcct gtagagcccg ctctctgcg acctctcaac 960
 tggctttcca ggtatgaatg cagatgtgac tatcatgcta aatttgactc tgtaacgggg 1020
 gaatgtgacg agtgtgaata tttgaatcgg ggcaaaaatg gctgtatctt tcataatgct 1080
 acacattgtc aaatttgtca cgctgttcct ccatgggaaa aggaaaatgt gtcagatttt 1140
 aatgattttg atgactgtaa taaagagcag taa 1173

<210> 51
 <211> 390
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 51
 Met Ala Leu Val Ser Trp Leu Val Glu His Gly Ile Thr Ser Glu Lys
 1 5 10 15
 Gln Trp Ile Gln Glu Asn Gln Glu Ser Tyr Leu Ser Phe Asn Ser Thr

20 25 30
Gly Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Ser Lys
35 40 45
Ile Met Ser Leu Thr Lys Ser Ala Ser Asp Tyr Leu Val Gly Gln Thr
50 55 60
Val Pro Glu Asp Ile Ser Glu Asn Arg Ile Trp Gln Ile Phe Asp Leu
65 70 75 80
Asn Gly Tyr Asp Pro Ala Tyr Ala Gly Ser Val Leu Tyr Gly Trp Cys
85 90 95
Thr Arg Ala Phe Gly Lys Arg Asn Thr Val Trp Leu Tyr Gly Pro Ala
100 105 110
Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ser His Thr Val Pro
115 120 125
Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
130 135 140
Cys Val Glu Lys Met Leu Ile Trp Trp Glu Glu Gly Lys Met Thr Ser
145 150 155 160
Lys Val Val Glu Pro Ala Lys Ala Ile Leu Gly Gly Ser Arg Val Arg
165 170 175
Val Asp Gln Lys Cys Lys Ser Ser Val Gln Val Asp Ser Thr Pro Val
180 185 190
Ile Ile Thr Ser Asn Thr Asn Met Cys Val Val Val Asp Gly Asn Ser
195 200 205
Thr Thr Phe Glu His Gln Gln Pro Leu Glu Asp Arg Met Phe Arg Phe
210 215 220
Glu Leu Met Arg Arg Leu Pro Pro Asp Phe Gly Lys Ile Thr Lys Gln
225 230 235 240
Glu Val Lys Asp Phe Phe Ala Trp Ala Lys Val Asn Gln Val Pro Val
245 250 255
Thr His Glu Phe Met Val Pro Lys Lys Val Ala Gly Thr Glu Arg Ala
260 265 270
Glu Thr Ser Arg Lys Arg Pro Leu Asp Asp Val Thr Asn Thr Asn Tyr
275 280 285
Lys Ser Pro Glu Lys Arg Ala Arg Leu Ser Val Val Pro Glu Thr Pro
290 295 300
Arg Ser Ser Asp Val Pro Val Glu Pro Ala Pro Leu Arg Pro Leu Asn
305 310 315 320
Trp Ser Ser Arg Tyr Glu Cys Arg Cys Asp Tyr His Ala Lys Phe Asp
325 330 335
Ser Val Thr Gly Glu Cys Asp Glu Cys Glu Tyr Leu Asn Arg Gly Lys
340 345 350
Asn Gly Cys Ile Phe His Asn Ala Thr His Cys Gln Ile Cys His Ala
355 360 365
Val Pro Pro Trp Glu Lys Glu Asn Val Ser Asp Phe Asn Asp Phe Asp
370 375 380
Asp Cys Asn Lys Glu Gln
385 390

<210> 52
<211> 2211
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial sequence; note =
synthetic construct

<400> 52
atgtcttttg ttgaccaccc tccagattgg ttggaatcga tcggcgacgg ctttcgtgaa 60
tttctcggcc ttgaggcggg tccccgaaa cccaaggcca atcaacagaa gcaagataac 120
gctcgaggtc ttgtgcttcc tgggtacaag tatcttggtc ctgggaacgg ccttgataag 180
ggcgatcctg tcaattttgc tgacgaggtt gcccagagac acgacctctc ctaccagaaa 240
cagcttgagg cgggcgataa cccttacctc aagtacaacc acgcggacgc agagtttcag 300


```

gagaaactcg cttctgacac ttcttttggg ggaaaccttg ggaaggctgt tttccaggct 360
aaaaagagga ttctcgaacc tcttggcctg gttgagacgc cggataaaac ggcgccctgcg 420
gcaaaaaaga ggcctctaga gcagagtcct caagagccag actcctcgag cggagttggc 480
aagaaaggca aacagcctgc cagaaagaga ctcaactttg acgacgaacc tggagccgga 540
gacgggcctc ccccagaagg accatcttcc ggagctatgt ctactgagac tgaatgcgt 600
gcagcagctg gcggaaatgg tggcgatgcg ggacaagggt ccgagggagt gggtaatgcc 660
tccggtgatt ggcattgcga ttccacttgg tcagagagcc acgtcaccac cacctcaacc 720
cgcacctggg tcctgccgac ctacaacaac cacctgtacc tgcggctcgg ctcgagcaac 780
gccagcgaca ccttcaacgg attctccacc ccctggggat actttgactt taaccgcttc 840
cactgccact tctcgccaag agactggcaa aggtcatca acaaccactg gggactgcgc 900
cccaaaagca tgcaagtccg catcttcaac atccaagtta aggaggtcac gacgtctaac 960
ggggagacga ccgtatccaa caacctcacc agcacgggtc agatctttgc ggacagcacg 1020
tacgagctcc cgtacgtgat ggatgcaggt caggagggca gcttgccctc tttccccaac 1080
gacgtgttca tgggtgcctca gtacgggtac tgcggactgg taaccggagg cagctctcaa 1140
aaccagacag acagaaatgc cttctactgt ctggagtact tttccagcca gatgctgaga 1200
accggaaaca actttgagat ggtgtacaag tttgaaaacg tgcccttcca ctccatgtac 1260
gctcacagcc agagcctgga taggctgatg aaccgcgtgc tggaccagta cctgtgggag 1320
ctccagtcta ccacctctgg aggaactctc aaccagggca attcagccac caactttgcc 1380
aagctgacca aaacaaactt ttctggctac cgcaaaaact ggctcccggg gcccatgatg 1440
aagcagcaga gattctccaa gactgccagt caaaaactaca agattcccca gggaagaaac 1500
aacagtctgc tccattatga gaccagaact accctcgacg gaagatggag caattttgcc 1560
ccgggaacgg ccattggcaac cgcagccaac gacgccaccg acttctctca ggcccagctc 1620
atctttgcgg ggccaacat caccggcaac accaccacag atgccaataa cctgatgttc 1680
acttcagaag atgaacttag ggccaccaac ccccgggaca ctgacctgtt tggccacctg 1740
gcaaccaacc agcaaaacgc caccaccgtt cctaccgtag acgacgtgga cggagtcggc 1800
gtgtaccggg gaatggtgtg gcaggacaga gacatttact accaaggggc catttgggac 1860
aaaattccac acacggatgg acactttcac ccgtctcctc tcattggcgg atttggactg 1920
aaaagccgc ctccacaaat attcatcaaa aacactcctg taccgccaac tcccgcaacg 1980
accttctctc cggccagaat caacagcttc atcaccagat acagcaccgg acaggtggct 2040
gtcaaaatag aatgggaaat ccagaaggag cgggtccaaga gatggaaccc agaggtccag 2100
ttcacgtcca actacggagc acaggactcg cttctctggg ctcccgacaa cgccggagcc 2160
tacaaagagc ccagggccat tggatcccga tacctcacca accacctcta g 2211

```

<210> 53

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; note =
synthetic construct

<400> 53

```

Met Ser Phe Val Asp His Pro Pro Asp Trp Leu Glu Ser Ile Gly Asp
1      5      10      15
Gly Phe Arg Glu Phe Leu Gly Leu Glu Ala Gly Pro Pro Lys Pro Lys
20      25      30
Ala Asn Gln Gln Lys Gln Asp Asn Ala Arg Gly Leu Val Leu Pro Gly
35      40      45
Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Asp Pro Val
50      55      60
Asn Phe Ala Asp Glu Val Ala Arg Glu His Asp Leu Ser Tyr Gln Lys
65      70      75      80
Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala Asp
85      90      95
Ala Glu Phe Gln Glu Lys Leu Ala Ser Asp Thr Ser Phe Gly Gly Asn
100     105     110
Leu Gly Lys Ala Val Phe Gln Ala Lys Lys Arg Ile Leu Glu Pro Leu
115     120     125
Gly Leu Val Glu Thr Pro Asp Lys Thr Ala Pro Ala Ala Lys Lys Arg
130     135     140
Pro Leu Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Val Gly
145     150     155     160

```

Lys Lys Gly Lys Gln Pro Ala Arg Lys Arg Leu Asn Phe Asp Asp Glu
 165 170 175
 Pro Gly Ala Gly Asp Gly Pro Pro Pro Glu Gly Pro Ser Ser Gly Ala
 180 185 190
 Met Ser Thr Glu Thr Glu Met Arg Ala Ala Gly Gly Asn Gly Gly
 195 200 205
 Asp Ala Gly Gln Gly Ala Glu Gly Val Gly Asn Ala Ser Gly Asp Trp
 210 215 220
 His Cys Asp Ser Thr Trp Ser Glu Ser His Val Thr Thr Thr Ser Thr
 225 230 235 240
 Arg Thr Trp Val Leu Pro Thr Tyr Asn Asn His Leu Tyr Leu Arg Leu
 245 250 255
 Gly Ser Ser Asn Ala Ser Asp Thr Phe Asn Gly Phe Ser Thr Pro Trp
 260 265 270
 Gly Tyr Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp
 275 280 285
 Trp Gln Arg Leu Ile Asn Asn His Trp Gly Leu Arg Pro Lys Ser Met
 290 295 300
 Gln Val Arg Ile Phe Asn Ile Gln Val Lys Glu Val Thr Thr Ser Asn
 305 310 315 320
 Gly Glu Thr Thr Val Ser Asn Asn Leu Thr Ser Thr Val Gln Ile Phe
 325 330 335
 Ala Asp Ser Thr Tyr Glu Leu Pro Tyr Val Met Asp Ala Gly Gln Glu
 340 345 350
 Gly Ser Leu Pro Pro Phe Pro Asn Asp Val Phe Met Val Pro Gln Tyr
 355 360 365
 Gly Tyr Cys Gly Leu Val Thr Gly Gly Ser Ser Gln Asn Gln Thr Asp
 370 375 380
 Arg Asn Ala Phe Tyr Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg
 385 390 395 400
 Thr Gly Asn Asn Phe Glu Met Val Tyr Lys Phe Glu Asn Val Pro Phe
 405 410 415
 His Ser Met Tyr Ala His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro
 420 425 430
 Leu Leu Asp Gln Tyr Leu Trp Glu Leu Gln Ser Thr Thr Ser Gly Gly
 435 440 445
 Thr Leu Asn Gln Gly Asn Ser Ala Thr Asn Phe Ala Lys Leu Thr Lys
 450 455 460
 Thr Asn Phe Ser Gly Tyr Arg Lys Asn Trp Leu Pro Gly Pro Met Met
 465 470 475 480
 Lys Gln Gln Arg Phe Ser Lys Thr Ala Ser Gln Asn Tyr Lys Ile Pro
 485 490 495
 Gln Gly Arg Asn Asn Ser Leu Leu His Tyr Glu Thr Arg Thr Thr Leu
 500 505 510
 Asp Gly Arg Trp Ser Asn Phe Ala Pro Gly Thr Ala Met Ala Thr Ala
 515 520 525
 Ala Asn Asp Ala Thr Asp Phe Ser Gln Ala Gln Leu Ile Phe Ala Gly
 530 535 540
 Pro Asn Ile Thr Gly Asn Thr Thr Thr Asp Ala Asn Asn Leu Met Phe
 545 550 555 560
 Thr Ser Glu Asp Glu Leu Arg Ala Thr Asn Pro Arg Asp Thr Asp Leu
 565 570 575
 Phe Gly His Leu Ala Thr Asn Gln Gln Asn Ala Thr Thr Val Pro Thr
 580 585 590
 Val Asp Asp Val Asp Gly Val Gly Val Tyr Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Ile Tyr Tyr Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly His Phe His Pro Ser Pro Leu Ile Gly Gly Phe Gly Leu
 625 630 635 640
 Lys Ser Pro Pro Pro Gln Ile Phe Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asn Pro Ala Thr Thr Phe Ser Pro Ala Arg Ile Asn Ser Phe Ile Thr

<400> 55
Thr Ala Pro Ala Ala Lys Lys Arg Pro Leu Glu Gln Ser Pro Gln Glu

50/60

Ile Lys Asn Thr Pro Val Pro Ala Asn Pro Ala Thr Thr Phe Ser Pro
 515 520 525
 Ala Arg Ile Asn Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ala
 530 535 540
 Val Lys Ile Glu Trp Glu Ile Gln Lys Glu Arg Ser Lys Arg Trp Asn
 545 550 555 560
 Pro Glu Val Gln Phe Thr Ser Asn Tyr Gly Ala Gln Asp Ser Leu Leu
 565 570 575
 Trp Ala Pro Asp Asn Ala Gly Ala Tyr Lys Glu Pro Arg Ala Ile Gly
 580 585 590
 Ser Arg Tyr Leu Thr Asn His Leu
 595 600

<210> 56
 <211> 1617
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 56
 atgcgtgcag cagctggcgg aaatggtggc gatgcgggac aaggtgccga gggagtgggt 60
 aatgcctccg gtgattggca ttgcgattcc acttggtcag agagccacgt caccaccacc 120
 tcaaccgcga cctgggtcct gccgacctac aacaaccacc tgtacctgcg gctcggctcg 180
 agcaacgcca gcgacacctt caacggattc tccacccctt ggggatactt tgactttaac 240
 cgcttccact gccacttctc gccaaagagac tggcaaaggc tcatcaacaa ccactgggga 300
 ctgcgcccc aagcatgca agtccgcac ttcaacatcc aagttaagga gggtcacgacg 360
 tctaacgggg agacgaccgt atccaacaac ctccaggca cgggtccagat ctttgcggac 420
 agcacgtacg agctcccgtg cgtgatggat gcaggtcagg agggcagctt gcctcctttc 480
 cccaacgacg tgttcatggt gcctcagtag gggtactgcg gactggtaac cggaggcagc 540
 tctcaaaacc agacagacag aaatgccttc tactgtctgg agtactttcc cagccagatg 600
 ctgagaaccg gaaacaactt tgagatgggt tacaagtgtg aaaacgtgcc cttccactcc 660
 atgtacgtc acagccagag cctggatagg ctgatgaacc cgctgctgga ccagtacctg 720
 tgggagctcc agtctaccac ctctggagga actctcaacc agggcaattc agccaccaac 780
 tttgccaagc tgaccaaaac aaacttttct ggctaccgca aaaactggct cccggggccc 840
 atgatgaagc agcagagatt ctccaagact gccagtcaaa actacaagat tccccaggga 900
 agaaacaaca gtctgctcca ttatgagacc agaactaccc tcgacggaag atggagcaat 960
 tttgccccgg gaacggccat ggcaaccgca gccaacgacg ccaccgactt ctctcaggcc 1020
 cagctcatct ttgcggggcc caacatcacc ggcaacacca ccacagatgc caataacctg 1080
 atgtttactt cagaagatga acttagggcc accaaccctt cggtagacga cgtggacgga 1140
 cacctggcaa ccaaccagca aaacgccacc accgttccta ccgtagacga cgtggacgga 1200
 gtcggcgtgt acccggaat ggtgtggcag gacagagaca ttactacca agggccattt 1260
 tgggccaaaa ttccacacac ggatggacac tttcaccctt ctctctcat tggcggattt 1320
 ggactgaaaa gcccgcctcc acaaatattc atcaaaaaca ctctgtacc cgccaatccc 1380
 gcaacgacct tctctccggc cagaatcaac agcttcatca cccagtacag caccggacag 1440
 gtggctgtca aaatagaatg ggaaatccag aaggagcggg ccaagagatg gaaccagag 1500
 gtccagttca cgtccaacta cggagcacag gactcgcttc tctgggctcc cgacaacgcc 1560
 ggagcctaca aagagcccag ggccattgga tcccgatacc tcaccaacca cctctag 1617

<210> 57
 <211> 538
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 57
 Met Arg Ala Ala Ala Gly Gly Asn Gly Gly Asp Ala Gly Gln Gly Ala
 1 5 10 15

Glu Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr Trp
 Ser Glu Ser His Val Thr Thr Thr Ser Thr Arg Thr Trp Val Leu Pro
 Thr Tyr Asn Asn His Leu Tyr Leu Arg Leu Gly Ser Ser Asn Ala Ser
 Asp Thr Phe Asn Gly Phe Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn
 Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn
 Asn His Trp Gly Leu Arg Pro Lys Ser Met Gln Val Arg Ile Phe Asn
 Ile Gln Val Lys Glu Val Thr Thr Ser Asn Gly Glu Thr Thr Val Ser
 Asn Asn Leu Thr Ser Thr Val Gln Ile Phe Ala Asp Ser Thr Tyr Glu
 Leu Pro Tyr Val Met Asp Ala Gly Gln Glu Gly Ser Leu Pro Pro Phe
 Pro Asn Asp Val Phe Met Val Pro Gln Tyr Gly Tyr Cys Gly Leu Val
 Thr Gly Gly Ser Ser Gln Asn Gln Thr Asp Arg Asn Ala Phe Tyr Cys
 Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Glu
 Met Val Tyr Lys Phe Glu Asn Val Pro Phe His Ser Met Tyr Ala His
 Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Leu Asp Gln Tyr Leu
 Trp Glu Leu Gln Ser Thr Thr Ser Gly Gly Thr Leu Asn Gln Gly Asn
 Ser Ala Thr Asn Phe Ala Lys Leu Thr Lys Thr Asn Phe Ser Gly Tyr
 Arg Lys Asn Trp Leu Pro Gly Pro Met Met Lys Gln Gln Arg Phe Ser
 Lys Thr Ala Ser Gln Asn Tyr Lys Ile Pro Gln Gly Arg Asn Asn Ser
 Leu Leu His Tyr Glu Thr Arg Thr Thr Leu Asp Gly Arg Trp Ser Asn
 Phe Ala Pro Gly Thr Ala Met Ala Thr Ala Ala Asn Asp Ala Thr Asp
 Phe Ser Gln Ala Gln Leu Ile Phe Ala Gly Pro Asn Ile Thr Gly Asn
 Thr Thr Thr Asp Ala Asn Asn Leu Met Phe Thr Ser Glu Asp Glu Leu
 Arg Ala Thr Asn Pro Arg Asp Thr Asp Leu Phe Gly His Leu Ala Thr
 Asn Gln Gln Asn Ala Thr Thr Val Pro Thr Val Asp Asp Val Asp Gly
 Val Gly Val Tyr Pro Gly Met Val Trp Gln Asp Arg Asp Ile Tyr Tyr
 Gln Gly Pro Ile Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His
 Pro Ser Pro Leu Ile Gly Gly Phe Gly Leu Lys Ser Pro Pro Pro Gln
 Ile Phe Ile Lys Asn Thr Pro Val Pro Ala Asn Pro Ala Thr Thr Phe
 Ser Pro Ala Arg Ile Asn Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln
 Val Ala Val Lys Ile Glu Trp Glu Ile Gln Lys Glu Arg Ser Lys Arg
 Trp Asn Pro Glu Val Gln Phe Thr Ser Asn Tyr Gly Ala Gln Asp Ser
 Leu Leu Trp Ala Pro Asp Asn Ala Gly Ala Tyr Lys Glu Pro Arg Ala

Ile Gly Ser Arg Tyr Leu Thr Asn His Leu
 530 515 520 525
 535

<210> 58
 <211> 150
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 58
 gtggcactcc cccccctgtc gcgttcgctc gttcgctggc tcgattgggg ggggtggcagc
 tcaaagagct gccagacgac ggccctctgg gccgtcgccc cccaatcga gccagcgaac
 gagcgaacgc gacagggggg ggagtgccac

<210> 59
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 59
 ctctagcaag ggggttttgt

<210> 60
 <211> 7
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 60
 agtgtgg

<210> 61
 <211> 158
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =
 synthetic construct

<400> 61
 aggtggtgat gtcattgttg atgtcattat agttgtcacg cgatagttaa tgattaacag
 tcatgtgatg tgtgttatcc aataggatga aagcgcgcga atgagatctc gcgagacttc
 cggggtataa aaggggtgag tgaacgagcc cgccgcca

<210> 62
 <211> 112
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; note =

synthetic construct

<400> 62
ggtggattct gggatatatt cgcctacct gctgccgaag gtccaaccag agcttcagtg 60
ggcgtggact aacctcgaag agtataaatt ggccgccctc aatctggagg ag 112

<210> 63
<211> 169
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 63
agtcaaagac ttttttgctt gggcaaaggt caaccaggtg ccggtgactc acgagtttat 60
ggttcccaag aaagtggcgg gaactgagag ggcggagact tctagaaaac gccactgga 120
tgacgtcacc aataccaact ataaaagtcc ggagaagcgg gcccggtc 169

<210> 64
<211> 4721
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 64
ttggccactc cctctatgcg cgctcgctcg ctcggtgggg cctgcggacc aaaggtccgc 60
agacggcaga gctctgctct gccggcccca ccgagcgagc gagcgcgcat agagggagtg 120
gccaaactcca tcaactagggg taccgcgaag cgcctccac gctgccgctg cagcgctgac 180
gtaaatcacg tcatagggga gtggtcctgt attagctgtc acgtgagtg cagcgctgac 240
ttttgcgaca ccacgtggcc atttgaggta tatatggccg agtgagcgag caggatctcc 300
attttgaccg cgaaatttga acgagcagca gccatgccgg gtttctacga gatcgtgatc 360
aaggtgccga gcgacctgga cgagcacctg ccgggcatit ctgactcgtt tgtgaactgg 420
gtggccgaga aggaatggga gctgcccccg gattctgaca tggatctgaa tctgatcgag 480
caggcacccc tgaccgtggc cgagaagctg cagcgcgact tcctgggtcca atggcgccgc 540
gtgagtaagg ccccgagggc cctgttcttt gttcagttcg agaagggcga gagctacttc 600
caccttcacg ttctgggtgga gaccacgggg gtcaagtcca tgggtgctagg ccgcttcctg 660
agtcagattc gggagaagct ggtccagacc atctaccgag gggtcgagcc cacgctgccc 720
aactggttcg cggtagacaa gacgcgtaat ggcgcggcg gggggaacaa ggtgggtggac 780
gagtgctaca tccccaaacta cctcctgccc aagacccagc ccgagctgca gtgggctgag 840
actaacatgg aggagtatat aagcgctgtt ttgaacctgg ccgaacgcaa acggctcgtg 900
gcgacgaccc tgaccacagt cagccagacg caggagcaga acaaggagaa tctgaacccc 960
aattctgacg cgcccgtgat cagggtcaaaa acctccgcgc gctacatgga gctgggtcggg 1020
tggctgggtg accggggcat cacctccgag aagcagtgga tccaggagga ccaggcctcg 1080
tacatctcct tcaacgccgc ctccaactcg cgggtcccaga tcaaggccgc gctggacaat 1140
gccggcaaga tcatggcgct gaccaaattc gcgcccgact acctgggtggg gccctcgctg 1200
cccgcggaca ttaaaaccaa ccgcatctac cgcctcctgg agctgaacgg gtacgatcct 1260
gcctacgccg gctccgtctt tctcggctgg gccagaaaa agttcgggaa gcgcaacacc 1320
atctggctgt ttgggcccgc caccaccggc aagaccaaca ttgcggaagc catcgccac 1380
gccgtgccct tctacggctg cgtcaactgg accaatgaga actttccctt caacgattgc 1440
gtcgacaaga tggatgatct gtgggaggag ggcaagatga cggccaaggt cgtggagtcc 1500
gccaaggcca ttctcggcgg cagcaagggt cgcgtggacc aaaagtgcaa gtcgtccgcc 1560
cagatcgacc ccaccccctg gatcgtcacc tccaacacca acatgtgcgc cgtgattgac 1620
gggaacagca ccaccttcga gcaccagcag ccgttgacgg accggatgtt caaatttgaa 1680
ctcaccgccg gtctggagca cgactttggc aaggtgacga agcaggaagt caaagagttc 1740
ttccgctggg ccagtgatca cgtgaccgag gtggcgcatg agttctacgt cagaaagggc 1800
ggagccagca aaagaccgc ccccgatgac gcggatataa gcgagcccaa gcgggcctgc 1860
ccctcagtcg cggatccatc gacgtcagac gcggaaggag ctccggtgga ctttgccgac 1920
aggtaccaa acaaatgttc tcgtcacgcg ggcattgatt agatgctgtt tccctgcaaa 1980

acgtgcgaga	gaatgaatca	gaatttcaac	atttgcttca	cacacggggt	cagagactgt	2040
ttagagtgtt	tccccggcgt	gtcagaatct	caaccgggtcg	tcagaaaaaa	gacgtatcgg	2100
aaactctgcg	cgattcatca	tctgctgggg	cgggcgcccc	agattgcttg	ctcggcctgc	2160
gacctgggtca	acgtggacct	ggacgactgc	gtttctgagc	aataaatgac	ttaaaccagg	2220
tatggctgcc	gatggttatc	ttccagattg	gctcgaggac	aacctctctg	agggcattcg	2280
cgagtgggtg	gacctgaaac	ctggagcccc	gaaacccaaa	gccaaccagc	aaaagcagga	2340
caacggcccg	ggtctggtgc	ttcctggcta	caagtacctc	ggacccttca	acggactcga	2400
caagggggag	cccgtcaacg	cggcggacgc	agcggccctc	gagcacgaca	aggcctacga	2460
ccagcagctc	aaagcgggtg	acaatccgta	cctgcggtat	aaccacgccg	acgccgagtt	2520
tcaggagcgt	ctgcaagaag	atacgtcatt	tgggggcaac	ctcgggagag	cagtcttcca	2580
ggccaagaag	cggtttctcg	aacctctcgg	tctggttgag	gaaggcgcta	agacggctcc	2640
tgcaaagaag	agaccggtag	agccgtcacc	tcagcgttcc	cccgaactct	ccacgggcat	2700
cggcaagaaa	ggccagcagc	ccgccagaaa	gagactcaat	ttcggtcaga	ctggcgactc	2760
agagtacgtc	cccgaacctc	aacctctcgg	agaacctcca	gcagcgccct	ctagtgtggg	2820
atctggtaca	gtggctgcag	gcggtggcgc	accaatggca	gacaataacg	aaggtgccga	2880
cggagtgggt	aatgcctcag	gaaattggca	ttgcgattcc	acatggctgg	gcgacagagt	2940
cattaccacc	agcacccgaa	cctgggcccc	gcccacctac	aacaaccacc	tctacaagca	3000
aatctccagt	gaaactgcag	gtagtaccaa	cgacaacacc	tacttcggct	acagcacccc	3060
ctgggggtat	tttgacttta	acagattcca	ctgccacttc	tcaccacgtg	actggcagcg	3120
actcatcaac	aacaactggg	gattccggcc	caagaagctg	cggttcaagc	tcttcaacat	3180
ccaggtcaag	gaggtcacga	cgaatgacgg	cggtacgacc	atcgctaata	accttaccag	3240
cacgattcag	gtattctcgg	actcgggaata	ccagctgccg	tacgtcctcg	gctctgcgca	3300
ccagggtcgc	ctgcctccgt	tcccggcgga	cgtcttcatg	attcctcagt	acggctacct	3360
gactctcaac	aatggcagtc	agtctgtggg	acgttccttc	ttctactgcc	tggagtactt	3420
cccctctcag	atgctgagaa	cgggcaacaa	ctttgagttc	agctacagct	tcgaggacgt	3480
gcctttccac	agcagctacg	cacacagcca	gagcctggac	cggctgatga	atccccctcat	3540
cgaccagtac	ttgtactacc	tggccagaac	acagagtaac	ccaggaggca	cagctggcaa	3600
tcgggaactg	cagttttacc	aggcggggcc	ttcaactatg	gccgaacaag	ccaagaattg	3660
gttacctgga	ccttgcttcc	ggcaacaaag	agtctccaaa	acgctggatc	aaaacaacaa	3720
cagcaacttt	gcttggactg	gtgccaccaa	atatcacctg	aacggcagaa	actcgttggt	3780
taatcccggc	gtcgccatgg	caactcacia	ggacgacgag	gaccgctttt	tcccatccag	3840
cggagtcctg	attttttgaa	aaactggagc	aactaacaaa	actacattgg	aaaatgtggt	3900
aatgacaaat	gaagaagaaa	ttcgtcctac	taatcctgta	gccacggaag	aatacgggat	3960
agtcagcagc	aacttacaag	cggctaatac	tgcagcccag	acacaagttg	tcaacaacca	4020
gggagcctta	cctggcatgg	tctggcagaa	ccgggacgtg	tacctgcagg	gtcccatctg	4080
ggccaagatt	cctcacacgg	atggcaactt	tcacccgtct	cctttgatgg	gcggcttttg	4140
acttaaacat	ccgcctcctc	agatcctgat	caagaacact	cccgttcccg	ctaatacctcc	4200
ggaggtgttt	actcctgcca	agtttgcttc	gttcatcaca	cagtacagca	ccggacaagt	4260
cagcgtggaa	atcgagtggg	agctgcagaa	ggaaaacagc	aagcgcctgga	acccggagat	4320
tcagtacacc	tccaactttg	aaaagcagac	tgggtgtggac	tttgccgttg	acagccaggg	4380
tgtttactct	gagcctcgcc	ctattggcac	tcgttacctc	acccgtaatc	tgtaattgca	4440
tgtaaatcaa	taaaccggtt	gattcgtttc	agttgaactt	tgggtctcctg	tgcttcttat	4500
cttatcggtt	tccatagcaa	ctgggttacac	attaactgct	tgggtgcgct	tcacgataag	4560
aacactgacg	tcaccgcggt	acccttagtg	atggagtgtg	ccactccctc	tatgcgcgct	4620
cgctcgctcg	gtggggcctg	cggaccaaaag	gtccgcagac	ggcagagctc	tgctctgccg	4680
gccccaccga	gcgagcgagc	gcgcatagag	ggagtggcca	a		4721

<210> 65

<211> 623

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial sequence; note =
synthetic construct

<400> 65

Met	Pro	Gly	Phe	Tyr	Glu	Ile	Val	Ile	Lys	Val	Pro	Ser	Asp	Leu	Asp
1				5				10						15	
Glu	His	Leu	Pro	Gly	Ile	Ser	Asp	Ser	Phe	Val	Asn	Trp	Val	Ala	Glu
		20						25					30		
Lys	Glu	Trp	Glu	Leu	Pro	Pro	Asp	Ser	Asp	Met	Asp	Leu	Asn	Leu	Ile
		35					40					45			

Glu	Gln	Ala	Pro	Leu	Thr	Val	Ala	Glu	Lys	Leu	Gln	Arg	Asp	Phe	Leu
50						55					60				
Val	Gln	Trp	Arg	Arg	Val	Ser	Lys	Ala	Pro	Glu	Ala	Leu	Phe	Phe	Val
65					70					75					80
Gln	Phe	Glu	Lys	Gly	Glu	Ser	Tyr	Phe	His	Leu	His	Val	Leu	Val	Glu
				85					90					95	
Thr	Thr	Gly	Val	Lys	Ser	Met	Val	Leu	Gly	Arg	Phe	Leu	Ser	Gln	Ile
			100					105					110		
Arg	Glu	Lys	Leu	Val	Gln	Thr	Ile	Tyr	Arg	Gly	Val	Glu	Pro	Thr	Leu
		115					120					125			
Pro	Asn	Trp	Phe	Ala	Val	Thr	Lys	Thr	Arg	Asn	Gly	Ala	Gly	Gly	Gly
	130					135					140				
Asn	Lys	Val	Val	Asp	Glu	Cys	Tyr	Ile	Pro	Asn	Tyr	Leu	Leu	Pro	Lys
145					150					155					160
Thr	Gln	Pro	Glu	Leu	Gln	Trp	Ala	Trp	Thr	Asn	Met	Glu	Glu	Tyr	Ile
				165					170					175	
Ser	Ala	Cys	Leu	Asn	Leu	Ala	Glu	Arg	Lys	Arg	Leu	Val	Ala	Gln	His
			180					185					190		
Leu	Thr	His	Val	Ser	Gln	Thr	Gln	Glu	Gln	Asn	Lys	Glu	Asn	Leu	Asn
		195					200					205			
Pro	Asn	Ser	Asp	Ala	Pro	Val	Ile	Arg	Ser	Lys	Thr	Ser	Ala	Arg	Tyr
	210					215					220				
Met	Glu	Leu	Val	Gly	Trp	Leu	Val	Asp	Arg	Gly	Ile	Thr	Ser	Glu	Lys
225					230					235					240
Gln	Trp	Ile	Gln	Glu	Asp	Gln	Ala	Ser	Tyr	Ile	Ser	Phe	Asn	Ala	Ala
				245					250					255	
Ser	Asn	Ser	Arg	Ser	Gln	Ile	Lys	Ala	Ala	Leu	Asp	Asn	Ala	Gly	Lys
			260					265					270		
Ile	Met	Ala	Leu	Thr	Lys	Ser	Ala	Pro	Asp	Tyr	Leu	Val	Gly	Pro	Ser
		275					280					285			
Leu	Pro	Ala	Asp	Ile	Lys	Thr	Asn	Arg	Ile	Tyr	Arg	Ile	Leu	Glu	Leu
	290					295					300				
Asn	Gly	Tyr	Asp	Pro	Ala	Tyr	Ala	Gly	Ser	Val	Phe	Leu	Gly	Trp	Ala
305					310					315					320
Gln	Lys	Lys	Phe	Gly	Lys	Arg	Asn	Thr	Ile	Trp	Leu	Phe	Gly	Pro	Ala
				325					330					335	
Thr	Thr	Gly	Lys	Thr	Asn	Ile	Ala	Glu	Ala	Ile	Ala	His	Ala	Val	Pro
			340					345					350		
Phe	Tyr	Gly	Cys	Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp
		355					360					365			
Cys	Val	Asp	Lys	Met	Val	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thr	Ala
	370					375					380				
Lys	Val	Val	Glu	Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg
385					390					395					400
Val	Asp	Gln	Lys	Cys	Lys	Ser	Ser	Ala	Gln	Ile	Asp	Pro	Thr	Pro	Val
				405					410					415	
Ile	Val	Thr	Ser	Asn	Thr	Asn	Met	Cys	Ala	Val	Ile	Asp	Gly	Asn	Ser
			420					425					430		
Thr	Thr	Phe	Glu	His	Gln	Gln	Pro	Leu	Gln	Asp	Arg	Met	Phe	Lys	Phe
		435					440					445			
Glu	Leu	Thr	Arg	Arg	Leu	Glu	His	Asp	Phe	Gly	Lys	Val	Thr	Lys	Gln
	450					455					460				
Glu	Val	Lys	Glu	Phe	Phe	Arg	Trp	Ala	Ser	Asp	His	Val	Thr	Glu	Val
465					470					475					480
Ala	His	Glu	Phe	Tyr	Val	Arg	Lys	Gly	Gly	Ala	Ser	Lys	Arg	Pro	Ala
				485					490					495	
Pro	Asp	Asp	Ala	Asp	Ile	Ser	Glu	Pro	Lys	Arg	Ala	Cys	Pro	Ser	Val
			500					505					510		
Ala	Asp	Pro	Ser	Thr	Ser	Asp	Ala	Glu	Gly	Ala	Pro	Val	Asp	Phe	Ala
		515					520					525			
Asp	Arg	Tyr	Gln	Asn	Lys	Cys	Ser	Arg	His	Ala	Gly	Met	Ile	Gln	Met
	530					535					540				
Leu	Phe	Pro	Cys	Lys	Thr	Cys	Glu	Arg	Met	Asn	Gln	Asn	Phe	Asn	Ile

545 Cys Phe Thr His Gly Val Arg Asp Cys Leu Glu Cys Phe Pro Gly Val
 550 560
 565 Ser Glu Ser Gln Pro Val Val Arg Lys Lys Thr Tyr Arg Lys Leu Cys
 575
 580 Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala
 590
 595 Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln
 600 605
 610 615 620

<210> 66
 <211> 737
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Description of Artificial sequence; note =
 synthetic construct

<400> 66
 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro
 20 25 30
 Lys Ala Asn Gln Gln Lys Gln Asp Asn Gly Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60
 Val Asn Ala Ala Asp Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80
 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
 100 105 110
 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
 115 120 125
 Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Ala Lys Lys Arg
 130 135 140
 Pro Val Glu Pro Ser Pro Gln Arg Ser Pro Asp Ser Ser Thr Gly Ile
 145 150 155 160
 Gly Lys Lys Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln
 165 170 175
 Thr Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro
 180 185 190
 Pro Ala Ala Pro Ser Ser Val Gly Ser Gly Thr Val Ala Ala Gly Gly
 195 200 205
 Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn
 210 215 220
 Ala Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val
 225 230 235 240
 Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His
 245 250 255
 Leu Tyr Lys Gln Ile Ser Ser Glu Thr Ala Gly Ser Thr Asn Asp Asn
 260 265 270
 Thr Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285
 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300
 Asn Trp Gly Phe Arg Pro Lys Lys Leu Arg Phe Lys Leu Phe Asn Ile
 305 310 315 320
 Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn
 325 330 335
 Asn Leu Thr Ser Thr Ile Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu

Pro Tyr Val 340 Leu Gly Ser Ala His 345 Gln Gly Cys Leu Pro 350 Pro Phe Pro
Ala Asp Val 355 Phe Met Ile Pro 360 Gln Tyr Gly Tyr Leu Thr 365 Leu Asn Asn
Gly 370 Ser Gln Ser Val 375 Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
385 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Glu Phe Ser Tyr Ser
405 Phe Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
420 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ala
435 Arg Thr Gln Ser Asn Pro Gly Gly Thr Ala Gly Asn Arg Glu Leu Gln
450 Phe Tyr Gln Gly Gly Pro Ser Thr Met Ala Glu Gln Ala Lys Asn Trp
465 Leu Pro Gly Pro Cys Phe Arg Gln Gln Arg Val Ser Lys Thr Leu Asp
485 Gln Asn Asn Asn Ser Asn Phe Ala Trp Thr Gly Ala Thr Lys Tyr His
500 Leu Asn Gly Arg Asn Ser Leu Val Asn Pro Gly Val Ala Met Ala Thr
515 His Lys Asp Asp Glu Asp Arg Phe Phe Pro Ser Ser Gly Val Leu Ile
530 Phe Gly Lys Thr Gly Ala Thr Asn Lys Thr Thr Leu Glu Asn Val Leu
545 Met Thr Asn Glu Glu Ile Arg Pro Thr Asn Pro Val Ala Thr Glu
565 Glu Tyr Gly Ile Val Ser Ser Asn Leu Gln Ala Ala Asn Thr Ala Ala
580 Gln Thr Gln Val Val Asn Asn Gln Gly Ala Leu Pro Gly Met Val Trp
595 Gln Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro
610 His Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly
625 Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro
645 Ala Asn Pro Pro Glu Val Phe Thr Pro Ala Lys Phe Ala Ser Phe Ile
660 Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu
675 Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser
690 Asn Phe Glu Lys Gln Thr Gly Val Asp Phe Ala Val Asp Ser Gln Gly
705 Val Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn
725 Leu 730 735

<210> 67
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 67
Gly Ser Ser Asn Ala Ser Asp Thr

1 5

<210> 68
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 68
Thr Thr Ser Gly Gly Thr Leu Asn Gln Gly Asn Ser Ala Thr
1 5 10

<210> 69
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 69
Asn Gly Arg Ala His Ala
1 5

<210> 70
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 70
Ser Ile Gly Tyr Pro Leu Pro
1 5

<210> 71
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 71
Lys Phe Asn Lys Pro Phe Val Phe Leu Ile
1 5 10

<210> 72
<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence; note =
synthetic construct

<400> 72
Asn Ile Ser Leu Asp Asn Pro Leu Glu Asn Pro Ser Ser Leu Phe Asp
1 5 10 15
Leu Val Ala Arg Ile Lys
20

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2005/031837

A. CLASSIFICATION OF SUBJECT MATTER
C12N15/864

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 2005/056807 A (THE GOVERNMENT OF THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE S) 23 June 2005 (2005-06-23) example 4	1-68
P,X	----- GIOVANNI DI PASQUALE, JOHN A. CHIORINI: "AAV transcytosis through barrier epithelia" XTH PARVOVIRUS WORKSHOP PROGRAM, 'Online! 9 September 2004 (2004-09-09), XP002364013 Retrieved from the Internet: URL: http://cme.ufl.edu/conf/parvovirus/program.shtml > 'retrieved on 2006-01-23! page 2 ----- -/--	1-68



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the International search

24 January 2006

Date of mailing of the International search report

16/02/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Guarinos Viñals, E

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2005/031837

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	GIOVANNI DI PASQUALE, JOHN A. CHIORINI: "AAV transcytosis through barrier endothelia and endothelium" 8TH ANNUAL MEETING AMERICAN SOCIETY OF GENE THERAPY, 'Online! 1 June 2005 (2005-06-01), XP002364014 Retrieved from the Internet: URL: http://www.asgt.org/am05/programm/finalprogram.pdf > 'retrieved on 2006-01-23! right-hand column, paragraph 1 -----	1-68
A	WO 01/70276 A (THE GOVERNMENT OF THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE) 27 September 2001 (2001-09-27) example 4 -----	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2005/031837

Box II Observations where certain claims were found unsearchable (Continuation of Item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-68 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of Item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2005/031837

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2005056807	A	23-06-2005	NONE	
WO 0170276	A	27-09-2001	AU 4592401 A US 6855314 B1	03-10-2001 15-02-2005

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.